


**Pfizer**

# FLEA and TICK POWDER

(Contains SEVIN\*)  
For Positive Control of  
**BROWN DOG TICKS, FLEAS, AND LICE**  
For Use on Dogs, Cats,  
and in Kennels.

ACTIVE INGREDIENT:  
Carbaryl (1-naphthyl N-methylcarbamate) . . . . . 5%  
INERT INGREDIENTS . . . . . 95%  
100%

**CAUTION:** Keep out of reach of children.  
SEE BACK PANEL FOR ADDITIONAL CAUTIONS.



**NET WEIGHT: 5 OZ.**

## DIRECTIONS FOR USE

The control of Brown Dog Ticks on dogs, in kennels and other areas is becoming more and more difficult.

PFIZER Flea and Tick Powder has been developed especially to aid in the control of Brown Dog Ticks on dogs and premises. It is also very effective against fleas and lice on dogs and cats.

**FOR CONTROL OF TICKS, FLEAS AND LICE ON DOGS AND CATS:** Dust powder liberally over the animal and rub thoroughly into the skin. Be sure to apply to the feet, between the toes, and on the legs. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed.

**IN KENNELS and DOG HOUSES:** Dust in and around sleeping quarters and other areas where ticks and fleas occur.

**CAUTION: HARMFUL IF SWALLOWED. AVOID BREATHING OF DUST. DO NOT CONTAMINATE FOOD OR WATER. WASH HANDS AFTER USE. KEEP OUT OF REACH OF CHILDREN.**

Avoid contact with animals' eyes. Sensitive cats may exhibit temporary salivation. Avoid treatment of or exposure to kittens, or puppies less than 4 weeks of age. Avoid prolonged or repeated contact with skin.

\* TRADEMARK OF UNION CARBIDE CORP. FOR CARBARYL (1-NAPHTHYL N-METHYLCARBAMATE).

**USDA Reg. No. 1007-77**

MADE IN U.S.A. ?



Distributed by  
**AGRICULTURAL DIVISION**  
NEW YORK, N.Y. 10017

