

**DIRECTIONS FOR USE**

**PEAR**

Begin spray application at 10% bloom at a dosage rate of 50 to 100 gallons of a 200 ppm solution/acre. This would be 8 oz./50 gallons or 1 lb./100 gallons.

Repeat spray applications at 4 to 6 day intervals. This may involve up to 8 to 10 applications.

**Limitations**

Do not apply within 60 days of harvest. Use limited to West Coast.

**PEACH**

Use on a weekly application schedule starting with shuck split at the rate of 150 ppm active ingredient 0.125 lb. (equivalent to 2 oz. oxytetracycline) per hundred gallons spray solution, or equivalent to 11.33 oz. of Agricultural Terramycin® per hundred gallons of water (equivalent to 150 ppm active ingredient).

Apply spray solution to point of run off; rate of 3 gallons per tree on basis of 80 trees per acre (240 gallons spray solution per acre). Gallons of spray per acre may be increased for larger trees. Do not exceed 500 gallons per acre. Use pressure sprayer capable of delivering the spray at least 250 lb. pressure per square inch through a hand held single nozzle gun, or 150 lb. pressure per square inch using a wind blast sprayer. For best results with air blast sprayer do not exceed three miles per hour ground speed or 100 miles per hour spray velocity.

**NOTE**

The spray application schedules are based on a definite biological growth period for peaches, the shuck split. Shuck split stage for peaches varies North to South by states, in individual states and by varieties. Applications are weekly after shuck split stage. This may involve up to 8 applications. Applications to cease three weeks before harvest.

**Limitations**

Do not apply within 3 weeks of harvest. Use limited to New Jersey, Maryland, North Carolina, South Carolina, Georgia, Missouri, Michigan, Illinois and Indiana.

 **Mycoshield**™  
Brand of **Agricultural Terramycin®**  
(oxytetracycline)

**FOR CONTROL OF**  
**FIRE BLIGHT OF PEAR AND BACTERIAL SPOT OF PEACH.**  
Active Ingredient: Oxytetracycline base equivalent\* ..... 17%  
Inert Ingredients: ..... 83%  
**100%**

\*From oxytetracycline calcium complex.  
This material is a non-sterile, suitably denatured antibiotic formulation intended for agricultural use in control of plant diseases. SELLER MAKES NO WARRANTY, EXPRESSED OR IMPLIED, AND NO WARRANTY OF MERCHANTABILITY, WHICH EXTENDS BEYOND THE DESCRIPTION ON THIS LABEL. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONS

LOT NO.

**Net Contents: 25 lbs. (11.36 kg)**

 **CHEMICALS DIVISION**  
PFIZER INC., NEW YORK, N.Y. 10017

To avoid p plastic co
<b>Concentra Desired</b>
ppm*
100 ppm 150 ppm 200 ppm

\*ppm = part per

**CAUTION**  
May cause aller  
Do not breathe  
rubber gloves.

**ENVIRONMEN**  
Keep out of lak.  
equipment or di

**STORAGE AN**  
Storage: Keep t  
when possible.

**Disposal: Do no**

EPA REG. NO. 10  
EPA EST. NO. 10

MADE IN U.S.A  
1



# Mycoshield<sup>TM</sup>

Brand of  
**Agricultural Terramycin<sup>®</sup>**  
(oxytetracycline)

**FOR CONTROL OF**

**FIRE BLIGHT OF PEAR AND BACTERIAL SPOT OF PEACH.**

Active Ingredient: Oxytetracycline base equivalent\* ..... 17%  
Inert ingredients: ..... 83%  
..... 100%

\*From oxytetracycline calcium complex.

This material is a non-sterile, suitably denatured antibiotic formulation intended for agricultural use in control of plant diseases. SELLER MAKES NO WARRANTY EXPRESSED OR IMPLIED, AND NO WARRANTY OF MERCHANTABILITY WHICH EXTENDS BEYOND THE DESCRIPTION ON THIS LABEL. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONS

LOT NO.

**Net Contents: 25 lbs. (11.36 kg)**



**CHEMICALS DIVISION**  
PFIZER INC., NEW YORK, N.Y. 10017

<b>MIXING</b>			
To avoid possible pesticide contamination use only clean metal or plastic containers in preparing all solutions.			
Concentration Desired	Quantity Agricultural Terramycin <sup>®</sup> Per Volume of Water		
	50 gals.	100 gals.	500 gals.
ppm*			
100 ppm	4 oz.	8 oz. (½ lb.)	2½ lbs.
150 ppm	6 oz.	12 oz.	3¾ lbs.
200 ppm	8 oz. (½ lb.)	16 oz. (1 lb.)	5 lbs.

\*ppm - part per million oxytetracycline base in prepared solution.

**CAUTION**

May cause allergic reactions. Avoid contact with skin and eyes. Do not breathe dust or spray mist. Wear dust mask, chemical goggles and rubber gloves. Wash thoroughly after handling.

**ENVIRONMENTAL HAZARD**

Keep out of lakes, ponds and streams. Do not contaminate water by cleaning equipment or disposal of waste.


**STORAGE AND DISPOSAL**

**Storage:** Keep tightly closed. Storage should be at a cool temperature when possible, and with minimum exposure to the atmosphere.

**Disposal:** Do not reuse container for other uses when empty.

EPA REG. NO. 1007-82  
EPA EST. NO. 1007-IN-1

MADE IN U.S.A.  
1



**DIRECTIONS FOR USE**

**PEAR**  
 Begin spray application at 10% bloom at a dosage rate of 50 to 100 gallons of a 200 ppm solution/acre. This would be 8 oz./50 gallons or 1 lb./100 gallons.  
 Repeat spray applications at 4 to 6 day intervals. This may involve up to 8 to 10 applications.

**Limitations**  
 Do not apply within 60 days of harvest.  
 Use limited to West Coast.

**PEACH**  
 Use on a weekly application schedule starting with shuck split at the rate of 150 ppm active ingredient 0.125 lb. (equivalent to 2 oz. oxytetracycline) per hundred gallons spray solution, or equivalent to 11.33 oz. of Agricultural Terramycin\* per hundred gallons of water (equivalent to 150 ppm active ingredient).

Apply spray solution to point of run off; rate of 3 gallons per tree on basis of 80 trees per acre (240 gallons spray solution per acre). Gallons of spray per acre may be increased for larger trees. Do not exceed 500 gallons per acre. Use pressure sprayer capable of delivering the spray at least 250 lb. pressure per square inch through a hand held single nozzle gun, or 150 lb. pressure per square inch using a wind blast sprayer. For best results with air blast sprayer do not exceed three miles per hour ground speed or 100 miles per hour spray velocity.

**NOTE**  
 The spray application schedules are based on a definite biological growth period for peaches, the shuck split. Shuck split stage for peaches varies North to South by states, in individual states and by varieties. Applications are weekly after shuck split stage. This may involve up to 8 applications. Applications to cease three weeks before harvest.

**Limitations**  
 Do not apply within 3 weeks of harvest.  
 Use limited to New Jersey, Maryland, North Carolina, South Carolina, Georgia, Missouri, Michigan, Illinois and Indiana.



Brand of  
**Agricultural Terramycin<sup>®</sup>**  
 (oxytetracycline)

**FOR CONTROL OF**  
**FIRE BLIGHT OF PEAR AND BACTERIAL SPOT OF PEACH.**  
 Active Ingredient: Oxytetracycline base equivalent\* ..... 17%  
 Inert Ingredients: ..... 83%  
 100%

\*From oxytetracycline calcium complex.  
 This material is a non-sterile, suitably denatured antibiotic formulation intended for agricultural use in control of plant diseases. SELLER MAKES NO WARRANTY EXPRESSED OR IMPLIED, AND NO WARRANTY OF MERCHANTABILITY, WHICH EXTENDS BEYOND THE DESCRIPTION ON THIS LABEL. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
 SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONS

LOT NO.  
**Net Contents: 25 lbs. (11.36 kg)**

**Pfizer** CHEMICALS DIVISION  
 PFIZER INC., NEW YORK, N.Y. 10017

<b>MIXING</b>			
To avoid possible pesticide contamination use only clean metal or plastic containers in preparing all solutions.			
Concentration Desired	Quantity Agricultural Terramycin <sup>™</sup> Per Volume of Water		
ppm*	50 gals.	100 gals.	500 gals.
100 ppm	4 oz.	8 oz. (½ lb.)	2½ lbs.
150 ppm	6 oz.	12 oz.	3¾ lbs.
200 ppm	8 oz. (½ lb.)	16 oz. (1 lb.)	5 lbs.

\*ppm part per million oxytetracycline base in prepared solution.

**CAUTION**  
 May cause allergic reactions. Avoid contact with skin and eyes. Do not breathe dust or spray mist. Wear dust mask, chemical goggles and rubber gloves. Wash thoroughly after handling.

**ENVIRONMENTAL HAZARD**  
 Keep out of lakes, ponds and streams. Do not contaminate water by cleaning equipment or disposal of waste.

**STORAGE AND DISPOSAL**  
**Storage:** Keep tightly closed. Storage should be at a cool temperature when possible, and with minimum exposure to the atmosphere.

**Disposal:** Do not reuse container or inner container when empty.

EPA REG. NO. 1007-82  
 EPA EST. NO. 1007-IN-1

MADE IN U.S.A.  
 1

