

DRAFT

**DIRECTIONS FOR USE**

is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

**ENTRY STATEMENT**

Do not apply this product in such a manner as to directly or through drift expose workers or the general public. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

**WORKER NOTIFICATION:** Written or oral warnings must be given to workers in an area expected to be in a treated area or in an area about to be treated with this product. Oral warnings must include the following information: Inform workers of areas or fields that must not be entered without appropriate protective clothing until sprays have dried. In case of accidental exposure, wash with plenty of water.

If oral warnings are given, written warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: **CAUTION:** Area treated with MYCOSHIELD on \_\_\_\_\_ (name of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, wash with plenty of water. For further information call \_\_\_\_\_ (telephone number).

**PEAR**  
Apply spray application at 10% bloom at a dosage rate of 50 to 100 gallons of a 200 ppm solution per acre. This would be 8 oz/50 gallons or 1 lb/100 gallons. Repeat spray applications at 5 to 6 day intervals. This may involve up to 8 to 10 applications.

**PEACH AND NECTARINE**

Apply a weekly application schedule starting with shuck split at the rate of 150 ppm (equivalent to 1.133 lb (equivalent to 2 oz oxytetracycline) per hundred gallons spray solution, or equivalent to 11.33 oz. of Agricultural Terramycin® per hundred gallons of water (equivalent to 150 ppm active ingredient).

Apply spray solution to point of run off, rate of 3 gallons per tree on basis of 80 trees per acre (240 gallons spray solution per acre). Gallons of spray per acre may be increased for larger trees. Do not exceed 500 gallons per acre. Use pressure sprayer capable of delivering the spray at least 250 lb pressure per square inch through a hand held single nozzle gun, or 150 lb pressure per square inch using a wind blast sprayer. For best results in air blast sprayer do not exceed three miles per hour ground speed or 100 miles per hour spray velocity.

Application schedules are based on a definite biological growth period for peaches, the shuck split. Shuck split stage for peaches varies North to South by states, individual states and by varieties. Applications are weekly after shuck split stage. This may involve up to 8 or 9 applications. Applications to cease three weeks before harvest.

MADE IN U.S.A.



Brand of

**Agricultural Terramycin®**  
(oxytetracycline)

**FOR CONTROL OF FIRE BLIGHT OF PEAR AND BACTERIAL SPOT OF PEACH AND NECTARINE**

Active Ingredient: Oxytetracycline base equivalent 31.5%

Inert Ingredients 68.5%

Total 100.0%

This material is a non-sterile, suitably denatured antibiotic formulation intended for agricultural use in control of plant diseases. Seller makes no warranty, express or implied of merchantability, fitness or otherwise concerning use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONS

LOT NO.

50034-252

**Net Contents: 25 lb (11.34 kg)**



**CHEMICAL DIVISION**  
PFIZER INC., NEW YORK, N.Y. 10017

**CAUTION**

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS**

May cause allergic reactions. Avoid contact with skin and eyes. Do not breathe dust or spray mist. Wear dust mask, chemical goggles and rubber gloves. Wash thoroughly after handling.

**ENVIRONMENTAL HAZARD**

Keep out of lakes, ponds and streams. Do not contaminate water when disposing of equipment wash waters.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Storage: Keep tightly closed. Storage should be at a cool temperature when possible, and with minimum exposure to the atmosphere.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Disposal:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

**MIXING**

To avoid possible pesticide contamination use only clean metal or plastic containers in preparing all solutions.

Concentration Desired	Quantity Agricultural Terramycin® Per Volume of Water		
	50 gals	100 gals	500 gals
150 ppm	6 oz	12 oz	3 1/2 lb
200 ppm	8 oz ( 1/2 lb )	16 oz ( 1 lb )	5 lb

\* ppm = part per million oxytetracycline base in prepared solution.


EPA REG NO 1007-82

EPA EST NO 1007-IN-1

6 88-355A

Active Ingredient: Oxytetracycline Calcium Complex.....31.5%  
(Equivalent to 17% Oxytetracycline)

Inert Ingredients: .....68.5  
100.0%



**BEST AVAILABLE COPY**

**ACCEPTED**  
AUG 27 1992  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended.