



**SEVIN<sup>®</sup>**

**NET WT. 25 KG.**

REGISTERED  
FEB 20, 1973  
1010-42

**FOR  
CONTROL  
OF INSECT  
PESTS**

APR 1970  
feb 26 1973  
100-10

# 85 SPRAYABLE

## CARBARYL INSECTICIDE

ACTIVE INGREDIENT Carbaryl (1-naphthyl methylcarbamate) 85% by wt.  
INERT INGREDIENTS 15% by wt.

### CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

HARMFUL IF INHALED OR SWALLOWED. Avoid Breathing of Dust or Spray.

Do Not Take Internally. Avoid Contact with Skin and Eyes.

Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do Not Use 2-PAM, opiates, or cholinesterase inhibiting drugs.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.

### WARRANTY

1. The manufacturer guarantees and warrants (a) that the active ingredient content and the total net weight are as stated within lawful limits and (b) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all States.
2. The manufacturer further warrants that the material herein is reasonably fit for use under normal conditions as directed hereon. The manufacturer neither makes nor authorizes any agent or representative to make any other warranties of FITNESS OR OF MERCHANTABILITY, guarantee or representation, express or implied concerning this material. This product is sold only on the basis that buyer assumes all risks of use or handling which result in loss or damage and which are beyond manufacturer's control. No claim of any kind, and whether or not based on negligence, shall be greater in amount than the purchase price of the material in respect of which such claim is made. In no event shall manufacturer or seller be liable for special, indirect or consequential damages resulting from the use or handling of this material.
3. No modification of this warranty and disclaimer is authorized, except by specific reference to them in writing by an employee of the manufacturer.

**DO NOT REUSE EMPTY CONTAINER. DESTROY BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.**

**UNION CARBIDE CORPORATION · AGRICULTURAL PRODUCTS · 270 PARK AVE., NEW YORK, N. Y. 10017**

E.P.A. Reg. No. 1016-42

U.S. Pat. Nos. 2,903,478 and 3,009,855

SE and Union Carbide\* are registered trade marks of Union Carbide Corporation

UCC

Made in U.S.A.

# UNION CARBIDE CORPORATION

<p><i>[Faint, illegible text in the leftmost column of the central row]</i></p>	<p><i>[Faint, illegible text in the second column of the central row]</i></p>	<p>UNION CARBIDE <b>SEVIN</b></p> <p>NET WT. 25 KG.</p> <p><b>85 SPRAYABLE</b></p> <p>FOR CONTROL OF INSECT PESTS</p> <p>CAUTION</p> <p>WARRANTY</p>	<p><i>[Faint, illegible text in the rightmost column of the central row]</i></p>

Job no. 13414 Date 1.11.73

COLORS  
GCM1 388BLUE  
90 BLACK  
COLORS SHOWN IN THIS LAYOUT  
ARE NOT AN ACCURATE COLOR MATCH

MASTERS and DIES  
**BYKRAFT**  
16 E. 42nd ST. N.Y. N.Y. 10017  
PHONE: 212 OX 7-4380

CAUTION: Check this miniature carefully for copy, color reference, size, and general layout. Our responsibility ends with the replacement of any incorrect die, or miniature. WE ARE NOT RESPONSIBLE FOR INCORRECTLY PRINTED CARTONS.

SEVIN 55 Sprinkle is a micro-fine granular for broadcast or water and applicable as an insecticide spray in hydraulic spray or mist blower, low-volume ground equipment and sprayer.

**READ SEVIN LABEL USE STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS**

**SPECIAL DIRECTIONS FOR PREPARING LOW-VOLUME SPRAYS**

**ON LOW-VOLUME GROUND EQUIPMENT:** Use 50-mesh or smaller screen in water system. Add about 5% of the required water volume to the spray tank. Check all nozzles that pump and maintain pressure by this operation while slowly adding the correct amount of SEVIN 55 Sprinkle. Add concentrate and diluent to thoroughly mix concentrate in water. Use water equivalent and apply at least 80 liters per hectare of the prepared spray. After spraying, flush tank and hose with clear water.

**FOR AIRBORNE APPLICATIONS:** For spraying from aircraft or other aerial equipment, SEVIN 55 should be used in a suitable hydraulic spray system. The amount of SEVIN 55 should be based on the amount of water and the amount of water to be sprayed per hectare. Consult the SEVIN 55 label for further information.

**PREVENTION OF INSECT RESISTANCE IN LOW-VOLUME EQUIPMENT AND AIRBORNE APPLICATIONS:** Use SEVIN 55 with spray formulation of other pesticides is not fully known. Before preparing such mixtures, consult the manufacturer's label for information. SEVIN 55 may reduce the other pesticide's and the resistance of water. DO NOT USE SEVIN 55 WITH OTHER PESTICIDES OR PRODUCTS. Incompatibility will reduce insect control and may cause substantial reductions in plant injury. Observe all warnings and restrictions on labeling of all products used in mixture.

**GENERAL CAUTIONS**

- 1. Avoid use under highly alkaline conditions. For optimum effect use with alkaline materials such as Bordeaux base lime sulfur and copper-iron fungicides.
- 2. Some phytotoxicity may occur on tender foliage in the presence of low or high humidity of warm-temperate climates following spraying.
- 3. SEVIN does not control spider mites. Other measures are effective with a recommended mixture.


**MAY KILL HONEY BEES IN SUBSTANTIAL NUMBERS**

This product is highly toxic to bees essential to direct treatment or residues on crops. Protective information may be obtained from your representative Agricultural Extension Service.

Do not use when value of bees as pollinators is high. If they must be treated, spray before dawn or late in the evening. Do not spray near hives. Do not spray near beehives. Do not spray near beehives.

**IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION AND LIMITATIONS**

Pre-harvest and grazing use restrictions are listed at the end of the label. For each crop use SEVIN 55 in accordance with the instructions on the label. Application can be made to the day of harvest and application can be made without loss of efficacy if application is made on the day of harvest. SEVIN 55 is used in accordance with other pesticides. Do not use SEVIN 55 in combination with other pesticides. Do not use SEVIN 55 in combination with other pesticides. Do not use SEVIN 55 in combination with other pesticides.



**DIRECTIONS FOR USE**

**IMPORTANT:** The directions on the label are based on tests and field experience. They are not intended to be applied to all crops and conditions. For optimum results, consult the SEVIN 55 label for specific directions for each crop and condition. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**COTTON INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants.

**TREE FRUIT INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per 100 liters of spray solution. Apply when insects are present. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT, VEGETABLE AND ORNAMENTAL:** Use 100 grams for each spray application. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**CITRUS INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per 100 liters of spray solution. Apply when insects are present. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**SMALL FRUIT INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants.

**TREE NUT INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per 100 liters of spray solution. Apply when insects are present. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**FRUIT AND VEGETABLE INSECT CONTROL**

Recommended dosage refer to grams of SEVIN 55 Sprinkle per hectare. Use at least 10 liters of water per hectare and increase to higher volumes on larger plants. Do not use SEVIN 55 in combination with other pesticides unless specifically stated on the label.

**ORCHARD AND FIELD CROPS INSECT CONTROL**

Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**GRASSHOPPERS**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.

**POULTRY INSECT CONTROL**

**CHICKENS, TURKEYS, PHEASANTS AND POULTRY**  
Apply 0.7 to 2.0 kg per acre as necessary to control grasshoppers on the above crops. A 1.4 kg rate is suggested for corn and sorghum. A 1.4 to 2.0 kg rate should be used when grasshoppers are common in the field. Do not apply to crops requiring special care.



**NET WT. 25 KG.**

**85 SPRAYABLE**

**CARBARYL METHOXY**

ACTIVE INGREDIENT: Carbaryl (methyl 1-naphthylcarbamate) 85% by wt.  
NET WT. 25 KG.

**CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

IRRITANT IF SWALLOWED OR INHALED. Avoid Breathing of Dust or Spray.  
Do Not Take Internally. Avoid Contact with Skin and Eyes.  
Wear regular long-sleeved work clothing. Change to clean, softening daily. Wash hands and face before eating, sleep, or after handling.  
NOTE FOR PHYSICIANS: Carbaryl is a moderate to severe irritant. Avoid use in children. Do not use in pregnant or nursing women.  
AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.

**WARRANTY**

1. The manufacturer warrants and agrees to keep the active ingredient content and the total net weight and its stated volume equal to that on the label for the period of one year from the date of manufacture of the product. If the active ingredient content or the total net weight or its stated volume is found to be less than that on the label, the manufacturer will refund the purchase price of the product or replace the product with a new one of equal value.  
2. The manufacturer warrants and agrees to keep the active ingredient content and the total net weight and its stated volume equal to that on the label for the period of one year from the date of manufacture of the product. If the active ingredient content or the total net weight or its stated volume is found to be less than that on the label, the manufacturer will refund the purchase price of the product or replace the product with a new one of equal value.  
3. The manufacturer warrants and agrees to keep the active ingredient content and the total net weight and its stated volume equal to that on the label for the period of one year from the date of manufacture of the product. If the active ingredient content or the total net weight or its stated volume is found to be less than that on the label, the manufacturer will refund the purchase price of the product or replace the product with a new one of equal value.

DO NOT REUSE EMPTY CONTAINER. REUSE BY SPRAYING WITH WATER OR DETERGENT. DO NOT BURN OR INCINERATE.

UNION CARBIDE CORPORATION AGRICULTURAL PRODUCTS 270 PARK AVE. NEW YORK, N.Y. 10017

**ORNAMENTAL INSECT CONTROL**

SEVIN 85 Sprayable is recommended for use on ornamental plants, shrubs and trees. It is effective against a wide range of insects including aphids, beet beetles, caterpillars, and grasshoppers. It is also effective against ticks and mites. Do not use on plants in bloom.

Use 100 grams per 100 liters of water when sprayed on trees, shrubs and plants. Do not use on plants in bloom.

Apple tree	100 g	100 l
Apricot tree	100 g	100 l
Banana tree	100 g	100 l
Blackberry	100 g	100 l
Blueberry	100 g	100 l
Cherry tree	100 g	100 l
Citrus tree	100 g	100 l
Coconut tree	100 g	100 l
Cucumber	100 g	100 l
Fig tree	100 g	100 l
Grape vine	100 g	100 l
Guava tree	100 g	100 l
Jackfruit tree	100 g	100 l
Jambhaya tree	100 g	100 l
Lemon tree	100 g	100 l
Lime tree	100 g	100 l
Mango tree	100 g	100 l
Orange tree	100 g	100 l
Peach tree	100 g	100 l
Pineapple tree	100 g	100 l
Plum tree	100 g	100 l
Quince tree	100 g	100 l
Raspberry tree	100 g	100 l
Strawberry tree	100 g	100 l
Tangerine tree	100 g	100 l
Watermelon tree	100 g	100 l

Use on shrubs and trees and spray on plants. Do not use on plants in bloom.

Apple tree	100 g	100 l
Apricot tree	100 g	100 l
Banana tree	100 g	100 l
Blackberry	100 g	100 l
Blueberry	100 g	100 l
Cherry tree	100 g	100 l
Citrus tree	100 g	100 l
Coconut tree	100 g	100 l
Cucumber	100 g	100 l
Fig tree	100 g	100 l
Grape vine	100 g	100 l
Guava tree	100 g	100 l
Jackfruit tree	100 g	100 l
Jambhaya tree	100 g	100 l
Lemon tree	100 g	100 l
Lime tree	100 g	100 l
Mango tree	100 g	100 l
Orange tree	100 g	100 l
Peach tree	100 g	100 l
Pineapple tree	100 g	100 l
Plum tree	100 g	100 l
Quince tree	100 g	100 l
Raspberry tree	100 g	100 l
Strawberry tree	100 g	100 l
Tangerine tree	100 g	100 l
Watermelon tree	100 g	100 l

**EXPORT MARKS**

NET: 25.715 Lbs. 25.00 Kgs.  
TARE: 0.7 Lbs. 2.00 Kgs.  
GROSS: 26.415 Lbs. 27.00 Kgs.  
DIMENSIONS: 17" x 17" x 18 inches  
45.57 x 45.45 x 45.72 CM