

## READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

## IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food. Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label directions.

If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat.

## DIRECTIONS FOR USE

Broadcast applications may be made with either ground equipment or aircraft. Treatments may be repeated as necessary. Dosages i. m 20 to 40 pounds per acre of Bait will control cutworms, armyworms, crickets, darkling ground beetles, grassnoppers and sowbugs.

Стор	Dosage Limitations to Meet Residue Tolerances	Days Between Final Application and Harvest
Cucumbers, squash, melons, peanuts	20 pounds per acre	None
Alfalfa, peas	30 pounds per acre	None
Cotton	30 pounds per acre	None
Beans, carrots, corn, eggplants, okra, peppers, potatoes, tomatoes and sweet corn	40 pounds per acre	None
Asparagus, strawberries	40 pounds per acre	1 Day
Broccoli. Brussels sprouts, cabbage, cauliflower, head lettuce, garden beets (roots), horseradish, parsnips, radishes, rutabagas, and turnips (roots)	40 pounds per acre	3 Days
Sugar beets, collards, endive (escarole), garden beets (tops), kale, leaf lettuce, parsley, spinach, Swiss chard and turnips (tops)	40 pounds per acre	14 Days

## **DISPOSAL OF CONTAINERS**

DO NOT REUSE EMPTY CONTAINER

Destroy it by burying with waste or burning. Stay away from smoke or fumes.