

LeGear®

USE DIRECTIONS

**FOR CONTROL OF
HORNFLIES, LICE & TICKS
ON HORSES, BEEF CATTLE
SHEEP AND GOATS
FACEFLIES ON BEEF CATTLE
HORNFLIES ON DAIRY COWS
AND
LICE ON SWINE**

ACCEPTED

JUN 06 1975

Under the Federal Insecticide,
Fungicide and Rodenticide Act
recommended, for the pesticide
registered under
EPA Reg. No.

DAIRY AND CATTLE DUST

CAUTION

Keep out of reach of children. Avoid breathing of dust. Avoid contact with skin, wash thoroughly after using. Avoid contamination of feed or food stuffs or feed or water troughs. Do not apply to dairy cattle later than five hours before milking. Do not reuse empty container. Destroy container by burying with waste or burning. Stay away from smoke or fumes while burning container.

Do not apply to animals under one month of age.

ENVIRONMENTAL CAUTION

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

LeGear®
O'NEAL, JONES & FELDMAN, INC.
1304 Ashby Road, St. Louis, Mo. 63132

ACTIVE INGREDIENTS:

- 5% Methoxyfenothrin Technical
- 4% Methoxyfenothrin (O, O dimethyl phosphor) derivative of diethyl mercaptosuccinate

INERT INGREDIENTS:

Equivalent to 4.4% of 2,2-bis(4-methoxyphenyl) 1,1,1-trichloroethane and 0.6% of other fillers and related compounds.

CAUTION. Keep out of reach of children. See side panel for additional cautions.

EPA Est. No. 70117-1
EPA Reg. No. 10119-50

CONTENTS: ONE POUND



FOR DIRECT APPLICATION

DAIRY ANIMALS - HORN FLIES (Also Beef Cattle and Horses) Use 2 heaping tablespoons (2 oz.) per animal for control of Horn Flies. Dust on poll, neck and back. Rub lightly to insure that the material is worked down beneath the hair. If application is made by dust gun, apply against the grain of the hair to force material into hair. Repeat application at 3 week intervals. Do not apply to Dairy Cattle later than five hours before or during milking. Do not contaminate water troughs. Do not apply to lactating goats.

DAIRY CATTLE - LICE, TICKS and HORN FLIES and as an aid for the control of Face Flies. For control of Horn Flies, Lice and Ticks, make liberal application to all parts of the animals body. Rub into hair as application is made. Repeat treatment as necessary to control infestation. As an aid for the control of Face Flies dust the faces of cattle thoroughly. Frequent repeated applications may be necessary.

HOGS - LICE Make liberal application of dust to hogs, pens and litter. One treatment may be sufficient. If not repeat in ten days or thereafter as needed for control.

HORSES - HORN FLIES, LICE and TICKS For control of Horn Flies, Lice and Ticks, make liberal application to all parts of the animals body. Rub into hair as application is made. Repeat treatment as necessary to control infestation.

SHEEP - LICE, KEDS, TICKS and HORN FLIES For control of lice, keds and ticks make thorough and liberal application to entire animal. Treatment just after shearing is preferred. Repeat as necessary. For control of Horn Flies apply 2 tablespoonsful (2 oz.) to the backs and necks of infested animals. Rub lightly. Repeat application, if necessary, no more often than once every three weeks.

GOATS - LICE, KEDS, TICKS and HORN FLIES For control of lice, keds and ticks make thorough and liberal application to entire animal. For control of Horn Flies apply 2 tablespoonsful (2 oz.) to the backs and necks of infested animals. Rub lightly. Repeat application, if necessary, no more often than once every three weeks.

BEFORE USE READ WARNING AND NOTICE STATEMENTS CAREFULLY

LeGear®

USE DIRECTIONS

FOR CONTROL OF
HORNFLIES, LICE & TICKS
ON HORSES, BEEF CATTLE,
SHEEP AND GOATS
FACEFLIES ON BEEF CATTLE
HORNFLIES ON DAIRY COWS
AND
LICE ON SWINE

ACCEPTED

JUN 28 1975

**DAIRY
AND
CATTLE
GUST**



FOR DIRECT APPLICATION

ACTIVE INGREDIENTS

INERT INGREDIENTS



O'NEAL JONES & FELDMAN INC.

CONTENTS: ONE POUND