



1021-46 PM17 1972

PYROCIDE® BOOSTER CONCENTRATE H, EMULSIFIABLE

Effective as a surface spray against exposed stages of confused flour beetles, grain weevils, cadelles, and many other insects infesting grain and food storage plants.

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427

ACCEPTED

TELEPHONE (612) 544-0341

ACTIVE INGREDIENTS:

- Pyrethrins
- *Piperonyl butoxide, technical
- Petroleum distillate

INERT INGREDIENTS:

MAK 1 4 1965
 This is the label for...
 use intended, for the product...
 registered under...
 EPA Reg. No. 1021-46

	1.20%
	12.00%
	78.80%
	8.00%
	<u>100.00%</u>

*Equivalent to 9.60% (butylcarbityl)(6-propylpiperonyl) ether and 2.40% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed. Cover or remove food and feedstuffs during spraying. In food processing plants do not spray while processing is underway. After spraying in food processing plants, etc. wash with potable water all equipment, benches, shelving and surfaces where exposed food will be handled. Do not store, pour or spill near heat or open flame. Keep container closed. Mix well before use or sampling (in drum).

DIRECTIONS FOR USE

BEST AVAILABLE COPY

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Dilute with water. Add some water to the sprayer tank first, then add PYROCIDE® BOOSTER CONCENTRATE H, EMULSIFIABLE, make up to volume with water and agitate the emulsion vigorously until a complete emulsion is obtained. Spray the water emulsion as directed below in grain storage bins, elevator boots, mill walls and floors, etc., paying special attention to cracks, crevices and other insect hiding places.

For making a heavy deposit, dilute one gallon of PYROCIDE® BOOSTER CONCENTRATE H, EMULSIFIABLE with four gallons of water. The resulting water emulsion should be used fresh the same day, and should not be allowed to stand overnight or longer.

To obtain a deposit of about 75 mg. of piperonyl butoxide per sq. ft., use one gallon of the water emulsion per 1,000 sq. ft. of surface. To obtain a deposit of about 100 mg. of piperonyl butoxide per sq. ft. use on and one-third gallons of the water emulsion per 1,000 sq. ft.

If more frequent applications are to be made, for example, once every two weeks, dilute one plus nine. If daily applications are to be used for contact spraying a more dilute emulsion may be used.

Alkaline surfaces such as cement, mortar or calcimined wood or metal may require more frequent applications than neutral surfaces.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place. Keep container closed.

Pesticide Disposal: Pesticide, spray mixture or rinse water that

**It is a violation of Federal law to use this product
in a manner inconsistent with its labeling.**

Dilute with water. Add some water to the sprayer tank first, then add PYROCIDE® BOOSTER CONCENTRATE H, EMULSIFIABLE, make up to volume with water and agitate the emulsion vigorously until a complete emulsion is obtained. Spray the water emulsion as directed below in grain storage bins, elevator boots, mill walls and floors, etc., paying special attention to cracks, crevices and other insect hiding places.

For making a heavy deposit, dilute one gallon of PYROCIDE® BOOSTER CONCENTRATE H, EMULSIFIABLE with four gallons of water. The resulting water emulsion should be used fresh the same day, and should not be allowed to stand overnight or longer.

To obtain a deposit of about 75 mg. of piperonyl butoxide per sq. ft., use one gallon of the water emulsion per 1,000 sq. ft. of surface. To obtain a deposit of about 100 mg. of piperonyl butoxide per sq. ft. use one and one-third gallons of the water emulsion per 1,000 sq. ft.

If more frequent applications are to be made, for example, once every two weeks, dilute one plus nine. If daily applications are to be used for contact spraying a more dilute emulsion may be used.

Alkaline surfaces such as cement, mortar or calcimined wood or metal may require more frequent applications than neutral surfaces.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place. Keep container closed.

Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

Container Disposal: Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and Local procedures.

Net Contents _____

Manufactured by:

BEST AVAILABLE COPY

MC LAUGHLIN GORMLEY KING COMPANY
8810 Tenth Avenue North
Minneapolis MN 55427