

UNION
FUNGIC
FOR CONTROL OF
ED UN
TO ATTACH

ACTIVE INGREDIENTS:

Pyrethrins	.100%
*Piperonyl butoxide, technical	.818%
2-N-octyl bicycloheptene dicarboximide	.182%
Petroleum distillate	98.900%

*Consists of 2,2,4,4-tetrahydro-6H-benzothiazolo[5,4-b]pyridin-6-yl (butylcarbityl)(6-propylpiperonyl) ether and .163% related compounds.
EPA 284 Insecticide Synergist, U.S. Pat. Nos. 2,545,283 and 2,545,287.

DIRECTIONS FOR USE

IMPORTANT READ THESE ENTIRE DIRECTIONS AND CONDITIONS OF SALE, INCLUDING THE WARRANTY AND LIMITATION OF DAMAGES PROVISION BEFORE USING DAIRY CATTLE SPRAY NO. 6141.

For Cattle: To protect dairy animals from stable flies, horn flies, horse flies, houseflies, mosquitoes and gnats, spray daily, using 1 to 2 ounces per animal. Cover the animal with spray, giving particular attention to backs, flanks and legs. Do not soak the skin or wet the hide. For control of lice, spray the animals with 2 ounces of DAIRY CATTLE SPRAY NO. 6141. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more frequently than once every three weeks.

For horses: Spray lightly or sponge horses, using 1 to 2 ounces per adult animal, proportionately less for smaller animals. Give special attention to shanks, legs, and shoulders. Apply until hair is moistened, but do not soak the hide.

For Pigs: Spray lightly where flies congregate. Repeat as needed.

For Barns, Stables, Milking parlors: To kill flies, mosquitoes, and gnats in the barn prior to milking, close all doors and windows and direct the spray toward the ceiling. The barn should remain closed for 10 to 20 minutes. Sweep up and destroy fallen insects. Repeat as necessary. When coarse spray is used, apply directly on walls, stanchions, ceiling, windows and door frames, wherever flies alight or congregate, until surfaces are moistened. Avoid direct contact with feed, food or food utensils. Spray outdoor surfaces also to discourage flies from resting there. Repeat as needed.

CAUTION**KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Avoid contamination of milk and milk handling equipment.

Avoid overspraying which will result in wetting the animal's hides. Do not pour, store or spill near open flame. Keep container tightly closed. Store in a cool place. Do not re-use empty container. Destroy container when empty.

CONDITIONS OF SALE. These Directions for Use, given above, reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. McLaughlin Gormley King Company and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

WARRANTY AND LIMITATION OF DAMAGES. McLaughlin Gormley King Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale; subject to the risks referred to therein. McLaughlin Gormley King Company makes no other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of McLaughlin Gormley King Company is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.