Specimen Label (4)-Modified Food Additive Petition 6H1946 SYNERGIZED PYRETHRIN SPRAY FOR MILLS, FOOD E ANTS AND THE HOME, F-6632

ACTIVE INCREDIENTS:

Pyrethrins

*Piperonyl butoxide, technical

'N-octyl bicycloheptene dicarboximide

Petroleum distillate

Petroleum distillate

*Equivalent to 0.16% (butylcarbityl) (6-propylpiperonyl)

*Equivalent to 0.16% (butylcarbityl) (6-propylpiperonyl)

and 0.04% related compounds.

*MGK 264, Insacticide Synergist.

This spray is especially designed for use in cafes, restaurants, food storage warehouses, flour mills, meat packing plants and the like. This spray is designed for use in fogging equipment, hand sprayers or in pressurized tank sprayers that give coarse, wetting particles for contact sprays.

DIRECTIONS FOR USE

Flies Mosquiicos Chats Small Flying Math

Flies. Mosquitoes. Grats. Small Flying Moths: Close room and discharge mist vigorously in all parts of room, using l_2 to 2 ounces for each 1000 cubic feet of space. Leave room closed for at least ten minutes. Ventilate room thoroughly before re-entry. Sweep up and destroy fallen insects. Repeat as needed.

IN PRESSURIZED TANK SPRAYERS FOR WFTTING CONTACT SPRAYS

Fleas (Infesting Premises): Remove and destroy animals' bedding and spray sleeping quarters freely, forcing the spray into all cracks in the floor and about the room. Repeat when re-infested.

Roaches. Waterbugs: Apply in all cracks and crevices in woodwork, underneath sinks, dark places behind pipes, and in all places where roaches and waterbugs infest, contacting as many insects as possible. Repeat the operation as often as necessary.

Tothes Moths, Carpet Beetles: Spray all articles to be protected, giving particular attenton to folds and seams. Interior surfaces of storage containers such as closets, chests and trunks should be sprayed regularly. The treatment should be repeated every 30 days unless the articles are stored in moth-tight containers.

Grain Weevils, Cadelles, Flour Peetles, Grain Mites (Accessible Stages): Wet surfaces thoroughly and treat all infested surface areas, insect runways and hiding places at the rate of 1 gallon of this material to 1000 square feet of area, or to the point of run-off. Repeat when necessary.

CAUTION

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Do not apply directly to food. When space spraying, cover or remove all food and food handling surfaces. Do not apply space sprays while food processing is underway. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions above. Thoroughly wash with potable water all treated chutes, processing surfaces, etc., that food will come into contact with prior to re-use. In case of impractivability of washing, do not allow food to come in contact with treated surfaces for at least 48 hours.

Remove pets and cover fish bowls before spraying.

This product is toxic to fish. Keep out of lakes, streams or pends. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not re-use empty container. Destroy when empty and discard in a safe place.

Note: This spray when driffed properly as described and applied per the directions and cautions given above, conforms with Food Additive Regulations 12:.1074, 121.1075 and 121.1205 and the food additive tolerances established thereunder.

For Reference Only:

Manufactured by
MC LAUGHLIN GORNLEY KING COMPANY
MINNEAPOLIS, MINN., U.S.A. 55414

EPA Reg. No. 1021-924

Net	Weight		
-----	--------	--	--