

Shite Tenth Avenue N. Minneapors, Minnesota 88427

ACTIVE INGREDIENTS:

Pyrethrins. *Piperonyl butoxide, technical IN-Octyl bicycloheptene dicarboximide Petroleum distillate Mineral Oil

PYROCIDET Fogging Concentrate 7104


For Use in Homes, Pestaurants, Food Processing Plants, Warenouses, and Industrial Installations.

FOR APPLICATION BY TRAINED PERSONNEL ONLY

TELEPHONE (612) 544 0341

3.00%

6.00%



*Equivalent to 4.80% (butylcarbityl)(6-propylp ether and 1.20% related compounds. MGK@ 264, Insecticide Synergist.

> DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General: This Concentrate is designed for use undiluted in mechanical sprayers which give particle size of aerosol size, such as the Micro-Gen Micron Generator. It can be diluted with refined kerosenes or mineral oil which meet Food Additive Regulation 121.1182 and can be applied in fogging equipment that gives larger than aerosol particle size, such as the Microsc¹ or Challenger machines. It can also be diluted with oil specified above and applied as a contact spray to kill crawling insects shown later on this label.

NOTE: This spray when diluted properly as described, and applied per the directions and cautions given above, conforms with Food Additive Regulations 121.1074, 121.1075 and 121.1205 and the food additive tolerances established thereunder.

Kills the Accessible Stages of INDIAN MEAL MOTH, DRIED FRUIT BEETLE, and CONFUSED FLOUR BEETLE in dried fruit products storage areas or processing areas. Close room or warehouse and shut off all ventilating systems. Use a spraying device similar to the Micro-Gen equipment to dispense this product. Do not apply to dried fruit directly. Direct spray into areas of the room frequented by these insects, hitting as many insects as possible. Then spray into the air above the trays at a rate of 1/2 gallon of spray per 50,000 cu. ft. of space. Repeat application every 30 days. Leave room closed for one hour after application. Do not remain in treated areas and ventilate before re-entry. Infested stored dried fruit should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Kills the Accessible Stages of RICE WEEVIL, YELLOW MEALWORM CADELLE, CONFUSED FLOUR BEETLE, SAW-TOUTHED GRAIN BEETLE, ANGOUMOIS GRAIN MOTH, MEDITERRANEAN FLOUR MOTH, and INDIAN MEAL MOTH in food warehouses, storage rooms and similar areas. Close room or warehouse and shut off all ventilating systems. Use a spraying device similar to the Micro-Gen equipment. Direct spray into corners, under pallets and around stacks of packaged goods, hitting as many insects as possible. Then spray at the rate of one fl. oz. of spray per 1,000 cu. ft. of space. Repeat applications as necessary. Leave room closed for one hour after application. Do not remain in treated areas and ventilate prior to re-entry. Infested package goods should be fumigated or treated by other effective means using an approved product intended for this purpose.

To Kill ROACHES, SILVEPFISH, EARWIGS, ANTS and SPIDERS: Apply in Micro-Gen or similar equipment adjusted to coarse spray. Apply the spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes, and in all places where these insects hide. Contact as many insects as possible. Do not allow spray to contact stored food. Repeat

To Kill ROACHES, SILVERFISH, EARWIGS, ANTS and SPIDERS: Apply in Micro-Gen or similar equipment adjusted to coarse spray. Apply the spray into all cracks and crevices in woodwork, wear, walls, floors, underneath sinks, behind pipes, and in all places where these insects hide. Contact as many insects as possible. Do not allow spray to contact stored food. Repeat treatment as necessary for control.

To Kill FLIES, MOSQUITOES and GNATS in open areas, dilute Concentrate with petroleum distillate at the rate of one part Concentrate to two parts of oil and mix well. Use at a time when the temperature is cool (75°F to less) and the wind velocity is approximately 5 mph. With equipment similar to the Micro-Gen equipment, spray at a dosage rate of 15 ounces per minute upwind of the area to be controlled.

If the equipment is truck mounted, keep the speed at approximately 5 mph and treat a swath width of no more than 300 feet.

CAUTION

Harmful if swallowed. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing. Wash hands, arms and face with soap and water after handling and before smoking. Wash all contaminated clothing with soap and water before re-use.

Remove pets, birds and cover fish aquariums before spraying.

is product is toxic to fish. Keep out of any body of water. Do not contaminate water by treaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Mix well in container before each use. DO NOT POUR OR SPILL NEAR OPEN FLAME.

STORAGE AND DISPOSAL

Storage: Store in a cocl, dry place. Keep container closed.

<u>Pesticide Disposal</u>: Pesticide, spray mixture or finse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

<u>Container Disposal</u>: Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and Local procedures.

Net Contents

Manufactured by:

MC LAUGHLIN GORMLEY KING COMPANY 8810 Tenth Avenue North Minneapolis MN 55427

EPA Reg. No. 1021-1198

EPA Est. No. 1021-MN-1