



1021-1302

08/09/1974

1/1



PYROCIDE® FOGGING FORMULA 7207

For Use In Meat, Poultry, Egg,
and Rabbit Processing Plants

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427

This synergized pyrethrins insecticide formulation is especially designed for use in Meat, Poultry, Egg and Rabbit Processing Plants. It may also be used in other indoor facilities where the below-named insects are present.

ACTIVE INGREDIENTS:	100%	
Pyrethrins		2.0%
*Piperonyl butoxide, technical		2.5%
1-N-octyl bicycloheptene dicarboximide		2.5%
Petroleum distillate		8.0%
Mineral Oil		85.0%

*Equivalent to 2.0% (butylcarbityl)(6-propylpiperonyl) ether and 0.5% related compounds.

1-MCK-264, Insecticide Synergist.

DIRECTIONS FOR USE

Because of the small micron size particles emitted from the MICRO-GEN, Mini Leco or other cold aerosol foggers, this dual synergized insecticidal formulation is extremely effective on those insects listed. The small particle size permits greater insect-insecticide contact than previously attainable by other methods of space spraying.

Prior to application, close all doors and windows. Turn off air-conditioning or ventilating systems. Cover or remove all food and food handling surfaces.

Space Spray to Kill Flies, Mosquitoes, Small Flying Moths and Gnats: Using a spraying device similar to the Micro-Gen or Mini Leco, apply 1/2 ounce per 1,000 cu. ft. of room space. The applicator should be centrally located or move around the room as necessary to achieve uniform distribution of the insecticide mist. Vacate premises and leave room closed for one hour. Ventilate room thoroughly prior to re-entry. Repeat as necessary for control.

Contact Spray to Kill Roaches, Silverfish, Waterbugs, Earwigs, Ants, Spiders, and Beetles in Sheltered Situations: With equipment described above, apply the spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Do not allow spray to contact stored food. Repeat dosage as necessary for control. Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-entry.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not apply directly to food. Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc. where exposed food will be handled, with potable water. Use a respirator approved by the U.S. Bureau of Mines.

Remove pets, birds and cover fish aquariums before spraying.

Do not re-use empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

Warm to room temperature and mix well in container before each use. Keep container closed and store in a cool place. DO NOT POUR OR SPILL NEAR OPEN FLAME.

Net Weight _____

Manufactured by

MC LAUGHLIN GORMLEY KING COMPANY
MINNEAPOLIS, MINN. U.S.A. 55427

EPA Reg. 1021-1302

EPA Est. 1021-MN-1