



PYROCIDE® Fogging Formula 7221

Poultry House Fogging Spray

8810 Tenth Avenue N / Minneapolis, Minnesota 55427

TELEPHONE (612) 544-0341

For Use in Poultry Houses and Other Animal Holding Areas Listed Below.

ACTIVE INGREDIENTS:

Pyrethrins	0.60%
*Piperonyl butoxide, technical	3.00%
Petroleum distillate	96.40%
	<u>100.00%</u>

*Equivalent to 2.4% (butylcarbityl)(6-propylpiperonyl) ether and 0.6% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Avoid breathing vapors or spray mist. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration and the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part II. Contains petroleum distillate. If swallowed do not induce vomiting. get immediate medical attention. Avoid contact with the eyes and skin. In case of contact immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. Wash all contaminated clothing with soap and warm water before re-use.

When using in animal housing areas, do not apply directly to food, water or feed supplements. Do not apply directly to animals. Wash teats of dairy animals before milking.

This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Physical Hazards

Do not store near heat or open flame.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place. Keep container closed.

Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

Container Disposal: Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and Local procedures.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General: This ready-to-use formula has been specifically designed for use in ULV aerosol generators that produce a large majority of the particles in the 5-30 micron range. A method of determining particle size is available upon request. This product is designed for use in poultry houses, dairy and beef barns, and swine housing areas.


For Use in Poultry Houses, Dairy and Beef Barns, Swine Housing: To Control houseflies, stable flies, horn flies, mosquitoes, gnats, small flying moths: Cover or remove exposed food, feedstuffs and watering troughs. Close poultry house or animal barn as tightly as possible. Using a suitable ULV cold aerosol generator, direct spray upward and apply until the room is filled with mist. Vacate premises, keep house closed for 10 to 30 minutes, and then ventilate thoroughly prior to re-entry. Repeat application as necessary.

Note: This product will kill the above listed insects in the house or barn at the time of treatment. Being non-residual, it will not kill insects that migrate into the treated premises from adjacent areas. In conjunction with space treatment for control of the above mentioned insects, good sanitation measures should be undertaken whenever practical to remove breeding areas and sources for reinvasion of treated premises.

Net Contents _____

Manufactured by:

MC LAUGHLIN GORMLEY KING COMPANY
8810 Tenth Avenue North
Minneapolis MN 55427



EPA Reg. No. 1021-1318

EPA Est. No. 1021-MN-1