

[1021-1318]

M cLAUGHLIN
G ORMLEY
K ING *Company*

8810 Lytle Avenue N. Minneapolis, Minnesota 55427



MGR® PYROCIDE® Fogging Formula 7221

For Use in Poultry Houses and Other Animal Holding Areas Listed Below.

ACTIVE INGREDIENTS:	100%	
Pyrethrins		.6%
*Piperonyl butoxide, technical		3.0%
Petroleum distillate		96.4%

*Equivalent to 2.4% (butylcarbityl) (6-propylpiperonyl) ether and 0.6% related compounds.

General: This ready-to-use formula has been specifically designed for use in ULV aerosol generators that produce a large majority of the particles in the 5-30 micron range. A method of determining particle size is available upon request. This product is designed for use in poultry houses, dairy and beef barns, and swine housing areas.

DIRECTIONS FOR USE

For Use In Poultry Houses, Dairy and Beef Barns, Swine Housing: To control house flies, stable flies, horn flies, mosquitoes, gnats, small flying moths:

Cover or remove exposed food, feedstuffs and watering troughs. Close poultry house or animal barn as tightly as possible. Using a suitable ULV cold aerosol generator, direct spray upward and apply until the room is filled with mist. Vacate premises, keep house closed for 10 to 30 minutes and then ventilate thoroughly prior to re-entry. Repeat application as necessary.

Note: This product will kill the above listed insects in the house or barn at the time of treatment. Being non-residual, it will not kill insects that migrate into the treated premises from adjacent areas. In conjunction with space treatment for control of the above mentioned insects, good sanitation measures should be undertaken whenever practical to remove breeding areas and sources for reinvasion of treated premises.

CAUTION!

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not apply directly to food. Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing. Wash hands, arms and face with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

When using in animal housing areas, do not apply directly to food, water or feed supplements. Do not apply directly to animals. Wash teats of dairy animals before milking.

This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Do not re-use drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

Mix well in drum before sampling and each use. Keep container closed and store in a cool place. Do not pour, spill, or store near open flame.

Manufactured by

MC LAUGHLIN GORMLEY KING COMPANY
MINNEAPOLIS, MINN., U.S.A. 55427

11/1/74

EPA Reg. 1021-1318

EPA Est. 1021-MN-1