



M. S. Red Squill Mouse Tracking Powder (1-10)

For Indoor Use Only

ACTIVE INGREDIENTS:

Scilliroside glycoside (from red squill) .05%

Stearic acid .30%

INERT INGREDIENTS:

99.65%

Total--100.00%

DIRECTIONS FOR USE

For Control of House Mice Indoors:

Stabilized Red Squill Tracking Powder must be placed in dry areas where the mice will pick up the powder.

First, determine dry areas of highest mouse activity, such as along walls, by crawling mice and burrows, in concealed places, or in spaces between floors and walls. These areas are applied directly on the floor, removed and placed on cardboard boxes or other objects to force mice to travel through constricted areas.

Next, apply the tracking powder in the areas of highest activity by applying 1/2 ounce or 5 level teaspoons (1/4 cup) to cover area approximately 10' x 10'. Apply powder in treated area for at least twenty days. Repeat treatments as necessary. After completion of control program, collect and dispose of dead mice by incineration, and clean up the remaining powder.

Red Squill Mouse Tracking Powder can be used only in areas where mice are active. It is not effective against mice in areas where they are not active. It is not effective against other rodents.

Next, apply the tracking powder in those areas of highest activity by sprinkling powder (1/2 ounce or 5 level teaspoons full) to cover an area approximately 6 inches by 12 inches. Keep powder in treated area for at least twenty days. Repeat treatment when re-infestation occurs. After completion of control program, collect and dispose of dead mice by burying or burning, and clean up remaining powder.

Red Squill Mouse Tracking Powder can be used only in areas where there is no possibility of food contamination or contamination of surfaces that come in direct contact with food. The tracking powder should be placed in locations not accessible to children, pets, wildlife, and domestic animals.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED: Do not inhale powder. If swallowed, promptly induce vomiting with salty water or mustard solution. CALL A PHYSICIAN IMMEDIATELY

NOTE TO PHYSICIAN: Persons poisoned should be treated the same as with an overdose of cardiac glycoside.

Wash hands thoroughly after working with bait concentrate and before eating and smoking.

This powder should be placed in areas inaccessible to children, pets, wildlife and domestic animals.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.

Dispose of empty container by perforation or by crushing and burying in a safe place. Do not re-use empty drum.

Manufactured by

MINIAPOLIS COMPANY, INC. COMPANY
MINNEAPOLIS, MINN., U.S.A. 55427

EPA Reg. No. 1021-1326

EPA Estab. No. 10-1-1326

6/1/75

