



MGK® BIG GAME REPELLENT POWDER - BGR®-P

5810 Tenth Avenue N. Minneapolis, Minnesota 55427

For Repelling Black-tailed deer (mule deer) and Roosevelt elk that browse or uproot Douglas Fir in the Northwest United States.

ACTIVE INGREDIENTS:	
Powdered Inedible Egg Solids	36%
INERT INGREDIENTS:	64%
	100%

ACCEPTED
 SEP 11 1980
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 registered for this pesticide
 under EPA Reg. No. 1021-1420

WARNING
 KEEP OUT OF REACH OF CHILDREN

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Avoid inhalation of dust. Harmful if swallowed. Wash contaminated skin with soap and water. In case of eye contact, flush eyes with plenty of water. Get medical attention.

Environmental Hazards

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place. Keep container closed.

Pesticide Disposal: Pesticide that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water. Consult Federal, State or Local disposal authorities.

25

Weather Conditions: BGR-P can only be applied to seedlings when the foliage is moist - either soon after a heavy dew or rain, preferably during a rain.

Timing of Treatment: To protect young seedlings from winter browsing, apply repellent before onset of browsing, usually November in the Northwest area. To protect against spring browsing of new growth, treat wet seedlings after bud break and before new shoots exceed 1" in length.

Method of Application: Using a suitable duster or shaker, apply about one gram of BGR-P to the terminal whorl of each seedling. If applying BGR-P during a rain, do not let the powder in the applicator become wet, as plugging will result.

For Repelling Black-tailed deer (mule deer) and Roosevelt elk that browse or uproot Douglas Fir in the Northwest United States.

ACTIVE INGREDIENTS: Powdered Inedible Egg Solids 36%
INERT INGREDIENTS: 64%
100%

ACCEPTED
SEP 11 1980
EPA Reg. No. 1021-1420

WARNING
KEEP OUT OF REACH OF CHILDREN

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Avoid inhalation of dust. Harmful if swallowed. Wash contaminated skin with soap and water. In case of eye contact, flush eyes with plenty of water. Get medical attention.

Environmental Hazards

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place. Keep container closed.

Pesticide Disposal: Pesticide that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water. Consult Federal, State or Local disposal authorities.

25

Weather Conditions: BGR-P can only be applied to seedlings when the foliage is moist - either soon after a heavy dew or rain, preferably during a rain.

Timing of Treatment: To protect young seedlings from winter browsing, apply repellent before onset of browsing, usually November in the Northwest area. To protect against spring browsing of new growth, treat wet seedlings after bud break and before new shoots exceed 1" in length.

Method of Application: Using a suitable duster or shaker, apply about one gram of BGR-P to the terminal whorl of each seedling. If applying BGR-P during a rain, do not let the powder in the applicator become wet, as plugging will result.

Net Contents _____

MC LAUGHLIN GORMLEY KING COMPANY
8810 Tenth Avenue North
Minneapolis MN 55427

26

EPA Reg. No. 1021-1420

EPA Est. No. 1021-MN-1