

1021-1420

01/12/1981

1/2



MGK® BIG GAME REPELLENT POWDER - BGR®-P

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427

For Repelling Black-tailed deer (mule deer), White-tailed deer and Roosevelt elk that browse or uproot Conifer Seedlings.

ACTIVE INGREDIENTS:

Powdered Inedible Egg Solids

36%

INERT INGREDIENTS:

64%

100%

WARNING
KEEP OUT OF REACH OF CHILDREN

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Avoid inhalation of dust. Harmful if swallowed. Wash contaminated skin with soap and water. In case of eye contact, flush eyes with plenty of water. Get medical attention.

Environmental Hazards

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place. Keep container closed.

Pesticide Disposal: Pesticide that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water. Consult Federal, State or Local disposal authorities.

Weather Conditions: BGR-P can be applied only to seedlings that are moist during a light rain, after a rain or heavy dew or after light misting of dry plants.

Timing of Treatment: To protect young seedling from winter browsing, apply repellent before onset of browsing. To protect against Spring browsing of new growth, treat wet seedlings after bud break and before new shoots exceed 1" in length.

Method of Application: To Treat plants in the field; Using a suitable shaker duster, apply a light dusting of powder to the plant. To heavy a dusting will cause the repellent to slide to the ground where it will not be an effective deterrent to browsing. If only the terminal whorl is to be treated apply about one gram of BGR to it. If applying BGR-P in the rain do not allow the powder in the applicator to become wet, as plugging will result.

To Treat Plants in the Nursery just before outplanting: Using one bundle of conifer seedling at a time, briefly immerse seedling tops in a bucket of water, then dust wet tops lightly with BGR-P striving to deposit some repellent on each seedling. Proceed with planting as usual.

Net Contents _____

McLaughlin Gormley King Company
8810 10th Avenue North
Minneapolis, Minnesota 55427

EPA Reg No. 1021-1420
EPA Est No. 1021-MN-1

ACCEPTED
JAN 12 1981
10-21-1420