1021-1420



MGK® BIG GAME REPELLENT POWDER - BGR-P

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427

TELEPHONE (612) 544-0341

1/2

For Repelling Black-tailed deer (mule deer), White-tailed deer and Roosevelt elk that browse or uproot Conifer Seedlings. Also repels White-tailed deer from listed ornamentals.

05/01/1981

ACTIVE INGREDIENTS: Powdered Inedible Egg Solids INERT INGREDIENTS:

36% 64% 100%

WARNING KEEP OUT OF REACH OF CHILDREN

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Avoid inhalation of dust. Harmful if swallowed. Wash contaminated skin with soap and water. In case of eye contact, flush eyes with plenty of water. Get medical attention.

Environmental Hazards

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place. Keep container closed.

Pesticide Disposal: Pesticide that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water. Consult Federal, State or Local disposal authorities.

<u>Weather Conditions</u>: BGR-P can be applied only to plants that are moist: during a light rain, after a rain or heavy dew, or after light misting of dry plants.

CONIFERS:

<u>Timing of Treatment</u>: To protect young seedling from winter browsing, apply repellent before onset of browsing. To protect against Spring browsing of new growth, treat wet seedlings after bud break and before new shoots exceed 1" in length.

The at statisted in the same the set of the state of the

Method of Application:

To Treat Plants in the Field: Using a suitable shaker duster, apply a light dusting of powder to the plant. Too heavy a dusting will cause the repellent to slide to the ground where it will not be an effective deterent to browsing. If only the terminal whorl is to be treated, apply approximately one gram of BGR-P to it. If applying BGR-P in the rain, do not allow the powder in the applicator to become wet, as plugging... will result.

To Treat Plants in the Nursery or Field Just Before Out-Planting: Using one bundle of conifer seedlings at a time, briefly immerse seedling tops in a bucket of water, then dust wet tops lightly with BGR-P, striving to deposit some repellent on each seedling. Proceed with planting as usual.

ORNAMENTAL SHRUB PLANTINGS - DORMANT USE ONLY:

To protect dormant Canadian Yew, arborvitae, douglas fir, flowering crab, hemlock, scotch pine and honey suckle, apply a light dusting when the plant is moist, to those parts of the shrub to be protected from browsing. Do not overtreat, since the excess powder will fall from the treated surface to the ground.

Net Contents

Manufactured by:

MC LAUGHLIN GORMLEY KING COMPANY 8810 Tenth Avenue North Minneapolis MN 55427

EPA Reg. No. 1021-1420

EPA Est. No. 1021-MN-1

ACCEPTED MAY 1 1981 Under the Federal Inascirida, Fungicide, and Redenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10-2/-