UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

August 19, 2005

Ms. Julie B. Schlekau, Registration Specialist McLaughlin, Gormley, and King Company 8810 Tenth Avenue North Minneapolis, MN 55427-4319

Subject:

Revise Advisory Statements Per PR Notice 2000-5 Evercide® Permethrin Multi-Purpose Spray 27782

EPA Reg. No. 1021-1742

Your submission dated March 11, 2005

Dear Ms. Schlekau:

We have reviewed the above referenced submission, in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended. The labeling is acceptable with the following comments:

• Under Precautionary Statements, delete the following two statements (i.e., 4th and 5th statements below "CAUTION")- "Avoid contact with eyes or clothing" and "Avoid contact with skin", due to repetition/redundance.

Please make the above stated changes to the product's labeling and submit two copies of the revised final printed label to the Agency. Enclosed, please find a copy of the stamped label for your records. If you have any questions concerning this action, contact Dr. B. A. Akinlosotu at (703) 605-0653.

Sincerely

George T. LaRocca, Product Manager 13 Insecticide Branch

Registration Division (7505C)

enclosure:

	CONCURREN	ICES	
SYMBOL . 7505C			
SURNAME - PLANTER W		1	
DATE - 81905			
			OFFICIAL FILE COPY

느



McLAUGHLIN **GORMLEY** KING COMPANY

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4372 U.S.A. - Telephone (763) 544-0341

[] indicates optional wording.

Controls Named Residential, Industrial, Garden, and Livestock Insects For use on Dogs and on Premises to Control Fleas and Ticks Controls Fleas and Ticks on Dogs for 14 Days Controls Pests for up to Four Weeks Kills Lice and Their Eggs

For use on Bedding, Furniture and Other Inanimate Objects Infested with Lice Not for Use on Humans

Ready-To-Use spray or wipe for use on horses & ponies; dogs and their premises.

ACTIVE INGREDIENT:

Permethrin (*(3-Phenoxyphenyl) methyl (+ or -) cis-trans-3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate) OTHER INGREDIENTS.....

99.75% 100.00%

Cis-trans isomers ratio: Min. 35% (+ or -) cis and Max. 65% (+ or -) trans EVERCIDE® - Registered trademark of McLaughlin Gormley King Company

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID		
IF INHALED:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. 	
IF IN EYES:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 	
IF ON SKIN OR	Take off contaminated clothing.	
CLOTHING:	 Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 	
IF SWALLOWED:	 Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control 	
	center or doctor.	
	Do not give any liquid to the person	
	Do not give anything by mouth to an unconscious person. Per or label with you when calling a poison central center or dector or	

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.

ACCEPTED with COMMENTS In EPA Letter Dated:

AUG 19 2005

Under the Federal Sameticide. Fancicide, and Redenticide Act. as amended, for the pesticide registered under EPA Reg. No. 1021-1742

0.25%

Manufactured by: Mc LAUGHLIN GORMLEY KING COMPANY 8810 Tenth Avenue North Minneapolis, MN 55427

Net Contents

EPA Est. No. 1021-MN-2

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if inhaled or absorbed through skin. Avoid contact with eyes, skin, clothing or breathing spray mist. Causes moderate eye irritation. Avoid contact with eyes or clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Vacate room after treatment and ventilate before reoccupying. Avoid contamination of feed and foodstuffs. Cover or remove fishbowls. This product is not for use on humans.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

For Use In the Non-Food areas of food/feed handling establishments, restaurants, and other areas where food is commercially prepared or processed. Do not use in the edible product areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation.

Nonfood areas are areas such as garbage rooms, layatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

In the home, all food processing surfaces and utensils should be covered. `ring treatment or thoroughly washed before use. Exposed food should be covered or rem. . . .

SHAKE WELL BEFORE USING

Hold container upright and spray from a distance of 12 to 15 inches. Remove birds and cover fish aquariums before spraying.

COCKROACHES, WATERBUGS, PALMETTO BUGS, SILVERFISH, FIREBRATS, CRICKETS, SCORPIONS, SPIDERS, MILLIPEDES, CENTIPEDES, SOWBUGS AND PILLBUGS: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible.

Exposed stages of CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, RED FLOUR BEETLES, RICE WEEVILS, GRANARY WEEVILS, GRAIN MITES: Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Keep area closed for 45 minutes to 1 hour after treatment. Infested stored food products in the home should be disposed of.

CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets.

BEDBUGS: Spray mattresses, particularly around tufts and seams. Allow to dry thoroughly before remaking bed. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Allow all sprayed articles to dry thoroughly before reuse. Do not treat blankets, sheets, pillows, pillow cases or mattress covers.

To control CLOTHES MOTHS and CARPET BEETLES on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunfight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage areas, directing the spray into cracks, joints and crevices. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all sprayed articles to dry thoroughly before use.

TO TREAT LIVESTOCK: To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of STABLE FLIES. HORSE FLIES, DEER FLIES, FACE FLIES, HOUSE FLIES, HORN FLIES, MOSQUITOES, AND GNATS. Spray on each side of animal being careful to spray back, withers and forelegs thoroughly. To protect from FACE FLIES, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.

FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment.

To control BLOOD SUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control POULTRY LICE, spray roosts, walls and nests or cages thoroughly, followed by spraying over the birds with a fine mist.

For control of BEDBUGS AND MITES in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

TO CONTROL FLEAS AND TICKS (Adults and Larvae) [ON DOG PREMISES]: Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

Protects Horses and First from attacks of Horse Flies, Stable Flies, Face Flies, Deer Flies, House Flies, Hom Fliè losquitoes and Gnats. Do not use on foals under 3 months of age.

To Protect Horses and Ponies from attacks of Stable Flies, Horse Flies, Deer Flies, Face Flies. House Flies, Hom Flies, Mosquitoes, and Gnats. Spray on each side being careful to spray back withers and forelegs thoroughly. To protect for Face Flies spray the face and head but do not spray into eyes. Repeat treatment when flies are troublesome. Wipe-On: Remove excess dust and dirt. Spray product on clean cloth and wipe animal, being careful to treat back, withers and forelegs. To protect from face flies, apply to face and head but avoid contact with eyes.

READ ENTIRE LABEL BEFORE EACH USE

FOR PETS: USE ONLY ON DOGS, HORSES AND PONIES

Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant or nursing animals.

To control FLEAS, TICKS AND LICE on dogs: Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to ensure penetrating the coat. Dampen the fur up to the head. Do not spray directly in/on eyes, mouth or genitalia of pets. Repeat application weekly if necessary to control infestations. NOTE: Do not use on puppies less than 12 weeks old.

To kill LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry thoroughly before use. Do not treat blankets, sheets, pillows, pillow cases or mattress covers.

For the control of GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill JAPANESE BEETLES hit by spray, as well as WHITEFLIES, APHIDS, EXPOSED THRIPS, SPIDERS, RED MITES AND LEAFMINERS.

May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatment of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivv, euonymus, fuschia. geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against WHITEFLIES, APHIDS, SPIDER AND RED MITES. EXPOSED THRIPS, LACE BUGS, JAPANESE BEETLES, PAVEMENT ANTS, CLOVER MITES AND ARMY WORMS. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or lawns and grounds.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry area. Do not transport or store below 32°F.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.