

1021-1749

06/06/2000

1/6



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (H7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
1021-1749

Date of Issuance:  
June 6, 2000

Term of Issuance: Conditional

Name of Pesticide Product:  
Evercide Permethrin Dust 2786

NOTICE OF PESTICIDE:

Registration  
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

McLaughlin Gormley King Company  
8810 Tenth Avenue North  
Minneapolis, MN 55427-4372

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A), provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2) Please make the following labeling changes:

- a) Revise the EPA Registration Number to read "EPA Reg. No. 1021-1749"
- b) Please revise your First Aid statement in accordance with PR Notice 2000-3:

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center for treatment advice.

Signature of Approving Official:

Date: June 6, 2000

- c) Under "Cole Crops", please put the stipulation that for Chinese Broccoli your product is for use "on gai Ion, white flowering" varieties and "Not for use in California".
  - d) Please remove aphid and flea beetle from your list of pests controlled under "Tomatoes". These pests are not listed on the cited labels for this vegetable.
  - e) Please add the following under "Asparagus (post harvest)": "Not for use in California".
  - f) Under "Peaches", please add the stipulation "Dust lightly. Apply up to 5 times per season".
  - g) Under "Pears-Dormant" please remove codling moth and green fruitworm from the list of pests controlled. These pests are not listed under this category on your cited label.
  - h) Please delete "Specially Formulated" on the front panel. The same product is marketed for commercial use and the product lists uses other than residential use.
- 3) Your CSF has been approved. Your product is toxicologically similar to EPA Reg. No. 28293-126 and the acute toxicity data for that product can be used in support of this product.
  4. Please submit two (2) copies of your final printed label for before you release the product for shipment. Please refer to the A-79 enclosure for a further description of final printed labeling. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing amended labeling constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

If you have any questions regarding this action, please contact Tracy Keigwin of my team at (703) 305-6605.

Sincerely,

George T. LaRocca  
Product Manager (13)  
Insecticide Branch  
Registration Division (7505C)

**MIGK** McLAUGHLIN  
GORMLEY  
KING COMPANY

EVERCIDE® Permethrin Dust 2786

3810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4372 U.S.A.

TELEPHONE (612) 544-0341

Optional marketing statements:

Specially formulated for residential use  
Contains PERMETHRIN  
KILLS beetles, caterpillars, bugs and many more.  
Ready to Use  
Controls ornamental pests  
control (kill) vegetable pests  
Use on home garden fruits, vegetables and ornamental plants (including roses)  
Use on fruit & nut trees

Use on dogs, cats and pet bedding to control fleas, ticks and lice

For control of Horn Flies and Lice on Beef and Dairy Cattle  
Aids in reducing Face Fly population on Beef and Dairy Cattle  
For control of Lice on Swine  
For control of Northern Fowl Mites on Poultry  
For use in dust bags, shaker can or mechanical dust applicator

ACCEPTED  
with COMMENTS  
by EPA Letter Dated

JUN - 6 2000  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
1021-1749

ACTIVE INGREDIENT:

Permethrin ..... 0.25%

\*\* OTHER INGREDIENTS ..... 99.75%

100.00%

EVERCIDE® - Registered trademark of McLaughlin Gormley King Company

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

FIRST AID

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.  
IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION**

**HUMANS:** Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with eyes and skin. Wash thoroughly with soap and water after handling. Avoid contamination of feed and foodstuffs.

**ANIMALS:** Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic organisms. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwater.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THE ENTIRE LABEL BEFORE EACH USE.

FOR PET USE - USE ONLY ON DOGS & CATS

HOME GARDENS

For use on home gardens. Not for use on plants being grown for commercial sale or commercial use.

EVERCIDE® Permethrin Dust 2786 [brand insecticide] controls a wide range of insect pests. Apply when insects first appear before damage from feeding occurs. Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Allow dust to settle in treated areas before reentering. Wash hands thoroughly after handling and before eating or smoking. Do not allow dust to wind drift to nontarget plants or water supplies. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. For plants above chest height, consider using [brand insecticide].

Vegetables and Melons

The number in parentheses ( ) following each crop grouping is the pre-harvest interval, e.g., the number of days that must be observed between application and harvest.

- Artichokes (0): To control (kill) *artichoke plume moth, leafminer*. Do not exceed 5 applications per season.
- Asparagus (1): To control (kill) *asparagus beetle, cutworm*. Do not exceed 4 applications per season.
- Asparagus (post-harvest): To control (kill) *larval and adult stage of asparagus beetle, tarnished plant bug, lygus bug, collards and adult stage of Japanese beetle*. Dust ferns or bush growth after spear harvest when insects are present.
- Celery - see leafy vegetables.
- Cole Crops (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Chinese Cabbage (tight-heading variation only), Chinese Broccoli (1): To control (kill) *beet armyworm, cabbage aphid, cabbage looper, cutworm, diamondback moth, flea beetle, imported cabbageworm, leafminer, plant bug, thrips*. Apply every 5 days or as needed. Do not exceed 8 applications per season on cauliflower, Brussels sprouts and broccoli, or 5 applications on cabbage.
- Corn - see sweet corn.
- Cucurbits (Balsam Pear - bitter melon: Chinese waxgourd - Chinese preserving melon; Citron melon; Cucumber; Gherkin; Edible gourds; Melons including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honeyballs, mango melon, muskmelon, Persian melon; Pumpkin, Squash - summer and winter; Watermelon) (0): To control (kill) *aphid, cabbage looper, cucumber beetle (adult), cutworm, leafhopper, leafminer, melonworm, pickleworm, plant bug, squash bug*. Do not exceed 8 applications per season.
- Eggplant (3): To control (kill) *Colorado potato beetle, vegetable leafminer, flea beetle*. Do not exceed 10 applications per season.
- Leafy Vegetables (Amaranth, Arugula, Celery, Chervil, Corn Salad, Chrysanthemum (edible-leaved and garland), Cress (garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Rhubarb, Spinach, Swiss chard (1): To control (kill) *alfalfa looper, aphid, beet armyworm, cabbage looper, corn earworm, cutworm, European corn borer, fall armyworm, green cloverworm, Southern armyworm, tobacco budworm, vegetable leafminer*. Do not exceed 5 applications per season.
- Melons - see Cucurbits.
- Onions (dry bulb & garlic) (1): To control (kill) *armyworm, cutworm, leafminer, onion maggot (adult), onion thrips, stink bug*. Do not exceed 6 applications per season.
- Peppers (bell) (3): To control (kill) *beet armyworm, aphid, cabbage looper, corn earworm, European corn borer, flea beetle, pepper weevil, vegetable leafminer*. Do not exceed 8 applications per season.
- Potatoes (14): To control (kill) *beet armyworm, cabbage looper, Colorado potato beetle, cutworm, European corn borer, potato aphid, potato flea beetle, potato leafhopper, potato psyllid, potato tuberworm, tarnished plant bug*. Do not exceed 8 applications per season.
- Spinach (1): To control (kill) *aster leafhopper, beet armyworm, cabbage looper, corn earworm, cutworm, European corn borer, leafhopper*. Do not exceed 5 applications per season.
- Sweet Corn (1): To control (kill) *corn earworm, corn rootworm (adult), European corn borer, fall armyworm, hop vine borer, leafhopper, Southern armyworm, stalk borer*. Dust every 5 days or as needed. Do not exceed 6 applications per season. (FL-only: 8)
- Tomatoes (0): To control (kill) *aphid, beet armyworm, cabbage looper, Colorado potato beetle, flea beetle, granulate cutworm, hornworm, vegetable leafminer, tomato fruitworm, tomato pinworm*. Do not exceed 6 applications per season.

Ornamentals (including Roses)

Contact insects directly whenever possible. Repeat treatment as needed.

Controls (Kills): *corn earworms, fall cankerworm, fall webworm, Japanese beetle (adults), leafminer and omnivorous leafroller* on the following plants:

- African violets, Ageratum, Arizona Cypress, Aster, Azalea, Begonia, Camellia, Carnation, Chrysanthemum, Coleus, Common Ninebark, Common Snowberry, Crassula, Dahlias, Delphinium, Dogwood, Exacum, Euonymous, English Ivy, Fuchsia, Geranium, Gladiolus, Gold Bells, Grape Vines (ornamental), Hypoestes, Ivy, Laurel, Lilac, Marigold, Mock Orange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Poinsettia, Rhododendron, Rose, Snapdragon, Stocks, Taxus, Wax Plant, Weeping Jew, Weeping Fig, Zinnia.

Fruit and Nut Trees

The number in parentheses ( ) following each crop grouping is the pre-harvest interval, e.g., the number of days that must be observed between application and harvest. Also includes relevant comment.

- Apples (Do not apply after petal fall): To control *apple ahpid, green fruitworm, redbanded leafroller, plum curculio, spotted tentiform leafminer, tarnished plant bug, white apple leafroller*. Do not exceed 1 application per season.
- Cherries (In the East, do not exceed 4 applications after petal fall): To control *green fruitworm, lesser peachtree borer, plum curculio, redbanded leafroller, rose chafer, tarnished plant bug*. Do not exceed 6 applications per season.
- Cherries (In the West, do not exceed 3 applications after petal fall): To control *green fruitworm, lesser peachtree borer, plum curculio, redbanded leafroller, rose chafer, tarnished plant bug*. Do not exceed 4 applications per season.
- Peaches (14): To control *green fruitworm, lesser peach tree borer, oriental fruit moth, peach twig borer, plum curculio, rose chafer, tarnished plant bug*.

Pears – Dormant (May be applied dormant through the delayed dormant growth periods only): To control *codling moth*, *green fruitworm*, and *pear psylla*. Do not exceed 2 applications per season.

Pears – Pre-bloom (May be applied from dormant through bud burst): To control *codling moth*, *green fruitworm*, and *pear psylla*. Do not exceed 2 applications per season.

Walnuts (1): To control *codling moth*, *navel orangeworm*, and *walnut husk fly*. Do not exceed 8 applications per season.

BEEF AND DAIRY CATTLE AND HORSES

Can be used in dust bags, shaker can and mechanical dust applicator. *Horn Flies, Lice, Face Flies*. Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through which the animals must pass daily for water, feed or minerals. Bags may also be placed in loafing sheds or in front of mineral feeders. For dairy cows, bags may be suspended in the exit through which the cows leave the milking barn. The bags should hang 4 to 6 inches below the back line of the cattle. For reduction of *Face Flies*, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted.

*Horn Flies, Lice* - Direct Application: Apply 2 oz. of dust per animal by shaker can or dust glove over the head, neck, shoulders, back and legs. Repeat as necessary.

SWINE

*Lice* on Swine Direct Application: Apply only 1 ounce per head as a uniform coat to the head, shoulder and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary but not more often than once every 10 days. In severe infestation, both animals and the bedding may be treated. Do not ship animals for slaughter within 5 days of treatment.

Poultry

To control *Northern Fowl Mites*, apply at a rate of 1 lb. per 100 birds. Ensure thorough treatment of vent area.

PETS

USE ONLY ON DOGS OR CATS: Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing, or medicated animals.

To control (kill) *fleas, ticks and lice*. Use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin paying close attention to legs and feet. Use 1/2 oz. on pets 20 pounds and under and 1 oz. on pets over 20 pounds. Reapply every two weeks. This product should not be applied by children. Do not use on puppies or kittens under 12 weeks of age. Dust pet bedding and doghouse using a shaker can or other dust applicator.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in cool, dry place preferably a locked storage area. Keep container closed.

DISPOSAL: Do not reuse empty container. Wrap container in several layers of newspaper and discard in garbage.

Net Contents \_\_\_\_\_

Manufactured by:

Mc LAUGHLIN GORMLEY KING COMPANY  
8810 Tenth Avenue North  
Minneapolis, MN 55427  
Made in USA

EPA Reg. No. 1021-

EPA Est. No. 1021-MN-2

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....