1021-170 Please read instructions before con	19	2-1	4-200	3			1)
United States Environmental Protection A					Registration Amendme	n	60. Approval expires 05-31-98 OPP Identifier Number
	Washing	ton, DC 20460		X	Other		
		Application 1			<u> </u>		10-14-14-14-14-14-14-14-14-14-14-14-14-14-
1. Company/Product Number 1021-1749			2. EPA Product Manager George LaRocca			3. Proposed	Classification
4. Company/Product (Name)			PM#			☑ None	Restricted
EVERCIDE Permethrin Dust 2786 5. Name and Address of Applicant (Include Zip Code)			6. Expedited Review. In accordance with FIFRA Se				ion 3/c)/3)/b)/i) my
1	MLEY KING COMPANY JE NORTH		1 '	ilar or identic	al in compositi		
Check if this	is a new address		Product Nan	ne			
			Section - II				
Amendment	- Explain below.				Final printed	labels in resp	onse to
_	•			_	Agency letter	dated	NOTIFICATION
	on in response to Agency let	Ket dated			"Me Too" Ap	plication.	FEB 1 4 2003
	- Explain below. nal page(s) if necessary. (F				Other - Expla	in below.	
			Section - III			· · · · · · · · · · · · · · · · · · ·	·
Material This Product W	ill Re Packaged In:		SOCEION - III		,		
Child-Resistant Packaging			Water Soluble	Packaging		2. Type of C	Container
☐ Yes*	Yes		l n	Yes		Metal	
∏ No	No No			No		Plastic	3
	If "Yes"	No. per	If "Yes"		No. per	Glass	
*Certification must be submitted	Unit Packaging wgt.	container I	Package wgt.		container	Paper	(Specify)
3. Location of Net Content	s information	4. Size(s) Reta	il Container		5. Location o	f Label Directi	
Label [Container	// 0.20(0) / (0.20				On Label	
6. Manner in Which Label	is Affixed to Product	Lithograph			Other	OU CYDening	accompanying product
		Paper glued		L	3.1.0.		
		Stenciled	Section - IV				
1 Contact Point (Complete	e items directly below for ide			tacted if nec	esseny to proc	ess this annii	cation.)
Name	Tomo and any addition to the	Title	10001	aoioa, ii noo	sodary, to proc		lo. (Include Area Code)
Thomas A. Lennan		Registration Specialist				(763) 544-	,
1	nts I have made on this fom knowingly false or misleadin w.				•	te.	6. Date Application Received (Stamped)
2. Signature Unimer A. Lennan		3. Title Registration Specialist					00 00 00 00 00 00 00 00 00 00 00 00 00
4. Typed Name	A. Man	5. Date					* C + C + C + C + C + C + C + C + C + C
Thomas A. Lennan		1/28/2003					a saise

2786-0203



EVERCIDE® Permethrin Dust 2786

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4319 U.S.A.

TELEPHONE (763) 544-0341

NOTIFICATION

FEB 1 4 2003

Optional marketing statements:

Contains PERMETHRIN KILLS beetles, caterpillars, bugs and many more Ready to Use Controls ornamental pests To control (kill) vegetable pests

Use on home garden fruits, vegetables and ornamental plants (including roses)

Use on fruit & nut trees

Use on dogs, cats and pet bedding to control fleas, ticks and lice

For control of Horn Flies and Lice on Beef and Dairy Cattle Aids in reducing Face Fly population on Beef and Dairy Cattle For control of Lice on Swine For control of Northern Fowl Mites on Poultry

For use in dust bags, [this] shaker can or mechanical dust applicator

Easy-to-use powder duster

No unpleasant odor

ACTIVE INGREDIENT:

Permethrin 0.25% OTHER INGREDIENTS 99.75% 100.00%

EVERCIDE® - Registered trademark of McLaughlin Gormley King Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

	FIRST AID .
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES:	 Hold eye open and rinse slowing and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 15 minutes, then continue rinsing eyes. Call a poison control center for treatment advice.
Have the product cor	 Call a poison control center for treatment advice. ntainer or label with you when calling a poison control center or doctor or going for treatment. For information regarding

emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

HUMANS: Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ANIMALS: Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic organisms. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwater.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

READ THE ENTIRE LABEL BEFORE EACH USE.

FOR PET USE - USE ONLY ON DOGS & CATS

HOME GARDENS

For use on home gardens. Not for use on plants being grown for commercial sale or commercial use.

EVERCIDE® Permethrin Dust 2786 [brand insecticide] [this product] controls a wide range of insect pests. Apply when insects first appear before damage from feeding occurs. Wearing household latex or rubber gloves, use a dust applicator such as a [this] shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Allow dust to settle in treated areas before reentering. Wash hands thoroughly after handling and before eating or smoking. Do not allow dust to wind drift to nontarget plants or water supplies. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. For plants above chest height, consider using [brand insecticide].

Vegetables and Melons

The number in parentheses () following each crop grouping is the pre-harvest interval, e.g., the number of days that must be observed between application and harvest.

Artichokes (0): To control (kill) artichoke plume moth, leafminer. Do not exceed 5 applications per season.

Asparagus (1): To control (kill) asparagus beetle, cutworm. Do not exceed 4 applications per season.

Asparagus (post-harvest): To control (kill) larval and adult stage of asparagus beetle, tarnished plant bug, lygus bug, collards and adult stage of Japanese beetle. Dust ferns or bush growth after spear harvest when insects are present. (Not for use in California).

Celery - see leafy vegetables.

Cole Crops (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Chinese Cabbage (tight-heading variation only), Chinese Broccoli (on gai Ion, white flowering). (Not for use in California). (1): To control (kill) beet armyworm, cabbage aphid, cabbage looper, cutworm, diamondback moth, flea beetle, imported cabbageworm, leafminer, plant bug, thrips. Apply every 5 days or as needed. Do not exceed 8 applications per season on cauliflower, Brussels sprouts and broccoli, or 5 applications on cabbage.

Corn - see sweet corn.

Cucurbits (Balsam Pear - bitter melon: Chinese waxgourd - Chinese preserving melon; Citron melon; Cucumber; Gherkin; Edible gourds; Melons including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honeyballs, mango melon, muskmelon, Persian melon; Pumpkin, Squash - summer and winter; Watermelon) (0): To control (kill) aphid, cabbage looper, cucumber beetle (adult), cutworm, leafhopper, leafminer, melonworm, pickleworm, plant bug, squash bug. Do not exceed 8 applications per season.

Eggplant (3): To control (kill) Colorado potato beetle, vegetable leafminer, flea beetle. Do not exceed 10 applications per season.

Eggplant (3): To control (kill) Colorado potato beetle, vegetable leafminer, flea beetle. Do not exceed 10 applications per season.

Leafy Vegetables (Amaranth, Arugula, Celery, Chervil, Corn Salad, Chrysanthemum (edible-leaved and garland), Cress (garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Rhubarb, Spinach, Swiss chard (1): To control (kill) alfalfa looper, aphid, beet armyworm, cabbage looper, com earworm, cutworm, European corn borer, fall armyworm, green cloverworm, Southern armyworm, tobacco budworm, vegetable leafminer. Do not exceed 5 applications per season.

Melons - see Cucurbits.

Onions (dry bulb & garlic) (1): To control (kill) armyworm, cutworm, leafminer, onion maggot (adult), onion thrips, stink bug. Do not exceed 6 applications per season.

Peppers (bell) (3): To control (kill) beet armyworm, aphid, cabbage looper, com earworm, European com borer, flea beetle, pepper weevil, vegetable leafminer. Do not exceed 8 applications per season.

Potatoes (14): To control (kill) beet armyworm, cabbage looper, Colorado potato beetle, cutworm, European com borer, potato aphid, potato flea beetle, potato leafnopper, potato psyllid, potato tuberworm, tamished plant bug. Do not exceed 8 applications per season.

Spinach (1): To control (kill) aster leafhopper, beet armyworm, cabbage looper, com earworm, cutworm, European corn borer, leafhopper. Do not exceed 5 applications per season.

Sweet Corn (1): To control (kill) com earworm, com rootworm (adult), European corn borer, fall armyworm, hop vine borer, leafhopper, Southern armyworm, stalk borer. Dust every 5 days or as needed. Do not exceed 6 applications per season. (FL-only: 8)

Tomatoes (0): To control (kill) beet armyworm, cabbage looper, Colorado potato beetle, granulate cutworm, hornworm, vegetable leafminer, tomato fruitworm, tomato pinworm. Do not exceed 6 applications per season.

Ornamentals (including Roses, Flowers, Shrubs and Small Trees)

Contact insects directly whenever possible.

Controls (Kills): aphid, bagworms, cabbage looper, cicadas, com earworms, cutworm, exposed thrips, fall cankerworm, fall webworm, inchworms, Japanese beetle (adults), leafhopper, leafminer, and omnivorous leafroller, mealybugs, spider mites, stinkbugs, adult strawberry root weevil, adult obscure root weevil, thrips and whitefly on the following plants:

African violets, Ageratum, Arizona Cypress, Aster, Azalea, Begonia, Camellia, Carnation, Chrysanthemum, Coleus, Common Ninebark, Common Snowberry, Crassula, Dahlias, Delphinium, Dogwood, Exacum, Euonymous, English Ivy, Fuchsia, Geranium, Gladiolus, Gold Bells, Grape Vines (ornamental), Hypoestes, Ivy, Laurel, Lilac, Marigold, Mock Orange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Petunia,

Fruit and Nut Trees

The number in parentheses () following each crop grouping is the pre-harvest interval, e.g., the number of days that must be observed between application and harvest. Also includes relevant comment.

Philodendron, Poinsettia, Rhododendron, Rose, Snapdragon, Stocks, Taxus, Wax Plant, Wandering Jew, Weeping Fig, Zinnia.

Apples (Do not apply after petal fall): To control apple ahpid, green fruitworm, redbanded leafroller, plum curculio, spotted tentiform leafminer, tamished plant bug, white apple leafroller. Do not exceed 1 application per season.

Cherries (In the East, do not exceed 4 applications after petal fall): To control green fruitworm. lesser peachtree borer, plum curculio, redbanded leafroller, rose chafer, tarnished plant bug. Do not exceed 6 applications per season.

Cherries (In the West, do not exceed 3 applications after petal fall): To control green fruitworm. lesser peachtree borer, plum curculio, redbanded leafroller, rose chafer, tamished plant bug. Do not exceed 4 applications per season.

Peaches (14): To control green fruitworm, lesser peach tree borer, oriental fruit moth, peach twig borer, plum curculio, rose chafer, tarnished plant bug. Dust lightly. Apply up to 5 times per season.

Pears – Dormant (May be applied dormant through the delayed dormant growth periods only): To control pear psylla. Do not exceed 2 applications per season.

Pears – Pre-bloom (May be applied from dormant through bud burst): To control codling moth, green fruitworm, and pear psylla. Do not exceed 2 applications per season.

Wainuts (1): To control codling moth, navel orangeworm, and wainut husk fly. Do not exceed 8 applications per season.

BEEF AND DAIRY CATTLE AND HORSES

Can be used in dust bags, [this] shaker can and mechanical dust applicator. Horn Flies, Lice, Face Flies: Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through which the animals must pass daily for water, feed or minerals. Bags may also be placed in loafing sheds or in front of mineral feeders. For dairy cows, bags may be suspended in the exit through which the cows leave the milking barn. The bags should hang 4 to 6 inches below the back line of the cattle. For reduction of Face Flies, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted.

Hom Flies, Lice - Direct Application: Apply 2 oz. of dust per animal by [this] shaker can or dust glove over the head, neck, shoulders, back and legs.

SWINE

Lice on Swine Direct Application: Apply only 1 ounce per head as a uniform coat to the head, shoulder and back by use of a [this] shaker can or suitable mechanical dust applicator. Do not apply more often than once every 10 days. In severe infestation, both animals and the bedding may be treated. Do not ship animals for slaughter within 5 days of treatment.

<u>POULTRY</u>

To control Northern Fowl Mites, apply at a rate of 1 lb. per 100 birds. Ensure thorough treatment of vent area.

PETS

USE ONLY ON DOGS OR CATS: Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing, or medicated animals.

To control (kill) fleas, ticks and lice: Use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin paying close attention to legs and feet. Use 1/2 oz. on pets 20 pounds and under and 1 oz. on pets over 20 pounds. Reapply every two weeks. This product should not be applied by children. Do not use on puppies or kittens under 12 weeks of age. Dust pet bedding and doghouse using a [this] shaker can or other dust applicator.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in cool, dry place preferably a locked storage area. Keep container closed.

DISPOSAL: Do not reuse empty container. Wrap container in several layers of newspaper and discard in garbage.

Net Contents

Manufactured by:
Mc LAUGHLIN GORMLEY KING COMPANY
8810 Tenth Avenue North
Minneapolis, MN 55427
Made in USA

EPA Reg. No. 1021-1749

EPA Est. No. 1021-MN-2

8810 Tenth Avenue North • Minneapolis, MN 55427-4372 U.S.A 763-544-0341 • 800-645-6466 • Fax 763-544-6437 • www.mgk.com

January 28, 2003

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. ENVIRONMENTAL PROTECTION AGENCY
Ariel Rios Building
1200 Pennsylvania Avenue NW
Washington, DC 20460

Subject:

EVERCIDE® Permethrin Dust 2786

EPA Reg. No. 1021-1749

Notification

Dear Sir or Madam:

By notification, we are revising our label to remove restrictions specific to the state of California since that state is now lifting those restrictions.

2 copies of our current label are enclosed, 1 copy has additions highlighted in yellow.

I certify that this notification is consistent with PR Notice 98-10, PR Notice 92-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, then this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

If there are any questions or you need anything further, please contact this office.

FEB 1 4 2003

Sincerely,

McLAUGHLIN GORMLEY KING COMPANY

Thomas A. Lennan

Registration Specialist

Enclosure