

1021-1749

9/30/2011

10413



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460-0001

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

September 30, 2011

Steven Ditto  
McLaughlin Gormley King Company  
8810 Tenth Avenue North  
Minneapolis, MN 55427

Subject: Amended Reregistration Label  
Product Name: EVERCIDE® Permethrin Dust 2786  
EPA Registration Number: 1021-1749  
EPA Decision Number: 413877

Dear Mr. Ditto:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the reregistration of the above referenced product in connection with the **Permethrin RED**, and has concluded that your submission is acceptable.

NOTE: This product is not being reregistered under sections 3(c)5 and 4(g) of FIFRA at this time.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on non-refillable containers. The code may appear either on the label or durably marked on the container itself, and can be added by non-notification per PRN 98-10.


Please note that the record for this product currently contains the Confidential Statements of Formulation (CSFs) listed below. Any previously dated CSFs are superseded.

- Basic CSF, dated 2/11/2010

A copy of your label stamped "Accepted" is enclosed along with copies of the acute toxicity and product chemistry reviews completed for the subject product. Products shipped after 12 months from the date of this amendment or the next printing of the label whichever occurs first, must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e).

If you have any questions about this letter, please contact either Linda Arrington at (703) 305 6249 or via e-mail at [Arrington.Linda@epa.gov](mailto:Arrington.Linda@epa.gov).

Sincerely yours,



Richard J. Gebken  
Product Manager (10)  
Insecticide Branch  
Registration Division (7505P)

Enclosures: Label stamped "Accepted," dated 9/30/2011  
Acute Toxicity Review, dated 5/27/2010  
Product Chemistry Review, dated 12/22/10

30813

2786\_1021-1749\_0811



McLAUGHLIN  
GORMLEY  
KING COMPANY

EVERCIDE® Permethrin Dust 2786

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4319 U.S.A. - Telephone (763) 544-0341

# EVERCIDE® Permethrin Dust 2786


## MASTER LABEL

SUB\_LABEL A: AGRICULTURAL USE

SUB LABEL B: RESIDENTIAL USE

ACTIVE INGREDIENT:		
Permethrin .....	0.25%	
OTHER INGREDIENTS .....	99.75%	
	<u>100.00%</u>	

EVERCIDE® - Registered trademark of McLaughlin Gormley King Company



### KEEP OUT OF REACH OF CHILDREN CAUTION

Net Contents \_\_\_\_\_  
Manufactured by:  
Mc LAUGHLIN GORMLEY KING COMPANY  
8810 Tenth Avenue North  
Minneapolis, MN 55427

EPA Reg. No. 1021-1749

EPA Est. No. 1021-MN-2

40 of 13

EVERCIDE® Permethrin Dust 2786

SUB-LABEL A: AGRICULTURAL USE

50413

2786\_1021-1749\_0811



McLAUGHLIN  
GORMLEY  
KING COMPANY

EVERCIDE® Permethrin Dust 2786

[Label for Agricultural Use]

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4319 U.S.A. - Telephone (763) 544-0341

[ ] OR ( ) indicates alternate or optional wording

[Contains PERMETHRIN]  
[Ready to Use]

[For control of Horn Flies and Lice on Beef and Dairy Cattle]  
[Aids in reducing Face Fly population on Beef and Dairy Cattle]

[For control of Lice on Swine]  
[For control of Northern Fowl Mites on Poultry]  
[For use in dust bags, [this] shaker can or mechanical dust applicator]  
[Easy-to-use powder duster]  
[No unpleasant odor]

ACTIVE INGREDIENT:	
Permethrin .....	0.25%
OTHER INGREDIENTS .....	99.75%
	100.00%

EVERCIDE® - Registered trademark of McLaughlin Gormley King Company

KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### FIRST AID

IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF IN EYES:	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 15 minutes, then continue rinsing eyes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

Net Contents \_\_\_\_\_  
Manufactured by:  
Mc LAUGHLIN GORMLEY KING COMPANY  
8810 Tenth Avenue North  
Minneapolis, MN 55427

EPA Reg. No. 1021-1749

EPA Est. No. 1021-MN-2

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

#### PERSONAL PROTECTIVE EQUIPMENT:

Some materials that are chemical resistant to this product are barrier laminate butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride, viton. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

Loaders, applicators, and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves
- NIOSH approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any N, R, P, or HE filter

60913

Application with aerial or motorized ground equipment is prohibited.

**USER SAFETY RECOMMENDATIONS:**

User should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.  
 User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.  
 User should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

Under some conditions, it may also have a potential for transport into surface water in runoff (primarily absorbed to suspended soil particles), for several months or more after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded area, and areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THE ENTIRE LABEL BEFORE EACH USE.

**PRECAUTIONS AND RESTRICTIONS:**

Do not make applications during rain.  
Do not water the treated area to the point of run-off.  
Use of handled power duster equipment is prohibited.

Except when treating livestock do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Except when treating livestock only protected handlers may be in the area during application.

Do not breathe dusts.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

When used in dairy barns or facilities: close milk bulk tank lids to prevent contamination from dust and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.

**BEEF AND DAIRY CATTLE AND HORSES**

Can be used in dust bags, [this] shaker can and mechanical dust applicator. *Horn Flies, Lice, Face Flies*: Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through which the animals must pass daily for water, feed or minerals. Bags may also be placed in loafing sheds or in front of mineral feeders. For dairy cows, bags may be suspended in the exit through which the cows leave the milking barn. The bags should hang 4 to 6 inches below the back line of the cattle. For reduction of *Face Flies*, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted.

*Horn Flies, Lice* - Direct Application: Apply 2 oz. of dust per animal by [this] shaker can or dust glove over the head, neck, shoulders, back and legs.

**SWINE**

*Lice* on Swine Direct Application: Apply only 1 ounce per head as a uniform coat to the head, shoulder and back by use of a [this] shaker can or suitable mechanical dust applicator. Do not apply more often than once every 10 days. In severe infestation, both animals and the bedding may be treated. Do not ship animals for slaughter within 5 days of treatment.

**POULTRY**

To control *Northern Fowl Mites*, apply at a rate of 1 lb. per 100 birds. Ensure thorough treatment of vent area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:**

Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

**Pesticide Disposal:**

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

**Container Disposal:**

Nonrefillable container. Do not reuse or refill this container.

Completely empty can by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill

EVERCIDE® Permethrin Dust 2786

SUB-LABEL B: RESIDENTIAL USE



9013

2786\_1021-1749\_0811



McLAUGHLIN  
GORMLEY  
KING COMPANY

EVERCIDE® Permethrin Dust 2786

[Label for Residential Use]

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4319 U.S.A. - Telephone (763) 544-0341

[ ] OR ( ) indicates alternate or optional wording

[Contains PERMETHRIN]  
[KILLS listed beetles, caterpillars, bugs] [Ready to Use]  
[Controls listed ornamental pests] [To control (kill)  
listed vegetable pests]  
[Use on listed fruits, vegetables and ornamental plants  
(including roses) in residential gardens]

[Use on listed fruit & nut trees]  
[Use on dogs and dog bedding to control fleas, ticks and lice]

ACTIVE INGREDIENT:		
Permethrin .....	0.25%	
OTHER INGREDIENTS .....	99.75%	
	<hr/>	100.00%

EVERCIDE® - Registered trademark of McLaughlin Gormley King Company

**DO NOT USE ON CATS**  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

FIRST AID

IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF IN EYES:	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 15 minutes, then continue rinsing eyes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

Net Contents \_\_\_\_\_  
Manufactured by:  
Mc LAUGHLIN GORMLEY KING COMPANY  
8810 Tenth Avenue North  
Minneapolis, MN 55427

EPA Reg. No. 1021-1749

EPA Est. No. 1021-MN-2

**PRECAUTIONARY STATEMENTS**

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION**

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long sleeved shirt, long pants, shoes plus socks, and chemical resistant gloves (made out of any waterproof material). Wash hands thoroughly with soap and water after using and before eating, drinking, chewing gum, using tobacco, or using the toilet.

**DO NOT USE ON CATS**

May be toxic or potentially fatal if applied to or ingested by cats. Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movements. If this occurs, immediate veterinary care should be provided.



ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THE ENTIRE LABEL BEFORE EACH USE.

**PRECAUTIONS AND RESTRICTIONS:**

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Do not make applications during rain.

Do not water the treated area to the point of run-off.

Use of handheld power duster equipment is prohibited.

Not for use on plants being grown for commercial sale or commercial use.

Do not allow adults, children, or pets to enter the treated area until dusts have settled.

**DOGS – Use on Dogs Only:**

Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing, or medicated animals.

To control (kill) fleas, ticks and lice: Use product only outside or in a well-ventilated area. Wearing chemical resistant gloves, dust entire animal avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin paying close attention to legs and feet. Use 1/2 oz. on pets 20 pounds and under and 1 oz. on pets over 20 pounds. Reapply every two

weeks. Children should not apply this product. Do not use on puppies under 12 weeks of age. Dust pet bedding and doghouse using a [this] shaker can or other dust applicator.

HOME GARDENS

For use on home gardens. Not for use on plants being grown for commercial sale or commercial use. EVERCIDE® Permethrin Dust 2786 [brand insecticide] [this product] controls a wide range of insect pests. Apply when insects first appear before damage from feeding occurs. [How to Apply:] Wearing chemical resistant gloves, use a dust applicator such as a [this] shaker can, dust gun, puffer or other bulbous duster, or other appropriate duster as available. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Allow dust to settle in treated areas before reentering. Do not allow dust to wind drift to non-target plants or water supplies. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. For plants above chest height, consider using [brand insecticide].

The number in parentheses ( ) following each crop grouping is the pre-harvest interval, e.g., the number of days that must be observed between application and harvest.

Vegetables and Melons

- Artichokes (0): To control (kill) *artichoke plume moth, leafminer*. The minimum retreatment interval is 10 days, the maximum number of applications per year is 3.
- Asparagus (1): To control (kill) *asparagus beetle, cutworm*. The minimum retreatment interval is 7 days, the maximum number of applications per year is 4.
- Asparagus (post-harvest): To control (kill) *larval and adult stage of asparagus beetle, tarnished plant bug, lygus bug, collards and adult stage of Japanese beetle*. Dust ferns or bush growth after spear harvest when insects are present. (Not for use in California).
- Broccoli (1): To control (kill) *beet armyworm, cabbage aphid, cabbage looper, cutworm, diamondback moth, flea beetle, imported cabbageworm, leafminer, plant bug, thrips*. The minimum retreatment interval is 4 days, the maximum number of applications per year is 5.
- Brussels Sprouts (1): To control (kill) *beet armyworm, cabbage aphid, cabbage looper, cutworm, diamondback moth, flea beetle, imported cabbageworm, leafminer, plant bug, thrips*. The minimum retreatment interval is 5 days, the maximum number of applications per year is 4.
- Cabbage (Tight Headed Varieties Only) (1): To control (kill) *beet armyworm, cabbage aphid, cabbage looper, cutworm, diamondback moth, flea beetle, imported cabbageworm, leafminer, plant bug, thrips*. The minimum retreatment interval is 5 days, the maximum number of applications per year is 2.
- Cantaloupe (0): To control (kill) *aphid, cabbage looper, cucumber beetle (adult), cutworm, leafhopper, leafminer, melonworm, pickleworm, plant bug, squash bug*. The minimum retreatment interval is 7 days, the maximum number of applications per year is 4.
- Cauliflower (1): To control (kill) *beet armyworm, cabbage aphid, cabbage looper, cutworm, diamondback moth, flea beetle, imported cabbageworm, leafminer, plant bug, thrips*. The minimum retreatment interval is 5 days, the maximum number of applications per year is 4.
- Celery (1): To control (kill) *alfalfa looper, aphid, beet armyworm, cabbage looper, corn earworm, cutworm, European corn borer, fall armyworm, green cloverworm, Southern armyworm, tobacco budworm, vegetable leafminer*. The minimum retreatment interval is 7 days, the maximum number of applications per year is 5.
- Chinese Broccoli (1): To control (kill) *beet armyworm, cabbage aphid, cabbage looper, cutworm, diamondback moth, flea beetle, imported cabbageworm, leafminer, plant bug, thrips*. The minimum retreatment interval is 4 days, the maximum number of applications per year is 5.
- Chinese Cabbage (Tight Headed Varieties Only) (1): To control (kill) *beet armyworm, cabbage aphid, cabbage looper, cutworm, diamondback moth, flea beetle, imported cabbageworm, leafminer, plant bug, thrips*. The minimum retreatment interval is 5 days, the maximum number of applications per year is 2.
- Corn - see sweet corn.
- Cucurbits (Balsam Pear - bitter melon; Chinese waxgourd - Chinese preserving melon; Citron melon; Cucumber; Gherkin; Edible gourds; Melons including hybrids such as casaba, crenshaw, honeydew, honeyballs, mango melon, muskmelon, Persian melon; Pumpkin, Squash - summer and winter; Watermelon) (0): To control (kill) *aphid, cabbage looper, cucumber beetle (adult), cutworm, leafhopper, leafminer, melonworm, pickleworm, plant bug, squash bug*. The minimum retreatment interval is 7 days, the maximum number of applications per year is 6.
- Eggplant (3): To control (kill) *Colorado potato beetle, vegetable leafminer, flea beetle*. The minimum retreatment interval is 7 days, the maximum number of applications per year is 4.
- Garlic (1): To control (kill) *armyworm, cutworm, leafminer, onion maggot (adult), onion thrips, stink bug*. The minimum retreatment interval is 10 days, the maximum number of applications per year is 4.
- Leafy Vegetables (Amaranth, Arugula, Chervil, Corn Salad, Chrysanthemum (edible-leaved and garland), Cress (garden and upland), Dandelion, Dock, Endive, Fennel, , Orach, Parsley, Purslane (garden and winter), Rhubarb,

Swiss chard (1): To control (kill) *alfalfa looper, aphid, beet armyworm, cabbage looper, corn earworm, cutworm, European corn borer, fall armyworm, green cloverworm, Southern armyworm, tobacco budworm, vegetable leafminer*. The minimum retreatment interval is 3 days, the maximum number of applications per year is 10.

Lettuce (head and leaf) - To control (kill) *alfalfa looper, aphid, beet armyworm, cabbage looper, corn earworm, cutworm, European corn borer, fall armyworm, green cloverworm, Southern armyworm, tobacco budworm, vegetable leafminer*. The minimum retreatment interval is 7 days, the maximum number of applications per year is 4.

Melons - see Cucurbits.

Onions (dry bulb ) (1): To control (kill) *armyworm, cutworm, leafminer, onion maggot (adult), onion thrips, stink bug*.

The minimum retreatment interval is 7 days, the maximum amount of active ingredient allowed per season is 1.0 pounds.

Peppers (bell) (3): To control (kill) *beet armyworm, aphid, cabbage looper, corn earworm, European corn borer, flea beetle, pepper weevil, vegetable leafminer*. The minimum retreatment interval is 5 days, the maximum number of applications per year is 4.

Potatoes (14): To control (kill) *beet armyworm, cabbage looper, Colorado potato beetle, cutworm, European corn borer, potato aphid, potato flea beetle, potato leafhopper, potato psyllid, potato tuberworm, tarnished plant bug*.

The minimum retreatment interval is 10 days, the maximum number of applications per year is 4.

Spinach (1): To control (kill) *aster leafhopper, beet armyworm, cabbage looper, corn earworm, cutworm, European corn borer, leafhopper*. The minimum retreatment interval is 3 days, the maximum number of applications per year is 3.

Sweet Corn (1): To control (kill) *corn earworm, corn rootworm (adult), European corn borer, fall armyworm, hop vine borer, leafhopper, Southern armyworm, stalk borer*. The minimum retreatment interval is 3 days, the maximum number of applications per year is 4.

Tomatoes (0): To control (kill) *beet armyworm, cabbage looper, Colorado potato beetle, granulate cutworm, hornworm, vegetable leafminer, tomato fruitworm, tomato pinworm*. The maximum number of applications per year is 6. Restriction: Do not apply to tomato varieties which produce mature fruit <1 inch in diameter.

#### Ornamentals (including Roses, Flowers, Shrubs and Small Trees)

Contact insects directly whenever possible.

Controls (Kills): *aphid, bagworms, cabbage looper, cicadas, corn earworms, cutworm, exposed thrips, fall cankerworm, fall webworm, inchworms, Japanese beetle (adults), leafhopper, leafminer, and omnivorous leafroller, mealybugs, spider mites, stinkbugs, adult strawberry root weevil, adult obscure root weevil, thrips and whitefly* on the following plants:

African violets, Ageratum, Arizona Cypress, Aster, Azalea, Begonia, Camellia, Carnation, Chrysanthemum, Coleus, Common Ninebark, Common Snowberry, Crassula, Dahlias, Delphinium, Dogwood, Exacum, Euonymus, English Ivy, Fuchsia, Geranium, Gladiolus, Gold Bells, Grape Vines (ornamental), Hypoestes, Ivy, Laurel, Lilac, Marigold, Mock Orange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Poinsettia, Rhododendron, Rose, Snapdragon, Stocks, Taxus, Wax Plant, Wandering Jew, Weeping Fig, Zinnia.

#### Fruit and Nut Trees

Do not graze livestock in treated areas or feed cover crops from treated orchards/areas to livestock.

Apples (Do not apply after petal fall): To control *apple aphid, green fruitworm, redbanded leafroller, plum curculio, spotted tentiform leafminer, tarnished plant bug, white apple leafroller*. The minimum retreatment interval is 10 days, the maximum number of applications per year is 2.

Cherries (3): To control *green fruitworm, lesser peachtree borer, plum curculio, redbanded leafroller, rose chafer, tarnished plant bug*. The minimum retreatment interval is 10 days, the maximum number of applications per year is 3.

Peaches (14): To control *green fruitworm, lesser peach tree borer, oriental fruit moth, peach twig borer, plum curculio, rose chafer, tarnished plant bug*. Dust lightly. The minimum retreatment interval is 10 days, the maximum number of applications per year is 3.

Pears – Dormant Through Delayed Dormant (Prebloom) Applications: To control *pear psylla*. The minimum retreatment interval is 10 days.

Walnuts (1): To control *codling moth, navel orangeworm, and walnut husk fly*. The minimum retreatment interval is 10 days, the maximum number of applications per year is 3.

**ANTS:**

Ants , Carpenter Ants:

Treat doors, around window frames, ant trails and hills, cracks and crevices and other areas of entry.

**[PRODUCT FACTS]****[Kills Ants & Carpenter Ants]****[Where to Apply:]** Treat doors, around window frames, ant trails and hills, cracks and crevices and other areas of entry.**[How to Apply:]** Wearing chemical resistant gloves, use a dust applicator such as a [this] shaker can. Apply when air is calm to avoid drift and contact with eyes and skin. Allow dust to settle in treated areas before reentering. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces.**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal.

**Pesticide Storage**

Store in cool, dry area, preferably a locked storage area. Keep container closed. Always store pesticides in the original container. Store away from food, pet food, feed, seed, and fertilizers.

**Pesticide Disposal and Container Handling**

Nonrefillable container. Do not reuse or refill this container.

If empty:

Place in trash or offer for recycling if available.

If partly filled:

Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.