1021-1751

9/21/2012

Page 1 7 7

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460-0001



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

SEP 2 1 2012

Mr. Lennan McLaughlin Gormley King Company 8810 Tenth Avenue North Minneapolis, MN 55427

Subject: Amended Reregistration Label Product Name: PYGANIC Crop Protection EC 1.4 EPA Registration Number: 1021-1751 EPA Decision Number: 412428

Dear Mr. Lennan,

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the reregistration of the above referenced product in connection with the Pyrethrin RED(s), and has concluded that your submission is acceptable.

NOTE: This product is **not** being reregistered under sections 3(c)5 and 4(g) of FIFRA at this time.

Please note that the record for this product currently contains the Confidential Statements of Formulation (CSFs) listed below. Any previously dated CSFs are superseded.

- Basic CSF, dated 8/2/2011
- Alternate CSF #1, dated 2/28/01

A copy of your label stamped "Accepted" is enclosed along with copies of the acute toxicity and product chemistry reviews completed for the subject product. Products shipped after 12 months from the date of this amendment or the next printing of the label whichever occurs first, must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e).

If you have any questions about this letter, please contact Jessica Rogala at (703) 347-0263 or via e-mail at <u>rogala.jessica@epa.gov</u>.

Sincerely,

Richard J. Gebken Product Manager (10) Insecticide Branch Registration Division (7504P)

227

7443_1021-1751_0812



| | crops and ornamentals as described below e sprayed at any season of the year nites from hiding | GROU | P | 3A | INSECTICIDE |
|---|---|--|--|---|--|
| Provides rapid know Contains pyrethrum Approved for use of | kdown & kill of plant pests – a botanical insecticide derived from chrysan n over 200 fruits and vegetables with other insect control products | themums | | | tipaq mos- interfection people aligner aligner aligner |
| Can be used on da User friendly contai Non-flammable Controls more than One gallon of produ Low odor | y of harvest ner for convenient and accurate use 40 insects ict treats up to 8 acres | | as HECHINA ROAMANEAN HIGH 391 10 HIGH Jack HAR | opicator with constation with constation ablact alter path any R.P. alternation and a count | |
| ACTIVE ING | and the second | ang ang | | notifictes wit plane | 1.40% 8.60% |
| PYGANIC™ | CAUTION | REACH OF CHILDRE | N IÓN | n enclination en nen exectioned neider storchest | |
| PYGANIC™ | KEEP OUT OF | REACH OF CHILDRE PRECAUC alguien para que se la ex | N CIÓN Kplique a us | sted en detalle. | Redon stater notion Redon ben ben Ludead chaiters Redon State |
| PYGANIC™ | KEEP OUT OF CAUTION Si usted no entiende la etiqueta, busque a (If you do not understand the label | REACH OF CHILDRE PRECAUC alguien para que se la ex | N CIÓN Kplique a us | sted en detalle. | Redon stater notion Redon ben ben Ludead chaiters Redon State |
| PYGANIC™ | KEEP OUT OF CAUTION Si usted no entiende la etiqueta, busque a (If you do not understand the label | REACH OF CHILDRE PRECAUC alguien para que se la ex i, find someone to explain IRST AID tor immediately for treat f able to swallow. to do so by the poisor | N xplique a us it to you in atment ac n control c | sted en detalle. detail) lvice. enter or docto | Historiana stateli fotiale Lukard and versi fotiale rausa fitter Lukar Safety Rucchin Machine Ineres geos teenas universi gioves termore resource gioves termore resource JTC |
| IF SWALLOWED: IF ON SKIN OR CLOTHING: | KEEP OUT OF CAUTION Si usted no entiende la etiqueta, busque a (If you do not understand the label F • Call a poison control center or doc • Have person sip a glass of water if • Do not induce vomiting unless told | REACH OF CHILDRE PRECAUC a alguien para que se la ex- a find someone to explain IRST AID tor immediately for treat a able to swallow. to do so by the poisor an unconscious person of water for 15-20 mir | N Kplique a us it to you in atment ac n control c n. hutes. | ated en detalle. a detail) lvice. center or docto | Kolige statel aduate Kolige statel aduate Lives Science Science Lives Science Russes Micros Balance, Inen gloves Balance asso |
| IF SWALLOWED: | KEEP OUT OF CAUTION Si usted no entiende la etiqueta, busque a (If you do not understand the label F • Call a poison control center or doc • Have person sip a glass of water if • Do not induce vomiting unless told • Do not give anything by mouth to a • Take off contaminated clothing. • Rinse skin immediately with plenty | REACH OF CHILDRE PRECAUC alguien para que se la ex i, find someone to explain IRST AID tor immediately for treat able to swallow. to do so by the poisor an unconscious person of water for 15-20 mir tor for treatment advice d gently with water for , after the first 5 minute | N cplique a use it to you in atment ac n control con- nutes. e. 15-20 min es, then c | ated en detalle. detail) lvice. enter or docto | Analos latars websit and out over the over many sectors breaking many and the over and the over th |

CAUTION

ACCTATION SEP 2 1 2012 Inder the Pederal Insecticide. Inder the Pederal Insecticide. Inder the Pederal Insecticide as Ensembled the pestacide Act, as Ensembled the pestacide Particide Inder Particide Inder

Page 1 of 5 7443_1021-1751_0812 PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of **barrier laminate, butyl rubber, nitrile rubber, or viton**. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Mixers, loaders, applicators, and other handlers must wear the following:

--long-sleeve shirt, --long pants,

--chemical resistant gloves,

--shoes and socks.

In addition to the above PPE, applicators using hand held foggers in an enclosed area must wear a half-face, full face or hood-style NIOSH approved respirator with:

--a dust/mist filtering cartridge (MSHA/NIOSH approve number prefis TC-21C), or

--a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or

--a cartridge or canister with any R,P or HE filter.

When handlers use aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240 (d) (4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

See engineering controls for additional requirements.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations:

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. **Remove clothing/PPE immediately if pesticide** gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 70.240 (d)(6)

Human flagging is prohibited. Flagging to support aerial application is limited to the use of the Global Positioning System (GPA) or Mechanical flaggers.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through runoff. This product has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Page 2 of 5 7443_1021-1751_0812

477

USE RESTRICTIONS:

.

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Only protected handlers may be in the area during application.
- Do not make application during the rain.
- Do not wet plants to point of runoff or drip.
- Not for use in residential misting systems.

SPRAY DRIFT MANAGEMENT FOR AGRICULTURAL CROPS

Avoiding spray drift at the application site is the responsibility of the applicator and the grower. The interactions of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Do not apply at wind speeds greater than 10 mph at the application site.

Do not make any type of application into temperature inversions.

Apply as a medium or coarser spray (ASABE standard 572)

Additional Requirements for Aerial Applications:

Do not release spray at a height greater than 10 feet above the ground or crop canopy.

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Aerial applicators must consider flight speed and nozzle orientation in determining droplet size.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Additional requirements for ground applications:

Do not release spray at a height greater than 4 feet above the ground or crop canopy.

Additional requirements for airblast applications:

Direct sprays into the canopy

Turn off outward pointing nozzles at row ends and when spraying outer rows.

PESTICIDE RESISTANCE MANAGEMENT

PyGanic® Crop Protection EC 1.4 contains a Group 3A insecticide. Insect biotypes with acquired resistance to Group 3A may eventually dominate the insect population if Group 3A insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by PyGanic® Crop Protection EC 1.4 or other Group 3A insecticides.

To delay insecticide resistance consider:

- Avoiding the consecutive use of PyGanic® Crop Protection EC 1.4n or other group 3A insecticides that have a similar target site of action, on the same insect species.
- Using tank mixtures or premixes with insecticides from a different target site of action Group as long as the involved products are all registered for the same use and have different sites of action.
- Basing insecticide use on a comprehensive IPM program.
- Monitoring treated insect population for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for insecticide resistance management and or IPM recommendations for the specific site and resistance pest problems.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls:

Chemical-resistant gloves, such as Barrier Laminate, butyl rubber, nitrile rubber or viton.

Shoes plus socks.

Do not apply this product through any type of chemigation system.

USED ALONE: Apply 16 ounces (1 pint) to 64 ounces (2 quarts) per acre by ground in sufficient water for thorough coverage. This product may be applied by air at the rate of 16 to 64 ounces per acre in a minimum of 5 gallons of water. Mix only enough for immediate use. Spraying should begin when the insects first appear. Do not wait until the plants are heavily infested. Repeat as required to maintain effective control. It is recommended that the final spray mix be buffered to a pH of 5.5-7.0.

USED AS A TANK MIX: This product may be tank mixed with other insecticides and acaricides to provide a flushing of insects from hiding and into contact with other spray residues. This application should conform to accepted use precautions and directions for both products.

To avoid possible harm to honey bees, it is advisable to apply in the early morning or late evening hours.

GROWING CROPS (OUTDOORS AND IN GREENHOUSES):

ROOT AND TUBER VEGETABLES: Including (but not limited to) Arracacha, Arrowroot, Purple Arrowroot, Japanese Artichoke, Jerusalem Artichoke, Beets, Sugar Beets, Edible Burdock, Carrots, Cassava (bitter or sweet), Celeriac (celery root), Chervil (turnip rooted), Chicory, Chuffa, Dasheen, Ginger, Ginseng, Horseradish, Leren, Parsley (turnip rooted), Parsnip, Potato, Radish, Japanese Radish (Daikon), Rutabaga, Salsify, Black Salsify, Spanish Salsify, Sweet Potato, Tanier, Turmeric, Turnip, Yam (true), Yam Bean.

<u>LEAVES OF ROOT AND TUBER VEGETABLES</u>: Including (but not limited to) Beet, Sugar Beet, Edible Burdock, Carrot, Cassava (bitter or sweet), Celeriac (celery root), Chervil (Turnip Rooted), Chicory, Dasheen, Parsnip, Radish, Japanese Radish (Daikon), Rutabaga, Black Salsify, Sweet Potato, Tanier, Turnip, Yam (true).

<u>LEAFY VEGETABLES</u>: Including (but not limited to) Amaranth (Leafy Amaranth, Chinese Spinach, Tampala), Arrugula, Celery, Celtuce, Chervil, Cilantro, Corn Salad, Chrysanthemum (edible leaved), Chrysanthemum (gardland), Cress (garden), Upland Cress (yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Fennel (Florence), Lettuce (head and leafy), Orach, Parsley, Purslane (garden & winter), Rhubarb, Spinach, Fine Spinach (Malabar, Ceylon), Spinach (New Zealand), Swiss Chard.

BRASSICA (COLE) LEAFY VEGETABLES: Including (but not limited to) Broccoli, Chinese Broccoli (Gai Lon), Broccoli raab (Rapini), Brussels Sprouts, Cabbage, Chinese Cabbage (Bok Choy), Chinese Mustard Cabbage (Gai Choy), Cauliflower, Collards, Kale, Kohlrabi, Mustard Greens, Rape Greens.

LEGUME VEGETABLES (SUCCULENT OR DRIED): Including (but not limited to) Adzuki Beans, Field Beans, Kidney Beans, Lima Beans, Moth Beans, Mung Beans, Navy Beans, Pinto Beans, Rice Beans, Runner Beans, Snap Beans, Tepary Beans, Urd Beans, Wax Beans, Asparagus Beans, Black-eyed Peas, Catjang, Chinese Longbeans, Cowpeas, Chowder Peas, Southern Peas, Yard-Longbeans, Broad Beans, (Fava Beans), Chick Peas (Garbanzo Beans), Guar, Jackbean (Sword Bean), Lablab Bean (Hyacinth Bean), Lentils, Peas (Garden Peas, Field Peas, Sugar Peas), Pigeon Peas, Soybeans.

FOLIAGE OF LEGUME VEGETABLES: Including (but not limited to) Plant part of any legume vegetable included in the legume vegetable group that will be used as animal feed including any variety of Beans, Field Peas, Soybeans.

FRUITING VEGETABLES: Including (but not limited to) Eggplant, Ground Cherry, Okra, Pepinos, Pepper (Bell Pepper, Chili Pepper, Cooking Peppers, Pimentos, Sweet Peppers), Tomatillo, Tomatoes.

<u>CUCURBIT VEGETABLES</u>: Including (but not limited to) Balsam Pear (Bitter Melon), Chinese Waxgourd, Citron Melon, Cucumber, Gherkin, Edible Gourds, Melons (including hybrids, Cantaloupe, Casaba, Crenshaw, Honeydew Melons, Honey Balls, Mango Melon, Muskmelon, Persian Melon), Pumpkin, Squash (summer & winter), Watermelon (including hybrids).

<u>CITRUS FRUITS</u>: Including (but not limited to) Calamondin, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquats, Lemons, Limes, Mandarin (Tangerine), Orange (sweet & sour), Pummelo, Satsuma Mandarin.

POME FRUITS: Including (but not limited to) Apple, Crabapple, Loguat, Pear, Oriental Pear, Quince.

STONE FRUITS: Including (but not limited to) Apricot, Cherry (sweet & sour), Nectarine, Peach, Plum, Prune, Chickasaw Plum, Damson Plum, Japanese Plum.

SMALL FRUITS AND BERRIES: Including (but not limited to) Blackberry, Blueberry, Cranberry, Currant, Dewberry, Elderberry, Gooseberry, Grape, Huckleberry, Loganberry, Olallie Berry, Raspberry(black & red), Strawberry, Youngberry.

ORIENTAL VEGETABLES: Including (but not limited to) Japanese Artichoke, Chinese Broccoli (Gai Lan), Chinese Cabbage (Bok Choy,), Chinese Mustard Cabbage (Gai Choy), Dasheen, Ginger, Ginseng, Chinese Longbeans, Mung Beans, Citron Melon, Balsam Pear (Bitter Melon), Japanese Radish (Daikon), Chinese Spinach, Chinese Waxgourd.

ADDITIONAL CROPS: Asparagus, Onions (bulb and green), Rice.

ORNAMENTALS: Including (but not limited to) African Violet, Aster, Azalea, Begonia, Calceolaria, Calendula, Calla, Camellia, Carnation, Cineraria, Chrysanthemum, Cypress, Daffodil, Dahlia, Dogwood, Elm, Eucalyptus, Fern, Ficus, Geranium, Gladiolus, Gypsophila, Holly, Juniper, Lily, Marigold, Oak, Palm, Peony, Petunia, Philodendron, Pine, Roses, Snapdragon, Sweet Pea, Tulips, Viburnum, Wandering Jew, Yew, Zinnia.

FOR THE CONTROL OF INSECTS: Including (but not limited to) Ants, Aphids, Armyworms, Asparagus Beetle, Blister Beetles, Cabbage Looper, Caterpillars, Cockroaches, 12-spotted Cucumber Beetle, Colorado Potato Beetles, Corn Earworm, Crickets, Cross-striped Cabbageworm, Cucumber Beetles, Deer Fly, Diamondback Larvae, Fireworms, Flea Beetles, Fruit Flies, Fruittree Leafroller, Grape Leafhopper, Green Peach Aphids, Greenhouse Thrips, Gypsy Moth (adults & larvae), Harlequin Bug, Heliothis sp., Imported Cabbageworm, Leafhopper, Leafrollers, Leaftiers, Lice, Mexican Bean Beetle, Potato Leafhopper, Psyllids, Skippers, Stink Bugs, Thrips, Vinegar Flies, Webworms, and Whiteflies.

627

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

<u>PESTICIDE STORAGE</u>: Store in a cool, dry area, preferably under lock and key. Keep container closed. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

<u>PESTICIDE DISPOSAL</u>: To avoid wastes, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

<u>CONTAINER HANDLING</u>: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill.

Triple Rinse as follows [for containers of 5 Gallons (18.9 L) or less]: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple Rinse as follows [for containers greater than 5 Gallons (18.9 L)]: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Net Contents _____ Manufactured by: Mc LAUGHLIN GORMLEY KING COMPANY 8810 Tenth Avenue North Minneapolis, MN 55427

EPA Reg. No. 1021-1751

EPA Est. No. 1021-MN-2