

1021-1758

09/02/2003

1/10

27783-0603a z

# MGK<sup>®</sup> McLAUGHLIN EVERCIDE<sup>®</sup> 2.5 PERMETHRIN CONCENTRATE/REPELLENT

## GORMLEY KING COMPANY

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4319 U.S.A.

TELEPHONE (763) 544-0341

[Denotes alternate language]

For Use on Residential Lawns, Gardens, Patios and Home Picnic Areas Only

Controls (Kills) Listed Pests for up to 4 Weeks

Kills on Contact and Keeps Killing for up to 4 Weeks

Water Based Formula

Kills Bugs on Your Lawns

Trees and Shrubs

Roses and Flowers

Fruits and Nuts

Vegetables

Aids in Reducing Annoyance from Mosquitoes and Other Flying Insects

Kills Gnats, Fleas, Deer Ticks, Brown Dog Ticks and other Crawling insects

Repels Crawling Insects from your Home Recreation Areas

[May be used on indoor and outdoor plants]

[Controls over 50 common insect pests]

[Controls insects on: roses &amp; ornamentals; fruits, nuts &amp; vegetables; trees, shrubs; lawns]

[Controls nuisance pests: ants, spiders, fleas, ticks, crickets &amp; earwigs]

Contains Permethrin, a long lasting insecticide that provides contact and residual control for up to 4 weeks of many hard-to-kill sucking and chewing insects.

**ACTIVE INGREDIENT:**

- Permethrin [\* (3-phenoxyphenyl)methyl(±)-cis,trans-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate].....

2.50%

**OTHER INGREDIENTS** .....

97.50%

100.00%

- Cis,trans ratio: Min. 35% (±) cis and max 65% (±) trans

EVERCIDE<sup>®</sup> - Registered trademark of McLaughlin Gormley King Co.

KEEP OUT OF REACH OF CHILDREN

**CAUTION****FIRST AID****IF ON SKIN OR CLOTHING:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

May be harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish. Do not apply directly to water. Drifts and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are actively visiting the treatment area.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SEP - 2 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1021-1758

**SHAKE WELL BEFORE USE.**

This product helps reduce annoying and troublesome outdoor insects. Do not spray animals. People and pets should not be allowed in treated areas until spray has dried.

This product can be applied using a hose-end sprayer [such as product name], pump up pressure sprayer or by watering can. Carefully dilute the recommended amount of concentrate and water according to the directions found under the desired insect to control. Use a clean sprayer. Apply insecticide when insects first appear and repeat not more frequently than every 3 days. Flush sprayer or watering can with clean water after each use.

This product kills most troublesome outdoor insect, which destroy lawns, gardens, and are an annoyance on patios and at picnics.

**Use on Lawns:** Bent, Bermuda, Bluegrass, Dichondra, Fescue, Irish Moss, Merion and St. Augustine

**Pests:** Ants Armyworms, Brown dog ticks, Chinch bugs, Crickets, Cutworms, Deer ticks, Fleas, Grasshoppers, Japanese beetles grubs, Mole crickets and Sod webworms. Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For heavy infestations, repeat application after 2 weeks. For ants, crickets and grasshoppers use 2-fl. oz. In 3 gallons of water to cover 1,000 sq. ft. For other listed insects, use 6 fl. oz. in 10 gallons of water to cover 1,000 sq. ft.

[When using (product name) 1. Set dial to 1 oz. 2. Pour product into sprayer to fill jar one-quarter to one-half full. DO NOT add water. 3. After spraying, unused product can be poured back into its original container.]

**Use on Trees and Shrubs:** Arizona Cypress, Azalea, Birch, Non-Bearing Citrus, Conifers, Elm, English Ivy, Euonymus, Weeping Fig, Fir, Honeysuckle, Lilac, Mock-Orange, Oak, Palm, Philodendron, Pine, Poinsettia, Tulip, Poplar, Rhododendron, Taxus.

**Pests:** Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Inchworms, Leafminers, Mealybugs, Spider mites, Whiteflies. Fall webworms on Cherry and Honeysuckle, Budworms on Conifer, Elm leaf beetles, Elm spanworms and Fall webworms on Elms, Tussock moths on Douglas and Grand Firs, Citrus black flies on Non-Bearing Citrus, Oakworms on Oaks, Pine beetles, Pine moths, Pine needleminers, and Needle scales on Pines.

Use 1 fl. oz. in 1 gallon of water. Apply as a thorough spray, wetting leaves and branches but do not let runoff occur. Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50 to 75 F and when there is little or no wind. Spray at the first sign of insects. To maintain effectiveness, spray as reinfestation occurs, but use intervals of 4-8 days.

[When using (product name) 1. Set dial to 1 oz. 2. Pour product into sprayer to fill jar one-quarter to one-half full. DO NOT add water. 3. After spraying, unused product can be poured back into its original container.]

**Use On Roses and Flowers:** Ageratum, Aster, Azalea, Begonia, Coleus, Common Ninebark and Snowberry, Exacum, Gladiolus, Gold Bells, Hypoestes, Ivy, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Pansy, Pea Shrub, Petunia, Poinsettia, Rhododendron, Rose, Snapdragon, Zinnia.

**Pests:** Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Fall webworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider mites, and Whiteflies.

Use 1 fl.oz. in 1 gallon of water. Apply as a thorough spray, wetting the plants thoroughly, but do not let runoff occur. Try to hit underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening, when the temperature ranges from 50 to 75 F and when there is little or no wind. Spray at the first sign of insects. To maintain effectiveness, spray as reinfestation occurs, but use intervals of 4-8 days.

[When using (product name) 1. Set dial to 1 oz. 2. Pour product into sprayer to fill jar one-quarter to one-half full. DO NOT add water. 3. After spraying, unused product can be poured back into its original container.]

**Use on Fruit and Nut Trees:** Almonds, Apples, Peaches, and Pears.

General Directions: Use 2 fl. oz. in 1 gallon of water. Apply when insects appear. Wet the plants thoroughly, but do not let runoff occur. Try to hit underside of leaves and penetrate dense foliage.

[When using (product name) 1. Set dial to 1 oz. 2. Pour product into sprayer to fill jar one-quarter to one-half full. DO NOT add water. 3. After spraying, unused product can be poured back into its original container.]

**Almonds:**

**Pests:** Navel orangeworms, Peach twig borers.

**How to Use:** Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest.

**Apples:**

**Pests:** Green fruitworms, Leafrollers (Oblique-banded; Redbanded), Plum curculio, Rosy apple aphids, Spotted tentiform leafminers, Tarnished plant bugs, White apple leafhoppers.

**How to Use:** Repeat as required to maintain control. Do not make more than 3 applications. Do not apply after petal fall.

**Peaches:**

**Pests:** Borers (lesser peachtree, Peach twig), Green fruitworms, Plum curculio, Oriental fruit moths, Tarnished plant bugs.

**How to Use:** Do not apply within 7 days of harvest. Do not make more than 8 applications per season.

**Pears:**

**Pests:** Codling moth, Green fruitworms, Pear psylla.

**How to Use:** Do not make more than 3 applications per season. Can be applied up to 14 days prior to harvest.

**Use on Vegetables:** Asparagus (pre-harvest), Asparagus (post-Harvest), Celery and Cole Crops (Broccoli, Brussels Sprouts, Cabbage, Cauliflower), Cucumber, Melons, Pumpkins, Squash, Cucurbits, Eggplant, Horseradish, Head Lettuce, Leafy Vegetables, Potatoes, Peppers, Spinach, Sweet Corn, Tomatoes.

**Asparagus (pre-harvest):****Pests:** Asparagus beetles, cutworms**How to Use:** Use 1½ fl. oz. in one gallon of water to cover 1,000-sq. ft. Apply for uniform coverage as needed. Do not make more than 4 applications per crop. Can be applied up to 3 days prior to harvest.**Asparagus (post-harvest):****Pests:** Beetles (Adult Japanese Beetle, Larval stages of Asparagus Beetle), Lygus bugs, Tarnished plant bugs.**How to Use:** Use 1 ½ fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Treat ferns or bush growth after spear harvest when insects are present.**Celery:****Pests:** Armyworms (beet, Fall, Southern), Cabbage loopers, Cutworms (black, granulate), Vegetable leafminers.**How to Use:** Use 1½ fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 3-5 days or as needed. Can be applied up to 3 days prior to harvest. Do not make more than 10 applications per season.**Cole Crops: Broccoli, Brussels Sprouts, Cabbage, and Cauliflower.****Pests:** Beet armyworms, cabbage aphids and loopers, Diamondback moths, Imported cabbageworms.**How to Use:** Use 1 fl. oz. in 1 gallon of water to cover 1,000-sq. ft. Apply every 5 days or as needed. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts, and Broccoli and 10 applications on Cabbage.**Cucumber, Melons, Pumpkins, Squash:****Pests:** Flea Beetle, Leafhopper**How to Use:** Use 2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply as needed but not more than 8 applications per season. Can be applied up to day of harvest.**Cucurbits: Balsam Pear (Bitter Melon), Chinese Waxgourd (Chinese Preserving Melon), Citron Melon, Cucumber, Gherkin, Edible Gourds, Melons including hybrids such as Cantaloupe, Casaba, Crenshaw, Honeydew, Honey Balls, Mango Melon, Muskmelon, Persian Melon, Pumpkin, Summer & Winter Squash, Watermelon.****Pests:** Aphids, Cabbage, Looper, Cucumber Beetle (adults), Cutworms, Leafhoppers, Leafminers, Melonworms, Pickleworm, Plant Bugs, Squash Bugs.**How to Use:** Use 2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply as needed to ensure uniform coverage. Do not exceed 8 applications per season. Can be applied up to the day of harvest.**Eggplant:****Pests:** Colorado Potato beetles, Vegetable leafminers.**How to Use:** Use 3 fl. oz. in 1 gallon of water to cover 1,000-sq. ft. Apply for uniform coverage as needed. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest.**Horseradish (Foliar Application):****Pests:** Imported crucifer weevils**How to Use:** Use 3 fl. oz. in 1 gallon of water to cover 1,000-sq. ft. Apply as needed. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.**Head Lettuce:****Pests:** Corn earworms, Beet armyworms, Loopers (Alfalfa, Cabbage), Tobacco budworms**How to Use:** Use 2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 5-10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest.**Leafy Vegetables: Amaranth, Arugula, Celery, Celtuce, Chervil, Corn Salad, Chrysanthemum (Edible-leaved and Garland), Cress (Garden and 'Iplant), Dandelion, Dock, Endive, Fennel, Lettuce (Head and Leaf), Okra, Parsley, Purslane (Garden and Winter), Rhubarb, Spinach, Swiss Chard.****Pests:** Alfalfa Looper, Aphids, Beet Armyworm, Cabbage Looper, Corn Ear Worm, Cutworms, European Corn Borer, Fall Armyworm, Green Cloverworm, Southern Armyworm, Tobacco Budworm, Vegetable Leaf Miner.**How to Use:** Use 2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 5-10 days as needed. Do not exceed 10 applications per season. Can be applied up to 1 day prior to harvest.**Melons: - see Cucurbits****Potatoes:****Pests:** Beetles (Colorado potato, Potato flea), Beet armyworms, Cabbage loopers, Cutworms, European corn borers, Leafhoppers (Aster. Potato), Potato aphids, Potato Psyllids, Potato tuberworms, Tarnished plant bugs.**How to Use:** Use 1½ fl. oz. in 1 gallon of water to cover 1,000-sq. ft. Apply as needed. Do not make more than 12 applications. Can be applied up to 7 days prior to harvest.**Peppers:****Pests:** Cabbage loopers, Corn earworms, European corn borers, Flea beetles, Pepper weevils, Vegetable leafminers.**How to Use:** Use 2 fl. oz. in 1 gallon of water to cover 1,000-sq. ft. Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest.**Spinach:****Pests:** Beet armyworms, Cabbage loopers, Corn earworms, Cutworms, European corn borers, Green cloverworms, Leafhoppers.**How to Use:** Use 1 1/2 fl. oz. in 1 gallon of water to cover 1,000-sq. ft. Apply for uniform coverage when needed, but not more than 7 applications per season. Can be applied up to 7 days prior to harvest.**Sweet Corn:****Pests:** Armtworms, (Fall, Southern), Corn earworms, European corn borers.**How to Use:** Use 1 1/2 fl. oz. in 1 gallon of water to cover 1,000-sq. ft. Apply every 5 days and when needed, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.

**Tomatoes:**

**Pests:** Beet and Southern Armyworm, Tomato Pinworm, Tomato Fruitworm, Horn Worms, Granulate Cutworms, Colorado Potato Beetle, Cabbage Looper.

**How to Use:** Use 1 ½ fl. oz. in 1 gallon of water per 1,000 sq. ft. Apply as needed for uniform coverage. Do not exceed 8 applications per season. Can be applied up to day of harvest.

**Barrier Spray For Nuisance Pests:** Such as Ants, Clover mites, Cluster flies, Crickets, Earwigs, Fleas, Flying moths, Gnats, House flies, Millipedes, Mosquitoes, Sowbugs (pillbugs), Spiders, Tent caterpillars, Ticks (Brown dog, Deer, Lone star).

**General Directions:**

Use 6-2/3 fl. oz. In 1 quart of water, treating 400 linear feet with a 6-inch spray band.

**Outside Surfaces of Buildings:**

**Pests:** Ants, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs).

**How to Use:** To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, picnic areas, garages, refuse piles, and other areas where these pests congregate or have been seen. To maintain effectiveness, reapply as reinfestation occurs.

**Other Outside Areas:**

**Pests:** Ants, Crickets, Earwigs, Fleas, Spiders, Ticks (Brown dog, Deer, Lone star).

**How to Use:** For treatment of localized infestations of these insects in areas where there are weeds, shrubs or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as reinfestations occur.

**Pests:** Cluster flies, Flying moths, Gnats, House flies, Mosquitoes.

**How to Use:** For use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, under patio furniture, in garages and other places where these insects alight or congregate.

**Pests:** Tent caterpillars.

**How to Use:** Application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent, however, they should die within several hours.

**House Plants Indoors:** To protect plants in non-commercial greenhouses and lathhouses, and for indoor landscaping. Ageratum, Ardisia, Aster, Baby's Breath, Begonia, Birdsnest Fern, Bleeding Heart, Carnation, Chrysanthemum, Coleus, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Fuchsia, Gladiolus, Grape Ivy, Hypoestes Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia.

**Pests:** Aphids, Armyworms, Corn earworms, Cutworms, Cabbageworms, Exposed thrips, Fall cankerworms, Fungus gnats, Inchworms, Japanese beetles, Leafminers, Leafrollers, Mealybugs, Scales, Spider mites, Spring cankerworms, Whiteflies.

**How to Use on Indoor House Plants Listed Above:** Use 1 fl. oz. in 1 gallon of water. Spray will act as contact and residual control. Use sufficient spray to obtain full coverage of all foliage. Adjust spray volume used to plant size. Spray directly on insects. If application is to be made indoors, remove infested plants from living areas prior to application. Spray plants in an area not likely to be occupied by people or pets (particularly fish bowls or tanks). When leaves dry, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury. Use with care on Ferns, Palms and Fuchsia. As with all plants listed above, especially carnations, chrysanthemums, and roses, there exists a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury. Repeat application may be necessary to control citrus blackflies on non-bearing citrus and adult Japanese beetles on roses.

**Product with Hose End Sprayer Attached:**

Make sure knob is twisted fully clockwise to OFF position. Shake well before attaching to hose. Turn Product Control Button clockwise until its point aligns with triangle, then push all the way in. Avoid squeezing bottle. Turn water on at faucet, aim nozzle toward lawn and turn knob counter clockwise to begin spraying. When finished push Product Control Button to OFF position from opposite side. Return knob and faucet to OFF position. Discharge water pressure by turning knob ON and OFF again. Turn Product Control Button counter clockwise away from triangle to prevent accidental discharge. Store out of reach of children.

**STORAGE AND DISPOSAL**

**STORAGE:** Store in a cool, dry place, preferably a locked storage area. Store above freezing. Do not store diluted spray. Store with container tightly closed. Keep pesticide in original container; do not transfer to food or drink container. Avoid contamination of food, feed, and domestic water supplies.

**CONTAINER DISPOSAL:**

If empty:	Do not reuse this container. Place in trash or offer for recycling if available.
If partly filled:	Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NET CONTENTS**

Manufactured by:  
**Mc LAUGHLIN GORMLEY KING COMPANY**  
 8810 Tenth Avenue North  
 Minneapolis, MN 55427  
 Made in U.S.A

**EVERCIDE® 2.5 Permethrin Concentrate/Repellent**

[Indicates Optional Text]

Ready-Spray®

- **Aids in Reducing Annoyance From Mosquitoes and Other Flying Insects**  
*[INSECT ART, mosquito]*

[Treats up to 16,000 sq. ft.]  
[Kills chinch bugs, mole crickets, plus other listed insects! See back.]  
[Kills ants, armyworms, beetles, caterpillars, plus others listed! See back.]  
[Kills on contact and keeps killing for up to 4 weeks]

Active Ingredient	
Permethrin*.....	2.50%
Other Ingredients. ....	<u>97.50%</u>
	100.00%

\* Cis, trans ratio: Min. 35% (+/-) cis and max 65% (+/-) trans

For Use on Residential Lawns, Gardens, Patios and Home Picnic Areas Only

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panel booklet for additional precautionary statements.

**NET 32 FL OZ (1 QT) 946 mL**

[BACK]

**Ready-Spray®**

- Contains Permethrin, a long lasting insecticide that provides contact and residual control [kill] for up to 4 weeks of many hard-to-kill sucking and chewing insects.

<b>PRODUCT FACTS</b>	
<b>KILLS INSECTS</b>	<p>Lawn pests: ants, armyworms, brown dog ticks, chinch bugs, crickets, cutworms, deer ticks, fleas, grasshoppers, Japanese beetle grubs, mole crickets and sod webworms..</p> <p>Tree, shrub, rose &amp; flower pests: aphids, bagworms, cicadas, exposed thrips, fall cankerworms and other listed pests.</p> <p>Nuisance pests (outdoors): ants, clover mites, cluster flies, crickets, earwigs, fleas and other listed pests.</p>
<b>WHERE TO USE</b>	On lawns, trees and shrubs, roses and flowers and outside surfaces of buildings.
<b>[PHONE/ COMPUTER ICON]</b>	<p>Questions, Comments or Medical Information?</p> <p>Call 1-800-225-2883</p> <p><a href="http://www.ortho.com">www.ortho.com</a></p>

EPA Reg. No. 1021-1758-

EPA Est.

Superscript is first letter of lot number.

Made in USA

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product helps reduce annoying and troublesome outdoor insects. Do not spray animals. People and pets should not be allowed in treated areas until spray has dried.

- Apply insecticide when insects first appear and repeat as necessary but not more frequently than every 3 days.

This product kills most troublesome outdoor insects, which destroy lawns, gardens, and are an annoyance on patios and at picnics.

<b>HOW TO USE THE Ready-Spray® SYSTEM</b>	
<b>CONNECT</b>	<ol style="list-style-type: none"> <li>1. Shake well before using.</li> <li>2. Connect sprayer to hose.</li> <li>3. Turn on water.</li> </ol>
<b>SPRAY</b>	<ol style="list-style-type: none"> <li>1. To begin spraying, point nozzle in the direction you want to spray.</li> <li>2. Bend small yellow plastic tab back and turn knob clockwise to ON position.</li> <li>3. Spray evenly over measured area.</li> </ol>
<b>FINISH</b>	<ol style="list-style-type: none"> <li>1. To stop spraying, turn knob counter-clockwise to OFF position.</li> <li>2. Turn off water.</li> <li>3. Relieve water pressure by bending yellow plastic tab back and turning knob to ON position until water slows to a drip. Then, turn knob back to OFF position.</li> <li>4. Disconnect sprayer from hose.</li> </ol>

**Barrier Spray For Nuisance Pests:** Such as Ants, Clover mites, Cluster flies, Crickets, Earwigs, Fleas, Flying moths, Gnats, House flies, Millipedes, Mosquitoes, Sowbugs (pillbugs), Spiders, Tent caterpillars, Ticks (Brown dog, Deer, Lone Star)

**General Directions:**

Use 32 fl oz. to treat 1,900 linear feet with a 6-inch spray band.

<b>OTHER OUTSIDE AREAS</b>	
<b>Pests:</b> Ants, Crickets, Earwigs, Fleas, Spiders, Ticks, (Brown dog, Deer, Lone star)	<b>How to Use:</b> For treatment of localized infestations of these insects in areas where there are weeds, shrubs or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as reinfestations occur.
<b>Pests:</b> Cluster flies, Flying moths, Gnats, House flies, Mosquitoes	<b>How to Use:</b> For use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, under patio furniture, in garages and other places where these insects alight or congregate. Repeat as necessary.
<b>Pest:</b> Tent caterpillars	<b>How to Use:</b> Application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent, however, they should die within several hours..

Outside Surfaces of Buildings	<b>Pests:</b> Ants, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs).	<b>How to Use:</b> To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, picnic areas, garages, refuse piles, and other areas where these pests congregate or have been seen. [To maintain effectiveness, reapply as reinfestation occurs.] [Repeat as needed to maintain effectiveness.]
-------------------------------	--	---

### USE ON LAWNS

Bent, Bermuda, bluegrass, dichondra, fescue, Irish moss, Merion and St. Augustine	<b>Pests:</b> Ants Armyworms, Brown dog ticks, Chinch bugs, Crickets, Cutworms, Deer ticks, Fleas, Grasshoppers, Japanese beetles grubs, Mole crickets and Sod webworms.	<b>How to Use:</b> Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For heavy infestations, repeat application after 2 weeks. For ants, crickets and grasshoppers, 32 fl oz treats 16,000 sq ft. For other listed insects, 32 fl oz. treats 5,300 sq ft.
---	--	--

### USE ON TREES & SHRUBS

Arizona Cypress, Azalea, Birch, Non-Bearing Citrus, Conifers, Elm, English Ivy, Euonymus, Weeping Fig, Fir, Honeysuckle, Lilac, Mock-Orange, Oak, Palm, Philodendron, Pine, Poinsettia, Tulip, Poplar, Rhododendron, Taxus	<b>Pests:</b> Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Inchworms, Leafminers, Mealybugs, Spider mites, Whiteflies. Fall webworms on Cherry and Honeysuckle, Budworms on Conifer, Elm leaf beetles, Elm spanworms and Fall webworms on Elms, Tussock moths on Douglas and Grand Firs, Citrus black flies on Non-Bearing Citrus, Oakworms on Oaks, Pine beetles, Pine moths, Pine needleminers, and Needle scales on Pines.	<b>How to Use:</b> Apply as a thorough spray, wetting leaves and branches but do not let runoff occur. Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50 to 75 F and when there is little or no wind. Spray at the first sign of insects. To maintain effectiveness spray as reinfestation occurs, but use intervals of 4-8 days.
--	---	---



**USE ON ROSES & FLOWERS**

Ageratum, Aster, Azalea, Begonia, Coleus, Common Ninebark and Snowberry, Exacum, Gladiolus, Gold Bells, Hypoestes, Ivy, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Pansy, Pea Shrub, Petunia, Poinsettia, Rhododendron, Rose, Snapdragon, Zinnia.	<b>Pests:</b> Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Fall webworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider mites, and Whiteflies.	<b>How to Use:</b> Apply as a thorough spray, wetting the plants thoroughly, but do not let runoff occur. Try to hit underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening, when the temperature ranges from 50 to 75 F and when there is little or no wind. Spray at the first sign of insects. To maintain effectiveness; spray as reinfestation occurs but use intervals of 4-8 days.
--	--	---

**STORAGE AND DISPOSAL**

**STORAGE:** Store in a cool, dry place, preferably a locked storage area. Store above freezing. Do not store diluted spray. Store with container tightly closed. Keep pesticide in original container; do not transfer to food or drink container. Avoid contamination of food, feed, and domestic water supplies.

**CONTAINER DISPOSAL:**

If empty: Do not reuse this container. Place in trash or offer for recycling if available.  
If partially filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.  
Never place unused product down any indoor or outdoor drain.

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

May be harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**FIRST AID****IF ON SKIN OR CLOTHING:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents call 1-800-225-2883.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish. Do not apply directly to water. Drifts and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

**NOTICE:** Buyer assumes all risk of use, storage or handling of this product not in accordance with directions.

**The ORTHO Guarantee**

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

*[phone icon]* **Questions, Comments or Medical Information ?**

**call 1-800-225-2883**

**[www.ortho.com](http://www.ortho.com)**

© 2002, The ORTHO Group. World Rights Reserved.

EVERCIDE®-Registered trademark of McLaughlin Gormley King Co.

EPA Reg. No. 1021-1758-

EPA Est.

Superscript is first letter of lot number.

Made in USA