



McLAUGHLIN  
GORMLEY  
KING COMPANY

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4372 U.S.A. - Telephone (763) 544-0341

Kills Flies & Lice  
Insecticide For Cattle, Sheep and Their Premises  
Synergized Formula For Effective Control

Specially Formulated for Weekly Treatments  
POUR ON - SPRAY - MIST - BACK RUBBER - PREMISE

ACTIVE INGREDIENTS:

- \* Permethrin [(3-Phenoxyphenyl) methyl (+ or -) cis-trans (2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate)..... 0.50%
  - \*\* Piperonyl Butoxide Technical ..... 1.00%
  - \*\*\* OTHER INGREDIENTS ..... 98.50%
- 100.00%
- \* Cis-trans isomers ratio: Min. 35% (+ or -) cis and Max. 65% (+ or -) trans
  - \*\* Equivalent to Min. 0.8% butylcarbityl(6-propylpiperonyl) ether and 0.2% related compounds
  - \*\*\* Contains petroleum distillates

EVERCIDE® - Registered trademark of McLaughlin Gormley King Company

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

**ACCEPTED**

MAY 11 2005

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under 1021-1783  
EPA Reg. No.

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION**

Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and other aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

PHYSICAL OR CHEMICAL HAZARDS

**Combustible.** Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply the product only as specified on the label.

READY TO USE - No dilution necessary.

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> <li>▪ Immediately call a poison control center or doctor.</li> <li>▪ Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>▪ Do not give any liquid to the person.</li> <li>▪ Do not give anything by mouth to an unconscious person.</li> </ul>
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> <li>▪ Take off contaminated clothing.</li> <li>▪ Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>▪ Call a poison control center or doctor for treatment advice.</li> </ul>
IF INHALED:	<ul style="list-style-type: none"> <li>▪ Move person to fresh air.</li> <li>▪ If person is not breathing, call 911 or an ambulance. Then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>▪ Call a poison control center or doctor for further treatment.</li> </ul>
NOTE TO PHYSICIAN: Contains petroleum distillate – vomiting may cause aspiration pneumonia. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.	

Net Contents \_\_\_\_\_  
Manufactured by:  
Mc LAUGHLIN GORMLEY KING COMPANY  
8810 Tenth Avenue North  
Minneapolis, MN 55427

EPA Reg. No. 1021-1783

EPA Est. No. 1021-MN-2

Apply To:	Target Insects	Application Instructions
Lactating and Non-Lactating Dairy Cattle, Beef Cattle and Calves	Lice, Horn Flies, Face Flies; Aids in control of Horse Flies, Stable Flies, House Flies, Mosquitoes and Black Flies	DOSAGE: Apply 1/2 fl. oz. (15cc) per 100 lbs. body wt. of animal up to a maximum of 5 fl. oz. for any one animal POUR ON: Pour correct dose along back and down face. READY TO USE SPRAY: Use undiluted in a mist sprayer to apply correct dose. Apply directly to neck, face, back, legs and ears. BACK RUBBER USE: Mix one pint per gallon #2 diesel or mineral oil. Keep rubbing device charged. Results improved by daily forced use.
Sheep****	Sheep Keds****, Lice****	Pour along back. Apply 1/4 fl. oz. (7.5cc) per 50 lbs. body wt. of animal, up to a maximum of 3 fl. oz. for any one animal. For optimum control, all animals in the flock should be treated after shearing Sheep.

1021-1783

05/11/2005

1/2

Premises - in around Horse Dairy, Swine, Sheep and Poultry premises. Animal Hospital Pens and Kennels and "outside" Meat Processing Premises.	House Flies, Stable Flies, Face Flies, Gnats, Mosquitoes, Black Flies, Fleas, and Little Houseflies ( <i>Fannia spp.</i> ). Aids in control of Ticks, Cockroaches, Ants, Spiders, and Crickets.	For use as a ready-to-use spot spray or premise spray, use undiluted in a mist sprayer. Apply directly to surface to leave a residual insecticidal coating, paying particular attention to areas where insects crawl or alight. One gallon will treat approximately 7,300 square feet.
--	--	---

Apply 1/2 fl. oz. per 100 lbs. body weight of animal, up to a maximum of 5 fl. oz. for any one animal to kill Lice Horn Flies and Face Flies. Aids in the insecticidal control of Horse Flies, Stable Flies, Mosquitoes and Black Flies on lactating and non-lactating dairy cattle, beef cattle, calves and sheep. For sheep, apply 1/4 fl. oz. per 50 lbs. body weight of animal, up to a maximum of 3 fl. oz. for any one animal

**Pour-On:** Pour along back line and down face. For Sheep Keds\*\*\*\* or Lice\*\*\*\*, pour along back line over neck, shoulders and rump.

**Ready-To-Use Spray:** Use undiluted in a mist sprayer to apply correct dose. Apply directly to neck, face, back, legs and ears.

For Pour-On and Ready-To-Use Spray applications, repeat treatment as needed but do not exceed 5 fl. oz. for any one animal on cattle or 3 fl. oz. for any one animal on sheep during any 7-day interval. For optimum insecticidal effectiveness against lice, a second treatment at 7-day intervals is recommended.

**Back Rubber Use:** Mix 1 pint per gallon #2 diesel or mineral oil. Keep rubber device charged. For Back Rubber applications, results are improved by daily forced use.

For cattle and sheep, repeat treatment as needed, but not more often than once every week. For optimum lice control, two treatments at a 7-day interval are recommended.

**Special Note:** EVERCIDE® Synergized Permethrin Pour-On 2803 [this product] is not effective in controlling cattle grubs. EVERCIDE® Synergized Permethrin Pour-On 2803 [this product] is an oil-based, ready-to-use product that may leave an oily appearance on the hair/coat of some animals. EVERCIDE® Synergized Permethrin Pour-On 2803 [this product] should be used in an integrated pest management system which may involve repeat treatments and the use of other pest control practices.

\*\*\*\*Not registered in the State of California for the control of Sheep Keds and Lice in Sheep.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Keep container sealed when not in use. Do not store near food or feed.

#### DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.  
If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.