

1021-1796

6/11/2004

1/5

74402-404

PYROCIDE® GENERAL PURPOSE AQUEOUS SPRAY 74402

MGK McLAUGHLIN GORMLEY KING COMPANY

8810 Tenth Avenue N., Minneapolis, Minnesota 55412-4519 U.S.A.

TELEPHONE (763) 544-0341

[] or () Indicates alternate or optional wording

Contains the botanical insecticide, Pyrethrum [an extract of a Chrysanthemum flower]
 A micro-emulsion, water-base spray
 Kills [controls] and repels Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Mosquitoes, and Gnats
 Kills Black Widow Spiders and Black Carpenter Ants
 Quick killing pyrethrum controls crawling and flying insects in homes, schools, hospitals and other listed areas
 Controls Fleas and Ticks indoors and around pet areas
 Controls Deer Ticks and other Ixodid species that may carry and transmit Lyme Disease
 For use in Animal Quarters, Dairies, Milk Rooms, Milking Parlors, Calving Areas, Dairy and Hog Barns
 For use on Beef Cattle, Dairy Cattle, Horses, Hogs, Poultry and other listed livestock
 Kills Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus

ACCEPTED
 JUN 1 2004
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 1021-1796

ACTIVE INGREDIENTS:

Pyrethrins	0.10%
* Piperonyl butoxide, Technical	1.00%

OTHER INGREDIENTS	98.90%
	100.00%

*Equivalent to min. 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds
 PYROCIDE® is a registered trademark of McLaughlin Gormley King Company

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by a poison control center or doctor • Do not give anything by mouth to an unconscious person.
IF ON SKIN	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.	

PRECAUTIONARY STATEMENTS	
CAUTION	
<u>HAZARDS TO HUMANS AND DOMESTIC ANIMALS</u>	
Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Avoid contamination of food and feedstuffs. Remove pets, birds, and cover fish aquariums before spraying.	
<u>ENVIRONMENTAL HAZARDS</u>	
This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate.	
<u>PHYSICAL OR CHEMICAL HAZARDS</u>	
Do not use this product in or on electrical equipment because of shock hazard.	

[Pyrocide® General Purpose Aqueous Spray 74402] contains the quick-killing botanical insecticide, Pyrethrum, [an extract of Chrysanthemum flowers], which provides rapid, effective control of insects upon direct contact, with the added benefit of flushing [repelling] these insects to aid in more complete control [kill]. This highly effective water-based formula can be used on flying and crawling insects indoors and outdoors.

[Pyrocide® General Purpose Aqueous Spray 74402] can be used to control fleas and ticks hidden in carpets, rugs, upholstery, draperies, floor cracks and cabinets. Also controls Deer Ticks and other Ixodid species that may carry and transmit Lyme Disease.

[Pyrocide® General Purpose Aqueous Spray 74402] can be used in homes, schools, hospitals, restaurants and other listed areas to control [kill], on contact, listed crawling pests and is a fast, effective product for flying insect control. Pyrethrum has quick-killing ability with no persistent residue.

[Pyrocyde® General Purpose Aqueous Spray 74402] can be used to reduce annoyance of mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, outdoors around homes, recreational and other listed areas. Ideal for use prior to outdoor activities.

[Pyrocyde® General Purpose Aqueous Spray 74402] can be used in food and non-food areas of food-processing plants, canneries, industrial installations, bakeries, rice and wheat mills, tobacco warehouses, grain elevators, warehouses, storage sites and other listed areas to control [kill], on contact, listed crawling pests and is a fast, effective product for flying insect control.

[Pyrocyde® General Purpose Aqueous Spray 74402] provides rapid kill of insects without objectionable residues.

FOR THE CONTROL OF INSECTS including but not limited to:

- | | | | | |
|---|----------------------|---------------------------|--|---------------------|
| Black Fly | Ants | Cadelles | Lesser Grain Borer | Blood Sucking Lice |
| Deer Flies | Bedbugs | Carpet Beetles | Trogoderma Beetles | Lice |
| Face Flies | Black Carpenter Ants | Cereal Beetles | Almond Moths | Poultry Lice |
| Fannia Flies | Box Elder Bugs | Cigarette Beetles | Angoumois Grain Moths | Granary Weevils |
| Fruit Flies | Cockroaches | Confused Flour Beetles | Chocolate Moths | Rice Weevils |
| Gnats | Crickets | Darkling Beetles | Clothes Moths | Cheese Skipper |
| Horn Flies | Earwigs | Dermeid Beetles | Indian Meal Moths | Dark Mealworms |
| Hornets | Firebrat | Dried Fruit Beetles | Med. Flour Moths | Yellow Mealworms |
| Horse Flies | Millipedes | Drugstore Beetles | Miller Moths | Black Widow Spiders |
| House Flies | Palmetto Bugs | Flat Grain Beetles | Small Flying Moths | Cellar Spiders |
| Lesser House Flies | Pillbugs | Khapra Beetles | Tobacco Moths | Cheese Mites |
| Midges | Silverfish | Merchant Grain Beetles | Cat Fleas | Grain Mites |
| Skipper Flies | Sowbugs | Red Flour Beetles | Dog Fleas | Hornets |
| Stable Flies | Waterbugs | Rice Flour Beetles | Brown Dog Ticks | Wasps |
| Mosquitoes- including the Culex species that may carry and transmit the West Nile Virus | | Rusty Grain Beetles | Sheep Ticks | Yellow-Jackets |
| | | Spider Beetles | Deer Ticks | |
| | | Warehouse Beetles | Ticks that may carry and transmit Lyme Disease | |
| | | Saw-toothed Grain Beetles | | |

INDOOR AREAS for use in but not limited to:

- | | | | | |
|--------------------------|-------------------------|----------------------------|--------------------------|---------------------------|
| Animal Areas | Condominiums | Grain Storage | Milk Sheds | Solariums |
| Animal Quarter | Crawl Spaces | Granaries | Milking Rooms | Sport Stadiums |
| Animal Shelters | Credeenza | Grocery Stores | Mortuaries | Stables |
| Animal Control Centers | Crematoriums | Homes | Motels | Stable Bins |
| Apartments | Dairy, Livestock Barns | Horse Stables | Movie Theaters | Storage Areas |
| Atriums | Dairies | Hog Barns | Museums | Stored Products |
| Attics | Day Care Centers | Horse Stables & Barns | Nursing Homes | Stored Food, bags |
| Auditoriums | Delicatessens | Horse Trailers | Office Buildings | Stores |
| Automobiles | Dens | Hospices | Other Public Buildings | Sun Porch |
| Bakeries | Department Stores | Hospitals | Outdoor Eating Buildings | Sun Room |
| Bars | Diners | Hotels | Pantry | Swine Houses |
| Basements | Dog Houses | Household contents | Parking Garages | Taverns |
| Bathrooms | Dog Sleeping Quarters | Houses | Pet Bedding | Tents |
| Bedding | Dormitories | Indoor Eating Buildings | Pet Carriers | Textile Mills, Warehouses |
| Bedrooms | Draperies | Industrial Installations | Pet Grooming Parlors | Theaters |
| Beef Barns | Dressers | Industrial Plants | Pet Sleeping Areas | Tobacco Factories |
| Beverage Plants | Dried Fruits in Storage | Institutional Dining Areas | Pet Stores | Tobacco Warehouses |
| Boats | Drive-Ins | Institutions | Play Houses | Tool Sheds |
| Bookcases | Drugstores | Jails | Play Rooms | Trailers |
| Bottling Plants | Dumpsters | Kennels | Pool Halls | Trains |
| Breweries | Dwellings | Kitchens | Porches | Transportation Vehicles |
| Brewery Warehouses | Egg Processing Plants | Laboratories | Poultry Houses | Tree Houses |
| Building Housing Animals | Fabric Storage Areas | Laundromats | Poultry Plants | Tree Nuts Storage Areas |
| Building Housing Poultry | Factories | Lavatories | Pounds | Trucks |
| Buses | Farm Structures | Libraries | Prisons | Truck Trailers |
| Cabinets | Floors | Liquor Stores | Public Storage | Tunnels |
| Cabins | Food-processing Plants | Livestock Barns | Quonset Huts | Utility Rooms |
| Cafeterias | Food Storage Areas | Livestock Hauling Equip | Rabbit Processing Plants | Utilities |
| Calving Barns | Freight Containers | Livestock Holding Areas | Railroad Cars | Vending Machines |
| Campers | Fruit Packing Sheds | Living Rooms | Recreational Vehicles | Verandas |
| Canneries | Funeral Parlors | Loafing Shed | Research Animal Quarters | Veterinaries |
| Carpets | Furniture | Locker Rooms | Restaurants | Voids in Attics |
| Cat Sleeping Quarters | Garages | Marinas | Retail Stores | False Ceilings |
| Cattle Barns | Garbage Bins | Mattresses | Rice Mills | Warehouses |
| Chests | Garbage Cans | Mausoleums | Rooms | Washrooms |
| Churches | Garbage Compactors | Meat Packing Plants | Rugs | Wheat Mills |
| City Missions | Gas Stations | Mess Halls | Schools | Wholesale Stores |
| Circus Tents | Gazebos | Mini Storage | Seed Storage | Wine Cellars |
| Closed Porches | Goat Holding Areas | Mobile Canteens | Sheep Holding Areas | Wineries |
| Closets | Grain Elevators | Mobile Homes | Ships | Zoos |
| Clothes Storage | Grain Harvesting Equip | Morgues | Ships Holds | |
| Club Houses | Grain Handling Equip | Milk Houses | Smoke Shops | |
| Commercial Buildings | Grain Mills | Milk Rooms | | |

OUTDOOR AREAS for uses in and around, but not limited to:

- | | | | | |
|-----------------|--------------------|---------------|-----------------------|---------------|
| Alleys | Circuses | Game Fields | Outdoor Eating Estab. | Side Yards |
| Amphitheaters | Construction Sites | Garbage Dumps | Parks | Side Walks |
| Amusement Parks | Corrals | Gazebos | Parking Lots | Stadiums |
| Ant Hills | Courtyards | Grandstands | Patios | Tennis Courts |

Back Yards	Curbs	Junk Yards	Playgrounds	Truck Stops
Balconies	Decks	Kennels	Ranges	Underpasses
Baseball Fields	Drive-In Restaurants	Lanais	Recreational Areas	Utility Room Outdoor
Beaches	Drive-In Theaters	Levees	Rest Areas	Weigh Stations
Bridges	Dumpsters	Manholes	Roadsides	Wild Animal Parks
Carwashes	Fair Grounds	Manure Piles	Rooftops	Wood Piles
Campgrounds	Feedlots	Marinas	Sanitary Sewers	Zoos
Camp Sites	Front Yards			

FOR USE ON but not limited to:

Beefalo	Chickens	Lambs	Pigs	Swine
Birds	Donkeys	Llamas	Pullets	Turkeys
Buffalo	Ducks	Mules	Quail	Exotics such as
Calves	Emus	Ostrich	Ratites	European
Cattle- Beef	Geese	Oxen	Roosters	Red Deer
Cattle- Dairy	Horses	Pheasants	Sheep	

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read all directions completely before use. For best results, follow directions for specific use areas. Do not use this product in or on electrical equipment due to possibility of shock.

GENERAL:

The use of this product in food-processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food-processing surfaces should be covered during treatment or thoroughly cleaned before using.

For maximum effectiveness, follow the dilution instructions and use a combination of spot, surface, and fogging treatments.

SPACE SPRAYING: Use with conventional mechanical or compressed air equipment to kill exposed accessible stages of, but not limited to: Flies (including Fruit Flies), Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Small Flying Moths, Fleas, Gnats, Wasps, Hornets, Clover Mites, Cheese Mites, Cheese Skippers, Box Elder Bugs, Earwigs, Ants and the following common Stored Product Pests: Granary Weevils, Rice Weevils, Confused Flour Beetles, Saw-Toothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths, Tobacco Moths, Yellow Meal Worms, Dark Meal Worms, Grain Mites, Cadelles, Red Flour Beetles and other listed insects. Close room and shut off air conditioning or ventilating equipment. Apply at the rate of 1 to 1-1/2 fluid ounces per 1000 cubic feet of space. Direct spray toward ceiling and upper corners of room. Leave room closed for 1/2 hour and ventilate thoroughly before re-entry.

SURFACE SPRAYING: To kill Cockroaches, Ants, Silverfish, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Saw-toothed Grain Beetles, Spiders Beetles, Cigarette Beetles and other listed insects, use a sprayer adjusted to a coarse, wet spray. Mix well and apply as a coarse wet spray using approximately 1 gallon of spray mixture per 750 square feet of surface. Direct spray first into all cracks and crevices, and then behind all equipment, cupboards, obstructions or dark harborage areas, walls, floors, underneath sinks, behind pipes, beneath stoves and refrigerators, around garbage cans, cabinets, along baseboards, door sills, window sills, door and window frames and floors, around and on drains, pipes, plumbing, behind bookcases, storage and other utility installation areas and in all places that harbor insects to provide maximum penetration and impingement on the insects. Contact as many insects as possible. Do not allow spray to contact stored food. To kill ants, treat doors and around window frames and other areas of entry. Spray on ants whenever possible.

CLOTHES MOTHS: Remove any infested articles from storage. Brush thoroughly and air for several hours in sunlight, if possible. Apply liberally to empty chests, closets, bureaus and other storage areas, directing the spray into cracks, joints, and crevices.

CARPET BEETLES: Directly spray as many insects as possible. Spray all surfaces of upholstered furniture. Thoroughly spray both top and bottom of rugs and carpets when possible. Also treat shelves, closets, floors, walls, windowsills, baseboards, areas under furniture and wherever these insects are seen or suspected.

BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, floors, walls, furniture and other infested areas.

FOR USE IN LISTED PET AREAS: Remove, destroy or wash pet's old bedding. To reduce infestation, thoroughly spray animal's bedding, sleeping quarters, kennel, around baseboards, window and door frames, wall cracks, local areas of floors, and other resting places until the surface is slightly moist. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 18 inches. Also spray the entire inside surface of doghouses. Put fresh bedding in pet's quarters after spray has dried. Allow treated areas to dry thoroughly before allowing pets to re-enter.

FLEA AND TICK INFESTATION IN RUGS AND CARPETS: Prior to use, vacuum all infested areas thoroughly and dispose of vacuum bag in outside trash. Spray rugs and carpets where fleas tend to hide, usually where animals frequent. Spray thoroughly into hiding places such as cracks and crevices, in walls and along floors, baseboards, and window and door frames. Spray at the rate of one square foot per second. To kill ticks and other labeled insects, apply directly to pests. Treat pets with an EPA registered [brand name] product in conjunction with this application.

For best results and to protect your pet against and to minimize reintroduction of adult fleas from outdoors, use EPA registered flea and tick products, ([such as] [brand name] flea or flea and tick collar, [brand name] pet powder [brand name] pet spray, [brand name] pet dip, [brand name] pet shampoo, flea shampoo or flea and tick pet shampoo), in conjunction with this application.

FOR USE IN CANNERIES: Thoroughly clean floors, conveying equipment and walls, paying special attention to cracks and crevices, prior to bringing in produce. Spray the entire space inside the cannery, including walls, ceiling and floors, paying special attention to forcing the spray into all cracks and crevices for the control of ants cockroaches, silverfish, crickets, spiders, fruit flies, moths, house flies, cheese mites and other listed insects.

FOR USE IN FOOD AND NONFOOD AREAS OF FOOD-PROCESSING PLANTS, Industrial installations, Bakeries, Rice and Wheat Mills, Restaurants, Tobacco Warehouses, Grain Elevators and Warehouses: Spray as directed to control listed flying and crawling insects. The use of this product in food-

processing or food-handling establishments should be confined to time periods when the plant is not in operation. Food should be covered or removed during treatment.

4/5

DRIED FRUIT PRODUCTS in storage or being processed: For control of, but not limited to: Almond Moths, Ants, Cereal Beetles, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Earwigs, Flies, Fruit Flies, Fungus Gnats, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles, Silverfish, Dried Fruit Beetles, Warehouse Beetles, Khapra Beetles and other listed insects. Do not apply directly to fruit. Apply as a fine mist or fog in the air above and around the trays, bins, and shelves at the rate of 1 to 10 fluid ounces per 1000 cubic feet of space.

TREATMENT OF STORED FOOD AREAS: Foods in multi-walled bags, cloth bags or other packaging stored in such areas as, but not limited to, listed areas: For control of accessible adult stages of, but not limited to: Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Dark Mealworms, Drugstore Beetles, Earwigs, Firebrats, Flies, Fruit Flies, Fungus Gnats, Grain Mites, Granary Weevils, Hide Beetles, Indian Meal Moths, Mediterranean Flour Moths, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles, Silverfish, Small Flying Moths, Spiders, Spider Beetles, Yellow Mealworms, Dried Fruit Beetles, Warehouse Beetles, Rusty Grain Beetles, Lesser Grain Beetles, Khapra Beetle and other listed pests. Infested stored products should be destroyed, fumigated or treated by other effective methods using an approved product intended for this purpose. As noted above, disperse first by directing spray into cracks and crevices, corners, pallets and around stacks or packaged goods hitting as many insects as possible and then apply as a space spray at a dosage of 1 to 1-1/2 fluid ounces per 1000 cubic feet of room space. Do not remain in treated area and ventilate before re-entry.

ON PEANUTS, NUTS, BEANS, SEEDS IN BULK, BAGS OR OTHER PACKAGING: For control of accessible stages of Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Dark Mealworms, Drugstore Beetles, Earwigs, Firebrats, Flies, Fruit Flies, Fungus Gnats, Grain Mites, Granary Weevils, Hide Beetles, Indian Meal Moths, Mediterranean Flour Moths, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles, Silverfish, Small Flying Moths, Spiders, Spider Beetles, Yellow Mealworms, Dried Fruit Beetles, Warehouse Beetles, Rusty Grain Beetles, Lesser Grain Beetles, Khapra Beetle and other listed insects, apply to the surface of stored or bagged products. Also thoroughly spray walls, floors and other surfaces of bins and storage and handling areas, being especially careful to treat cracks, crevices and similar hiding places. Then apply as a space spray by directing the spray toward the ceiling and upper corners of the area and behind any obstructions.

FOR USE IN FEDERALLY-INSPECTED MEAT AND POULTRY PLANTS: Apply with conventional mechanical or compressed air equipment at the rate of 1 to 1-1/2 fluid ounces per 1000 cubic feet of room space. Apply spray into all cracks and crevices in woodwork, walls, and floors, and underneath sinks, behind pipes and in all places that shelter insects. Contact as many insects as possible. Repeat as necessary. Before use, exposed food products and packaging materials must be covered or removed. After use, equipment and utensils must be washed with an effective cleaning compound and rinsed with potable water. Use as a space spray only when plant is not in operation.

FOR IN TRANSIT PROTECTION: To control accessible stages of listed insects, clean and spray truck beds, boxcars, ship holds and other listed transit equipment before loading. Thoroughly treat walls and floors and lightly spray each layer of stacked carton or containers. Apply at the rate of 1 gallon per 750 square feet.

FOR USE IN STORAGE FACILITIES AND EQUIPMENT:

SANITATION MEASURES: This product can be used as a surface spray in warehouse bins, trucks, cargo ships, mills, bin hoppers, elevators and conveying equipment as a clean-up prior to use. All grain-infested accumulations should be removed from bin hoppers in mills and elevators. All storage and conveying equipment should be thoroughly cleaned by sweeping out waste grain, cobwebs and other debris from the walls and rafters as well as on the floor and doorframes, paying special attention to material lodged in cracks and crevices. All debris should be removed and disposed of in outside trash.

FOR FARMS, particular attention should be given to cleaning up grain residues from wagons, harvesting equipment and feed troughs and around used feed and grain bags. Newly harvested grain should not be placed in the same bin with carry-over grain and all carry-over grain stocks that are not treated with grain protectant should be fumigated or treated with EPA approved products [such as] [brand name]. These cleaning operations should be done two or three weeks before harvest.

After above sanitation measures have been employed, spray as directed.

FOR USE IN ANIMAL QUARTERS, Barns, Dairies, Milking Parlors, Milking Rooms, Poultry Houses and other listed animal areas. For rapid control [kill] of Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Gnats and other listed flying insects, follow dilution chart and spray at the rate of 5 to 10 seconds per 1000 cubic feet of space, directing the spray toward the ceiling and upper corners. Do not direct spray towards animals or birds. For best results, close doors and windows before spraying and keep them closed for 10 to 15 minutes. Applicator should vacate the treated area and ventilate it prior to returning.

FOR USE ON LIVESTOCK:

To protect beef cattle, dairy cattle, horses and other livestock from attacks and to kill and repel Stable Flies, Horse Flies, Deer Flies, House Flies, Horn Flies, Lesser House Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, and Gnats, spray about 3 seconds on each side of animal, being careful to thoroughly spray back, withers and forelegs.

To protect from Face Flies, spray the face and hair or spray on a cloth and rub on the face, being certain to avoid contact with the eyes. Repeat treatment as necessary when flies are troublesome.

For effective control of biting and sucking lice on cattle, horses, sheep, goats, and hogs, spray to thoroughly wet the hair of the animal including the head and brush of the tail. Repeat treatment in 10 days to kill newly hatched lice.

To control poultry lice and exposed stages of Darkling Beetles, thoroughly spray roosts, walls, nests and cages, followed by spraying the birds with a fine mist.

For control of bedbugs and mites on poultry and in poultry houses, spray crevices of roost poles, cracks in walls and cracks in nests where bedbugs and mites hide, followed by spraying the birds with a fine mist.

To control sheep "tick" or ked, thoroughly wet all portions of the body by spraying directly into the wool. Treat at a rate to sufficiently wet the animal.

To kill fleas and ticks on livestock and to obtain protection against reinfestation, spray to sufficiently wet the animal.

5/4

FOR USE IN BARNs, DAIRIES, MILKING PARLORS, MILKING ROOMS, POULTRY HOUSES and other listed animal housing areas: Do not use in milk rooms where raw milk may be exposed.

FOR USE WITH HAND-HELD MECHANICAL FOGGERS to control Houseflies, Horn Flies, Stable Flies, Horse Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Gnats and other listed insects. Direct spray toward the ceiling and upper corners of the room until the area is filled with mist, about 2 fluid ounces per 1000 cubic feet of space. For best results, close doors and windows before spraying and keep them closed for 10 to 15 minutes after spraying. Applicator should vacate the treated area and ventilate it before reoccupying.

FOR USE THROUGH AUTOMATIC SPRAYING SYSTEMS

Automatic spraying systems must not be programmed to release pesticide where food or feed is directly exposed.

To Kill Houseflies, Horn Flies, Stable Flies, Horse Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Gnats and other listed insects, set nozzles to deliver 1 fluid ounce per minute. Position nozzles to cover a maximum of 2000 cubic feet of space per nozzle. Set timer to operate in accordance with equipment directions.

OUTDOOR USE: This spray can be used as is or diluted at the rate of 1 part concentrate to 2 parts water for a concentration of 0.05% Pyrethrins.

Caution should be taken to avoid spraying when wind conditions might create a mist that can blow back to applicator. Do not apply this product in or around bodies of water. Do not spray directly on plants. Spray surfaces and contact as many insects as possible.

FLIES, GNATS, MOSQUITOES, SMALL FLYING MOTHS, Midges, small flying insects and other flying insects: Hold container 18 – 24 inches from shrubs, bushes, grassy areas and other areas infested with these pests. Do not apply directly to plants. Spray with a slow, sweeping motion while moving away from the treated area. To reduce annoyance from these pests, treat open porches and patios in a similar manner. Kills and repels flies, mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, midges, gnats and, when used intermittently during outdoor activities, will keep surrounding areas free from these pests. Allow spray to dissipate before occupying the treated area.

ANTS (including Black Carpenter Ants) : Spray ant trails, anthills and other areas of entry and infestation. Spray on ants where possible.

SPIDERS: Spray directly on spiders and webs.

SOWBUGS, MILLIPEDES, BEETLES, EARWIGS, Pillbugs, Centipedes, Scorpions, Cockroaches, Crickets, Waterbugs, Palmetto Bugs: Thoroughly spray surfaces and into hiding places such as cracks, crevices, moist areas, openings around pipes, spigots, storage areas, sheds and other areas where these pests tend to congregate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

STORAGE: Store in a cool, dry place. Keep container closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse or equivalent and offer for recycling or reconditioning, or dispose of in a sanitary landfill or by other procedures approved by State and Local authorities.

Net Contents _____

Manufactured by:

Mc LAUGHLIN GORMLEY KING COMPANY
8810 Tenth Avenue North
Minneapolis, MN 55427

EPA Reg. No. 1021-1796

EPA Est. No. 1021-MN-2