



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 31, 2022

Tom Lennan
MGK
Aspen Lane North
Minneapolis, MN 55428

Subject: Label Amendment – Agency initiated action to clarify rates
Product Name: PYROCIDE® Concentrate 74406
EPA Registration Number: 1021-1800
Application Date: 01/18/2022
Decision Number: 581554

Dear Mr. Lennan:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Autumn Metzger at 202-566-2830, or Metzger.autumn@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Elizabeth Fertich". The signature is written in a cursive style with a large initial "E".

Elizabeth Fertich
Product Manager 4
Invertebrate-Vertebrate Branch 1
Registration Division
Office of Pesticide Programs

Enclosure

PYROCID[®] Concentrate 74406

[COMMERCIAL SECTION OF LABEL] [ABN: VAMPYRE[®] Misting Concentrate]

[] or () indicates alternate or optional wording
 [Graphic: MGK logo] [MGK[®] Insect Control Solutions]
 [Contains the botanical insecticide Pyrethrum [an extract of a Chrysanthemum flower]]
 [A micro-emulsion, water-based spray]
 [Kills and repels Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Mosquitoes, and Gnats]
 [Provides rapid kill of mosquitoes and other listed insects without objectionable residues]
 [For use with [Brand Name], [sold separately]]
 [Kills Black Widow Spiders and Black Carpenter Ants]
 [Quick killing Pyrethrum controls crawling and flying insects in the home, schools, hospitals and other listed areas]

[Controls Fleas and Ticks indoors and around pet areas]
 [Controls Deer Ticks and other Ixodid species that may carry and transmit Lyme disease]
 [For use in Animal Quarters, Dairies, Milk Rooms, Milking Parlors, Calving Areas, and Dairy and Hog Barns]
 [For use on Beef Cattle, Dairy Cattle, Horses, Hogs, Poultry, and other listed livestock]
 [Kills Mosquitoes, including the Culex species that may carry and transmit the West Nile virus]
 [Kills mosquitoes that may transmit Zika virus]
 [Kills Aedes mosquitoes]
 [Kills mosquitoes, including the Aedes mosquito that may carry and transmit Zika virus]

ACTIVE INGREDIENTS:

Pyrethrins	3.00%
* Piperonyl butoxide	30.00%
OTHER INGREDIENTS	67.00%
	100.00%

*(butylcarbityl) (6-propylpiperonyl) ether and related compounds

PYROCID[®], MGK[®], VAMPYRE[®] - registered trademarks of McLaughlin Gormley King Company

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	


Manufactured by

 7325 Aspen Lane North
 Minneapolis MN 55428

EPA Reg. No. 1021-1800

EPA Est. No. 1021-MN-2

Net Contents: 1 Gallon



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt,
- Long pants,
- Chemical-resistant gloves made of barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or Viton, and
- Shoes and socks.

In addition to the above PPE, applicators using a high-pressure hand wand in an enclosed area must wear at least a half-face NIOSH-approved respirator with:

- Any R or P filter with a NIOSH approval number prefix TC-84A.

In addition to the above PPE, applicators using handheld foggers in an enclosed area must wear a half-face, full-face, or hood-style NIOSH-approved respirator with:

- A NIOSH-approved particulate filter with any R or P filter with NIOSH approval number prefix TC-84A
- A NIOSH-approved powered air purifying respirator with an HE filter with NIOSH approval number prefix TC-21C.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through runoff. This product has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use this product in or on electrical equipment because of shock hazard.

[Pyrocid® Concentrate 74406] contains the quick killing botanical insecticide, Pyrethrum [an extract of Chrysanthemum flowers], which provides rapid effective control of insects upon direct contact with an added benefit of flushing [repelling] these insects to aid in a more complete control [kill]. This highly effective water-based formula can be used on flying and crawling insects indoors and outdoors.

[Pyrocid Concentrate 74406] can be used to control fleas and ticks hidden in carpets, rugs, upholstery, draperies, floor cracks and cabinets. Also controls Deer Ticks and other Ixodid species that may carry and transmit Lyme disease.

[Pyrocid Concentrate 74406] can be used in the home, schools, hospitals, restaurants and other listed areas to control [kill], on contact, listed crawling pests and is a fast, effective product for flying insect control. Pyrethrum has quick killing ability with no persistent residue.

[Pyrocid Concentrate 74406] can be used to reduce annoyance of mosquitoes, including the Culex species that may carry and transmit the West Nile virus, outdoors around homes, in recreational areas, and other listed areas. Ideal for use before outdoor activities.

[Pyrocid Concentrate 74406] can be used in food and non-food areas of food processing plants, canneries, industrial installations, bakeries, rice and wheat mills, tobacco warehouses, grain elevators, warehouses, storage sites and other listed areas to control [kill], on contact, listed crawling pests and is a fast, effective product for flying insect control.

[Pyrocid Concentrate 74406] provides rapid kill of insects without objectionable residues.

To protect your pet against and to minimize reintroduction of adult fleas from outdoors, use EPA registered flea and tick products ([such as] flea or flea and tick collar, pet powder, pet spray, pet dip, pet shampoo, flea shampoo or flea and tick pet shampoo), in conjunction with this application and prior to re-entry.

Remove pets and birds and cover fish aquariums before spraying.

FOR THE CONTROL OF INSECTS including: Almond Moths, Angoumois Grain Moths, Ants, Bed Bugs, Black Carpenter Ants, Black Flies, Black Widow Spiders, Blood Sucking Lice, Boxelder Bugs, Brown Dog Ticks, Cadelles, Carpet Beetles, Cat Fleas, Cellar Spiders, Cereal Beetles, Cheese

Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Clothes Moths, Clover Mites, Cockroaches, Confused Flour Beetles, Crickets, Dark Mealworms, Darkling Beetles, Deer Flies, Deer Ticks, Dermestid Beetles, Dog Fleas, Driedfruit Beetles, Drugstore Beetles, Earwigs, Face Flies, Fannia Flies, Firebrats, Flat Grain Beetles, Fleas, Flies, Fruit Flies, Fungus Gnats, Gnats, Grain Mites, Granary Weevils, Hide Beetles, Horn Flies, Hornets, Horse Flies, House Flies, Indian Meal Moths, Khapra Beetles, Lesser Grain Beetles, Lesser Grain Borers, Lesser House Flies, Lice, Mediterranean Flour Moths, Merchant Grain Beetles, Midges, Miller Moths, Millipedes, Mosquitoes – including the Culex species that may carry and transmit the West Nile virus, Palmetto Bugs, Pillbugs, Poultry Lice, Red Flour Beetles, Rice Flour Beetles, Rice Weevils, Rusty Grain Beetles, Sawtoothed Grain Beetles, Sheep Ticks, Silverfish, Skipper Flies, Small Flying Moths, Sowbugs, Spider Beetles, Spiders, Stable Flies, Ticks that may carry and transmit Lyme disease, Tobacco Moths, Trogoderma Beetles, Warehouse Beetles, Wasps, Waterbugs, Yellow Mealworms and Yellowjackets.

INDOOR AREAS for use in: Animal Areas, Animal Control Centers, Animal Quarters, Animal Shelters, Apartments, Atriums, Attics, Auditoriums, Automobiles, Bakeries, Bars, Basements, Bathrooms, Bedding, Bedrooms, Beef Barns, Beverage Plants, Boats, Bookcases, Bottling Plants, Breweries, Brewery Warehouses, Buildings Housing Animals, Buildings Housing Poultry, Buses, Cabinets, Cabins, Cafeterias, Calving Barns, Camp Sites, Campers, Canneries, Carpets, Cat Sleeping Quarters, Cattle Barns, Chests, Churches, Circus Tents, City Missions, Closed Porches, Closets, Clothes Storage, Club Houses, Commercial Buildings, Condominiums, Crawl Spaces, Credenzas, Crematoriums, Dairies, Dairy and Livestock Barns, Day Care Centers, Decks, Delicatessens, Dens, Department Stores, Diners, Dog Houses, Dog Sleeping Quarters, Dormitories, Draperies, Dressers, Dried Fruits in Storage, Drive-Ins, Drugstores, Dumpsters, Dwellings, Egg Processing Plants, Fabric Storage Areas, Factories, False Ceilings, Farm Structures, Floors, Food Processing Plants, Food Storage Areas, Freight Containers, Fruit Packing Sheds, Funeral Parlors, Furniture, Garages, Garbage Bins, Garbage Cans, Garbage Compactors, Gas Stations, Gazebos, Goat Holding Areas, Grain Elevators, Grain Handling Equipment, Grain Harvesting Equipment, Grain Mills, Grain Storage, Granaries, Grocery Stores, Hog Barns, Homes, Horse Stables and Barns, Horse Trailers, Hospices, Hospitals, Hotels, Household Contents, Houses, Indoor Eating Buildings, Industrial Installations, Industrial Plants, Institution Dining Areas, Institutions, Jails, Kennels, Kitchens, Laboratories, Lanais, Laundromats, Lavatories, Libraries, Liquor Stores, Livestock Barns, Livestock Hauling Equipment, Livestock Holding Areas, Living Rooms, Loafing Sheds, Locker Rooms, Marinas, Mattresses, Mausoleums, Meat Packing Plants, Mess Halls, Milk Houses, Milk Rooms, Milk Sheds, Milking Rooms, Mini Storage, Mobile Canteens, Mobile Homes, Morgues, Mortuaries, Motels, Movie Theaters, Museums, Nursing Homes, Office Buildings, Other Public Buildings, Outdoor Eating Buildings, Pantries, Parking Garages, Patios, Pet Bedding, Pet Carriers, Pet Grooming Parlors, Pet Sleeping Areas, Pet Stores, Planes, Play Houses, Play Rooms, Pool Halls, Porches, Poultry Houses, Poultry Plants, Pounds, Prisons, Public Storage, Quonset Huts, Rabbit Processing Plants, Railroad Cars, Recreational Vehicles, Research Animal Quarters, Restaurants, Retail Stores, Rice Mills, Rooms, Rugs, Schools, Seed Storage, Sheep Holding Areas, Ship's Holds, Ships, Smoke Shops, Solariums, Sport Stadiums, Stable Bins, Stables, Storage Areas, Stored Food (bags), Stored Products, Stores, Sun Porches, Sun Rooms, Swine Houses, Taverns, Tents, Textile Mills, Theaters, Tobacco Factories, Tobacco Warehouses, Tool Sheds, Trailers, Trains, Transportation Vehicles, Tree Houses, Tree Nuts Storage Areas, Truck Trailers, Trucks, Tunnels, Utilities, Utility Rooms, Vending Machines, Verandas, Veterinary Clinics, Voids in Attics, Warehouses, Washrooms, Wheat Mills, Wholesale Stores, Wine Cellars, Wineries and Zoos.

OUTDOOR AREAS for use in and around: Alleys, Amphitheaters, Amusement Parks, Ant Hills, Back Yards, Balconies, Baseball Fields, Beaches, Bridges, Campgrounds, Carwashes, Circuses, Construction Sites, Corrals, Courtyards, Curbs, Drive-In Restaurants, Drive-In Theaters, Dumpsters, Fair Grounds, Feedlots, Front Yards, Game Fields, Garbage Dumps, Gazebos, Grandstands, Junkyards, Kennels, Levees, Manholes, Manure Piles, Marinas, Outdoor Eating Establishments, Parking Lots, Parks, Patios, Playgrounds, Ranges, Recreational Areas, Rest Areas, Roadsides, Rooftops, Sanitary Sewers, Sidewalks, Side Yards, Stadiums, Tennis Courts, Truck Stops, Underpasses, Utility Rooms Outdoors, Weigh Stations, Wild Animal Parks, Wood Piles and Zoos.

FOR USE ON: Beefalo, Birds, Buffalo, Calves, Cattle- Beef, Cattle- Dairy, Chickens, Donkeys, Ducks, Emus, Exotics such as European Red Deer, Geese, Horses, Lambs, Llamas, Mules, Ostrich, Oxen, Pheasants, Pigs, Pullets, Quail, Ratites, Roosters, Sheep, Swine and Turkeys.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product
in a manner inconsistent with its labeling.

READ ALL DIRECTIONS COMPLETELY BEFORE USE. For best results, follow directions for specific use areas.

RESTRICTIONS:

- **ENTRY RESTRICTIONS:**
 - For surface or direct spray use: Except when applying to livestock, do not enter or allow others to enter treated area until sprays have dried.
 - For Space Spray Use: Do not enter or allow others to enter until vapors, mists, and aerosols have dispersed, and the treated area has been thoroughly ventilated.
- **APPLICATION RESTRICTIONS:**
 - Except when applying to livestock, do not apply this product in a way that will contact workers or other persons, either directly or through drift.
 - Except when applying to livestock, only protected handlers may be in the area during application.
 - Do not make applications during rain.
 - Do not remain in treated area. Exit area immediately and remain outside the treated area until aerosols, vapors and/or mists have dispersed.
 - Do not wet articles to point of runoff or drip.
 - Do not use treated articles until spray has dried.
 - Remove or cover exposed food and drinking water before application.
 - Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces, or wash them before use.
 - Do not water the treated area to the point of runoff.
 - When used in dairy barns or facilities: Close milk bulk lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
 - Do not make space spray applications when facility is in operation. Prior to space spray applications, cover or remove food. Prior to space spray applications, cover food processing surfaces or clean after treatment and before use.
 - Do not apply directly into sewers or drains, or to any area like a gutter where drainage to storm sewers, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
 - Except when applying to sewers or drains as instructed on this label, do not apply directly to sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
 - Except for use with outdoor fogging devices, all outdoor applications must be limited to spot or crack and crevice treatments only, except for the following permitted uses:
 - Treatment to soil or vegetation around structures;

- Applications to lawns, turf, and other vegetation;
 - Applications to building foundations up to a maximum height of 3 feet.
- Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack and crevice applications, only.
- Do not use in aircraft cabins.

GENERAL: This concentrated product must be diluted with water. When making dilutions, add concentrate to the tank and add water. Agitate well while adding water and apply as directed:

0.012% Pyrethrins—add 0.5 fl. oz. of concentrate/gallon of water (3.9 ml/L)
 0.020% Pyrethrins—add 1.0 fl. oz. of concentrate/gallon of water (7.8 ml/L) -
 0.04% Pyrethrins – add 1.6 fl. oz. of concentrate / gallon of water (13 ml/L)
 0.05% Pyrethrins – add 2 fl. oz. concentrate / gallon of water (15.86 ml/L)
 0.10% Pyrethrins – add 4.3 fl. oz. concentrate / gallon of water (34.10 ml/L)
 0.30% Pyrethrins – add 13 fl. oz. concentrate / gallon of water (103.09 ml/L)
 0.40% Pyrethrins—add 17 fl. oz. Concentrate / gallon of water (218.75 ml/L)
 0.50% Pyrethrins – add 22 fl. oz. concentrate / gallon of water (174.46 ml/L)
 0.065 Pyrethrins—add 28 fl. z of concentrate/ gallon of water (128.75 ml/L)_

Confine use of this product in food processing establishments to time periods when the plant is not in operation. Remove or cover food during treatment. Cover all food processing surfaces during treatment or thoroughly clean before use following treatment.

For maximum effectiveness, following the dilution chart and using a combination of spot, surface treatment and fogging are recommended.

SPACE SPRAYING IN FOOD HANDLING ESTABLISHMENTS AND COMMERCIAL ESTABLISHMENTS: Use with conventional mechanical or compressed air equipment to kill exposed accessible stages of: Flies (including Fruit Flies), Mosquitoes including the Culex species that may carry and transmit the West Nile virus, Small Flying Moths, Fleas, Gnats, Wasps, Hornets, Clover Mites, Cheese Mites, Cheese Skippers, Boxelder Bugs, Earwigs, Ants, and the following common Stored Product Pests: Granary Weevils, Rice Weevils, Confused Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths, Tobacco Moths, Yellow Mealworms, Dark Mealworms, Grain Mites, Cadelles, Red Flour Beetles, and other listed insects. Close room and shut off air conditioning or ventilating equipment. Apply at the rate of 1 to 1.6 fl. oz. of concentrate per 1,000 cu. ft. (1.04 ml/m³ to 1.67 ml/m³) of space and direct spray toward upper corners of room and ceiling, filling the room with mist. Leave room closed for ½ hour and ventilate thoroughly before re-entry. Do not exceed the maximum application rate of 0.0033 lb pyrethrins per 1,000 cu. ft. or 1.6 fl. oz of concentrate per 1,000 cu. ft. of space.

SURFACE SPRAYING: To kill Cockroaches, Black Carpenter Ants, Silverfish, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles and other listed insects, use a sprayer adjusted to a coarse wet spray. Mix well and apply as a coarse wet spray, using approximately 1 gallon of spray mixture per 1,000 sq. ft. of surface (40.75 ml/m²). To provide maximum penetration and impingement on the insects, direct the spray first into all cracks and crevices, behind all equipment, cupboards, obstructions or dark harborage areas, on walls and floors, underneath sinks, behind pipes, beneath stoves and refrigerators, around garbage cans and cabinets, along baseboards, door sills, window sills, door and window frames and floors, around and on drains, pipes, and plumbing, behind bookcases, storage and other utility installation areas, and in all places that harbor insects. Contact as many insects as possible. Do not allow spray to contact stored food. If surface application only is to be used, spray floor and other surfaces. To kill Ants, treat doors and around window frames and other areas of entry. Spray on Ants whenever possible. Apply at the rate of 28 fl. oz. of concentrate/gallon of water per 1,000 sq. ft. Do not exceed the maximum application rate of 0.056 lbs. pyrethrins per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1,000 sq. ft.

CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply liberally to empty chests, closets, bureaus, and other storage areas, directing the spray into cracks, joints, and crevices. Apply at the rate of 28 fl. oz. of concentrate/gallon of water per 1,000 sq. ft. Do not exceed maximum application rate of 0.056 lbs. pyrethrins per 1,000 sq. ft. or 28 fl. oz. concentrate per 1,000 sq. ft.

CARPET BEETLES: Directly spray as many insects as possible. Spray all surfaces of upholstered furniture. Where possible, thoroughly spray both top and bottom of rugs and carpets. Also treat shelves, closets, floors, walls, windowsills, baseboards, areas under furniture, and wherever these insects are seen or suspected. Apply at a rate of 28 fl. oz. of concentrate/gallon of water per 1,000 sq.ft. Do not exceed the maximum application rate of 0.056 lbs. pyrethrins per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1000, sq. ft

BED BUGS: Spray mattresses lightly, paying special attention to tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, floors, walls, furniture, and other infested areas. Apply at a rate of 28 fl. oz. of concentrate/gallon of water per 1,000 sq.ft. Do not exceed the maximum application rate of 0.056 lbs. pyrethrins per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1,000 sq. ft

FOR USE IN LISTED PET AREAS: For best results remove, destroy, or wash pet's old bedding. To reduce infestation, thoroughly spray pet [dog] [cat] [puppy] [kitten] [animal] bedding, sleeping quarters, kennel, around baseboards, window and door frames, into wall cracks, on local areas of floors, and other animal resting places until the surface is slightly moist. Treat these areas at the rate of two seconds per linear foot at the distance of 12 to 18 inches (30.5 to 45.7 cm). Also, spray the entire inside surface of doghouses. Put fresh bedding in pet's quarters after spray has dried. Allow treated areas to dry thoroughly before allowing pets to re-enter. Apply at a rate of 28 fl. oz. of concentrate/gallon of water per 1,000 sq.ft.

Do not exceed the maximum application rate of 0.056 lbs. Pyrethrins/ 1,000 sq. ft. or 0.56 lbs. PBO per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1,000 sq. ft.

FLEA AND TICK INFESTATION ON RUGS AND CARPETS: Before use, vacuum all infested areas thoroughly and dispose of vacuum bag in outside trash. Spray rugs and carpets where fleas tend to hide and usually where animals frequent. Spray thoroughly into hiding places such as cracks and crevices, in walls, on floors, and along baseboards, window and door frames. To kill ticks and other labeled insects, apply directly to bugs. Apply at a rate of 28 fl. oz. of concentrate/gallon of water per 1,000 sq.ft.

Do not apply to pets. Treat pets with an EPA registered [brand name] product in conjunction with this application.

For best results, to protect your pet against and to minimize reintroduction of adult fleas from outdoors, use EPA registered flea and tick products, ([such as] [brand name] flea or flea and tick collar, [brand name] pet powder, [brand name] pet spray, [brand name] pet dip, [brand name] pet shampoo, flea shampoo or flea and tick pet shampoo), in conjunction with this application.

FOR KENNELS AND OTHER COMMERCIAL OUTDOOR ANIMAL AREAS: To control listed pests. Apply at a rate of 28 fl. oz. of concentrate/gallon of water per 1,000 sq.ft. Do not exceed the maximum application rate of 0.056 lbs. Pyrethrins/ 1,000 sq fr. or 0.56 lbs. PBO per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1, 000 sq. ft.

Apply liberally to surfaces, [paying] [with] special attention to cracks and crevices and other insect hiding places. Allow treated areas to dry thoroughly before allowing pets [animals] to re-enter.

FOR USE IN CANNERIES: Before bringing in produce, thoroughly clean floors, conveying equipment, and walls, paying special attention to cracks and crevices. Apply at a rate of 28 fl. oz. of concentrate / gallon of water per 1,000 sq. ft. and spray the entire area inside the cannery, including walls, ceilings, and floors, paying special attention to forcing the spray into all cracks and crevices for control of ants, cockroaches, silverfish, crickets, spiders, fruit flies, moths, house flies, cheese mites, and other listed insects. Do not exceed maximum application rate of 28 fl. oz. of concentrate per 1,000 sq. ft.

FOR USE IN FOOD AND NON-FOOD AREAS OF FOOD PROCESSING PLANTS, Industrial installations, Bakeries, Rice and Wheat Mills, Restaurants, Tobacco Warehouses, Grain Elevators and Warehouses: Spray as directed to control listed flying and crawling insects. Confine the use of this product in food processing or food handling establishments to time periods when the plant is not in operation. Cover or remove food during treatment. When applying as a space spray, apply at the rate of 1 to 1.6 fl. oz. of concentrate per 1,000 cu. ft. (1.04 ml/m³ to 1.67 ml/m³), not to exceed 0.0033 lb. pyrethrins or 1.6 fl. oz of concentrate per 1,000 cu ft. of space. . When applying as a surface spray apply at the rate of 28 fl. oz. of concentrate/gallon of water per 1,000 sq. fr. do not to exceed the maximum application rate of 28 fl. oz. of concentrate per 1,000 sq. ft.

DRIED FRUIT PRODUCTS in storage or processing: For control of [[listed insects]: Almond Moths, Ants, Cereal Beetles, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Earwigs, Flies, Fruit Flies, Fungus Gnats, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Sawtoothed Grain Beetles, Silverfish, Driedfruit Beetles, Warehouse Beetles, Khapra Beetles, and other listed insects]. Remove and destroy infested products. Do not apply directly to fruit. Apply as a fine mist or fog in the air above and around trays, bins, and shelves at the rate of 0.5 fl. oz. of concentrate per 1,000 cu. ft. of space (0.522 ml/m³). Repeat application at this dosage as required. Leave the room closed for one hour. Do not remain in treated areas and ventilate the area before re-entry. Sweep up and destroy fallen insects. Do not exceed the maximum application rate of 0.001 lbs. pyrethrins per 1,000 sq. ft. or 0.5 fl. oz. of concentrate per 1000, cu. ft of space.

TREATMENT OF STORED FOOD AREAS: Foods in multi-walled bags, cloth bags or other packaging stored in such areas as listed areas [for the control of listed pests]. [For the control of accessible adult stages of: Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Dark Mealworms, Drugstore Beetles, Earwigs, Firebrats, Flies, Fruit Flies, Fungus Gnats, Grain Mites, Granary Weevils, Hide Beetles, Indian Meal Moths, Mediterranean Flour Moths, Mosquitoes including the Culex species that may carry and transmit the West Nile virus, Red Flour Beetles, Rice Weevils, Sawtoothed Grain Beetles, Silverfish, Small Flying Moths, Spiders, Spider Beetles, Yellow Mealworms, Driedfruit Beetles, Warehouse Beetles, Rusty Grain Beetles, Lesser Grain Beetles, Khapra Beetles, and other listed pests.] Destroy, fumigate, or treat infested stored products using effective methods or an approved product intended for this purpose. As noted above, disperse first by directing spray into cracks and crevices, and around corners, pallets, and stacks of packaged goods, hitting as many insects as possible. Then apply as a space spray at a dosage of 1 to 1.6 fl. oz. of concentrate per 1,000 cu. ft. (1.04 ml/m³ to 1.67 ml/m³) of room space. Repeat application as necessary for insect control. Do not remain in treated area and ventilate before re-entry. Sweep up and destroy fallen insects.

ON PEANUTS, NUTS, BEANS, SEEDS IN BULK, BAGS OR OTHER PACKAGING: For the control of [listed insects] [accessible stages of Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Dark Mealworms, Drugstore Beetles, Earwigs, Firebrats, Flies, Fruit Flies, Fungus Gnats, Grain Mites, Granary Weevils, Hide Beetles, Indian Meal Moths, Mediterranean Flour Moths, Mosquitoes including the Culex species that may carry and transmit the West Nile virus, Red Flour Beetles, Rice Weevils, Sawtoothed Grain Beetles, Silverfish, Small Flying Moths, Spiders, Spider Beetles, Yellow Mealworms, Driedfruit Beetles, Warehouse Beetles, Rusty Grain Beetles, Lesser Grain Beetles, Khapra Beetles, and other listed insects]. Apply as a surface spray to the surface of stored or bagged products at a rate of 28 fl. oz. of cocentrate / gallon of water per 1,000 sq ft. Do not exceed the maximum application rate of 056 lb. pyrethrins or 28 fl. oz. of cocentrate per 1,000 sq ft. . Also spray walls, floors, and surfaces of bins and storage and handling areas thoroughly, being especially careful to treat cracks, crevices, and similar hiding places. Then apply as a space spray at the rate of 1 to 1.6 fl. oz. of concentrate per 1,000 cu. ft. (1.04 ml/m³ to 1.67 ml/m³) by directing the spray toward the ceiling and upper corners of the area and behind any obstructions. For the space spray, do not exceed the maximum application rate of 0.0033 pyrethrins or 1.6 fl. oz of concentrate per 1,000 cu. ft. of space.

FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS: Apply with conventional mechanical or compressed air equipment at the rate of 1 to 1.6 fl. oz. of concentrate per 1,000 cu. ft. (1.04 ml/m³ to 1.67 ml/m³) of room space. Apply spray into all cracks and crevices in woodwork, walls, and floors, underneath sinks, behind pipes, and in all places that shelter insects. Contact as many insects as possible. Repeat as necessary. Before use, cover or remove exposed food products and packaging materials. After use wash equipment and utensils with an effective cleaning compound and rinse with potable water. Use as a space spray only when plant is not in operation. Do not exceed the maximum application rate of 0.0033 lb pyrethrins or 1.6 fl. oz of concentrate per 1,000 cu ft. of space.

FOR IN TRANSIT PROTECTION: To control accessible stages of listed insects, clean and spray truck beds, boxcars, ship's holds and other listed transit equipment before loading. Thoroughly treat walls and floors, and lightly spray each layer of [stacked] cartons or containers. Apply at a rate of 28 fl. oz. of concentrate/ gallon of water per 1, 000 sq. ft. . Do not exceed maximum application rate of 0.056 lb pyrethrins or 28 fl. oz. of concentrate per 1,000 sq. ft.

FOR USE IN STORAGE FACILITIES AND EQUIPMENT:

SANITATION MEASURES: This product can be used as a surface spray in warehouse bins, trucks, cargo ships, mills, bin hoppers, elevators, and conveying equipment as a cleanup prior to use. In mills and elevators, remove all infested accumulations of grain from the bin hoppers. All storage and conveying equipment should be thoroughly cleaned by sweeping out waste grain, cobwebs, and other debris from the walls, rafters, floors, and door frames, paying special attention to material lodged in cracks and crevices. Remove all debris and dispose of in outside trash. Apply at the rate of 28 fl. oz. of concentrate per 1,000 sq ft. Do not exceed maximum application rate of 0.056 lb. of pyrethrin or 28 fl. oz. of concentrate per 1,000 sq. ft.

FOR FARMS, pay particular attention to cleaning up around used feed and grain bags and cleaning grain residues from wagons, harvesting equipment, and feed troughs. Do not place newly harvested grain in the same bin with carry-over grain. Fumigate or treat all carry-over grain stocks not previously treated with grain protectant with EPA approved products [such as] [brand name]. Perform these cleaning operations two or three weeks before harvest.

After above sanitation measures are employed, spray as directed.

FOR USE IN ANIMAL QUARTERS, Barns, Dairies, Milking Parlors, Milking Rooms, Poultry Houses, and other listed animal areas. For rapid control [kill] of Flies, Mosquitoes including the Culex species that may carry and transmit the West Nile virus, Gnats, and other listed flying insects: Dilute with water to give a 0.10% or 0.50% concentration and apply as a fine mist or spray. Direct spray towards ceiling and upper corners of area until filled with mist, using 1.6 fl. oz. of concentrate per 1,000 cu. ft. (1.67 ml/m³) of space. Do not direct spray towards animals or birds. For best results, close doors and windows before spraying and keep them closed for 10 to 15 minutes following application. Applicator should vacate the treated area and ventilate it before reoccupying. Do not use in milk rooms where raw milk may be exposed.

FOR USE ON LIVESTOCK: Create a solution containing 0.40% pyrethrins by adding 17 fl. oz of concentrate / gallon of water. Do not apply to livestock animals more than once per day.

To protect beef cattle, dairy cattle, horses, and other livestock from attacks and to kill and repel Stable Flies, Horse Flies, Deer Flies, House Flies, Horn Flies, Lesser House Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile virus, and Gnats, spray 3 seconds on each side of animal, being certain to spray the back, withers and forelegs thoroughly.

To protect from Face Flies, spray the face and hair or spray on a cloth and rub on the face, being certain to avoid contact with eyes. Repeat treatment as necessary when flies are troublesome.

For effective control of Biting and Sucking Lice on cattle, horses, sheep, goats and hogs, spray to thoroughly wet the hair of the animal, including the head and brush of the tail. Repeat treatment in 10 days to kill newly hatched lice.

To control Poultry Lice and exposed stages of Darkling Beetles, thoroughly spray roosts, walls, nests, and cages. Apply at a rate of 28 fl oz of concentrate / Gallon of water per 1,000 sq ft. of area Do not exceed the maximum application rate of 0.056 lb. pyrethrins or 28 fl. oz of concentrate per 1,000 sq. ft. Follow by spraying over the birds with a fine mist apply at a rate of 0.4 fl oz of concentrate / gallon of water per, 1,000 cu. ft. Do not exceed 0.0008 lb. pyrethrins or 0.4 fl oz of concentrate per 1,000 cu ft. of space.

For control of Bed bugs and Mites on poultry and in poultry houses, spray crevices of roost poles and cracks in walls and nests where Bed bugs and Mites hide. Apply at a rate of 28 fl. oz. of concentrate/gallon per 1000 sq. ft. of area. Do not exceed the maximum application rate of 0.056 lb. pyrethrins or 28 oz of concentrate per 1000 sq. ft. Follow by spraying over the birds with a fine mist applied at a rate of 0.4 fl.oz of concentrate/gallon water per 1000 cu. ft Do not exceed 0.0008 lb. pyrethrins or 0.4 fl. oz. of concentrate per 1000 cu. ft. of space.

To control Sheep "Tick" or Ked, create a solution containing 0.40%pyrethrins by adding 17 fl. oz of concentrate/gallon water and spray with enough solution to sufficiently wet the animal.

To kill Fleas and Ticks on livestock and to obtain protection against reinfestation, spray to sufficiently wet animal.

FOR USE IN BARNs, DAIRIES, MILKING PARLORS, MILKING ROOMS, POULTRY HOUSES, and other listed animal housing areas, follow dilution chart and apply as directed. Do not use in milk rooms where raw milk may be exposed.

0.10% Pyrethrins – Add 4.3 fl. oz. per gallon of water

0.50% Pyrethrins – Add 22 fl. oz. per gallon of water

DIRECTIONS FOR USE WITH HANDHELD MECHANICAL FOGGERS to control House Flies, Horn Flies, Stable Flies, Horse Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile virus, Gnats, and other listed insects. Dilute concentrate and apply as a fine mist or fog. Direct spray toward the ceiling and upper corners of the room until the area is filled with mist, using 1.6 fl. oz. per 1,000 cu. ft. (1.67 ml/m³) of space. For best results, close doors and windows before spraying and keep them closed for 10 to 15 minutes after spraying. Applicator should vacate the treated area and ventilate it before reoccupying.

DIRECTIONS FOR USE IN AUTOMATIC ULV SPRAYING SYSTEMS, IN AREAS SUCH AS ANIMAL QUARTERS, DAIRY AND BEEF BARNs, POULTRY HOUSES, SWINE HOUSING, WAREHOUSES, ZOOS AND OTHER LISTED LIVESTOCK OR ANIMAL HOUSING AREAS.

USE RESTRICTIONS FOR AUTOMATIC ULV SPRAYING SYSTEMS AND RESIDENTIAL MISTING SYSTEMS:

Do not apply this pesticide when people, pets and food are present.

- Do not use in an evaporative cooling system.
- Do not set nozzles to direct mist near outside air conditioner systems or other home air intakes.
- Direct nozzles to spray towards the target area and away from swimming pools, water bodies, or eating and cooking areas.
- When using this product, installers and service technicians must comply with the license, certification, or registration requirements of the state(s), tribe(s), or local authority(ies) where they are installed.
- If the system works when a person operates a remote activation device, then application of this pesticide when people, pets and food are present is prohibited.
- If system works on an automatic timer, set the timing for application when people, pets and food are unlikely to be present.
- Do not use in misters located within 3 feet of air vents, air conditioner units, or windows.
- If used in a system with a reservoir tank for the end-use dilution, the system reservoir tank must be locked. Securely attach the end-use pesticide label and a dilution statement to the system reservoir tank in a weather protected area or plastic sleeve. The dilution statement must be phrased as follows: this container holds ___ parts [product name] to ___ parts water.
- If used in a direct injection system, the pesticide container must be locked. Securely attach the end-use label to the pesticide container in a weather protected area or plastic sleeve.
- Automatic spraying systems must not be programmed to release pesticides where food or feed is directly exposed.

To kill Flies, House Flies, Horn Flies, Stable Flies, Horse Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile virus, Gnats and other listed insects: Add concentrate to tank and add water to obtain desired dilution. Agitate well while adding water. Dispense diluted concentrate through automatic spraying system. Set nozzles to deliver 1 fl. oz. per minute. Position nozzles to cover a maximum of 2,000 cu. ft. (56.63 m³) of space per nozzle. Set timer to operate in accordance with equipment directions. Equipment must be calibrated to deliver no more than the maximum application rate of 0.0033 lbs. Pyrethrins per 1,000 cu. ft. and 0.033 lbs. of PBO per 1,000 cu. ft., or no more than 1.6 fl. oz. of concentrate per 1,000 cu. ft. of space per day.

DIRECTIONS FOR USE IN OUTDOOR RESIDENTIAL MISTING SYSTEMS [Unit / Misters]:

NOTE: NOT REGISTERED for use in Automatic ULV Spraying Systems in the State of New York.

Use to control or temporarily reduce annoyance from accessible stages of: Mosquitoes including the Culex species that may carry and transmit the West Nile virus, Flies, Gnats, Face Flies, Fannia Flies, Deer Flies, Horse Flies, Stable Flies, Horn Flies, Hornets, Wasps, Yellowjackets, Firebrats, Small Flying Moths, Earwigs, Fleas, Ticks, Clover Mites, Brown Dog Ticks, Crickets, Cockroaches, Silverfish, Spiders, Darkling Beetles, and other nuisance insects.

Automatic Spraying System Dilution Table

A. System Tank size (in gallons)	B. Conversion Factor (fl. oz. of concentrate per gallon of water)	A. x B. = fl. oz. of concentrate for that tank size, to be mixed with water to achieve 0.044% pyrethrin solution.
30 (113.7 L)	1.88 (55.59 ml)	56.40 (1,668 ml)
55 (208.4 L)	1.88 (55.59 ml)	103.40 (3,058 ml)
125	1.88 (55.59 ml)	235 (6,949 mls)
250	1.88 (55.59 ml)	470 (13,898 mls)

When filling tank, partially fill the system [unit / mister] reservoir tank with water and add concentrate. Then fill the rest of the tank with water and agitate well.

Contact the system manufacturer for guidance or assistance to ensure compliance with the following requirement:

This product must only be used in systems that have been calibrated to apply no more than the maximum application rate of .00011 lbs. pyrethrins per 1,000 cu. ft. per day or 0.00058 lbs. piperonyl butoxide per 1,000 cu. ft. per day, whichever is lower (do not exceed 2 fl. oz. of diluted product per 1,000 cu. ft. per day).

For best performance, the spray solution should be in the pH range of 5.5 to 7.0.

This product will control insects directly contacted by its spray solution. Since many of the targeted insects can migrate in and out of treatment areas, it is recommended that this concentrate be used as part of an Integrated Pest Management (IPM) program. Control practices including eliminating breeding and harborage sites combined with appropriate use of contact insecticides, Insect Growth Regulators (IGRs) and / or residual insecticides will provide the most effective control of target insects.

OUTDOOR USE DILUTION TABLE:

Solution percentage	Ratio of concentrate to water	Fl. oz. of concentrate per gallon of water
0.02% Pyrethrins	1 part concentrate to 128 parts water	1 fl. oz of concentrate / gallon of water (7.8 ml/L) Apply at the rate of 1 gallon / 1,000 sq. ft.

Do not spray when wind conditions might create a mist that can blow back to applicator. Do not apply this product in or around bodies of water. Do not spray directly on plants. Spray surfaces and contact as many insects as possible.

SMALL FLYING INSECTS AND OTHER FLYING INSECTS including FLIES, GNATS, MOSQUITOES, SMALL FLYING MOTHS, and MIDGES: Apply at a rate of 1.0 fl. oz. of concentrate/gallon of water per 1,000 sq. ft. of area. Hold sprayer nozzle 18 to 24 inches (45.72 to 60.96 cm) from shrubs, bushes, grassy areas, and other areas infested with these pests. Do not apply directly to plants. Spray with a slow sweeping motion while moving away from the treated area. To reduce annoyance from these pests, treat open porches, patios and under decks in a similar manner. Kills and repels Flies, Mosquitoes including the Culex species that may carry and transmit the West Nile virus, Midges, and Gnats when used intermittently during outdoor activities and will keep surrounding areas free from these pests. Allow spray to dissipate before occupying the treated area. Do not exceed 0.002 lb. pyrethrins or 1 fl. oz. of concentrate per 1000 sq. ft. of area.

ANTS: Spray ant trails, hills, and other areas of entry and infestation. Directly spray ants where possible.

SPIDERS: Directly spray spiders and webs.

SOWBUGS, MILLIPEDES, BEETLES, EARWIGS, Pillbugs, Centipedes, Scorpions, Cockroaches, Crickets, Waterbugs, and Palmetto Bugs: Thoroughly spray surfaces and into hiding places such as cracks and crevices, moist areas, openings around pipes and spigots, in storage areas and sheds, and other areas where these pests tend to congregate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool, dry place. Always store pesticides in the original container. Store away from food and pet food.

PESTICIDE DISPOSAL:

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING AND DISPOSAL:

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill.

Triple rinse as follows [for containers of 5 gallons (18.9 L) or less]: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse as follows [for containers greater than 5 gallons (18.9 L)]: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

[For totes/55 gal containers]

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Or for refillable containers]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into rinsate collection system. Repeat this rinsing procedure two more times. Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.



NSF Listed. NSF No. [137614][137615]



PYROCID[®] Concentrate 74406

[RESIDENTIAL SECTION OF LABEL]

[] or () indicates alternate or optional wording

[Contains the botanical insecticide Pyrethrum [an extract of a Chrysanthemum flower]]
 [A micro-emulsion, water-based spray]
 [Kills and repels Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Mosquitoes, and Gnats] [Provides rapid kill of mosquitoes and other listed insects without objectionable residues.] [For use with [Brand Name], [sold separately]]
 [Kills Black Widow Spiders and Black Carpenter Ants]
 [Quick killing Pyrethrum controls crawling and flying insects in the home, schools, hospitals and other listed areas]

[Controls Fleas and Ticks indoors and around pet areas]
 [Controls Deer Ticks and other Ixodid species that may carry and transmit Lyme disease]
 [Kills Mosquitoes, including the Culex species that may carry and transmit the West Nile virus]
 [Kills mosquitoes that may transmit Zika virus]
 [Kills Aedes mosquitoes]
 [Kills mosquitoes, including the Aedes mosquito that may carry and transmit Zika virus]

ACTIVE INGREDIENTS:

Pyrethrins	3.00%
* Piperonyl butoxide	30.00%
OTHER INGREDIENTS	67.00%
	<u>100.00%</u>

*(butylcarbityl) (6-propylpiperonyl) ether and related compounds

PYROCID[®], MGK[®] - registered trademarks of McLaughlin Gormley King Company

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	

Manufactured by

 7325 Aspen Lane North
 Minneapolis MN 55428

EPA Reg. No. 1021-1800

EPA Est. No. 1021-MN-2

Net Contents 1 Gallon. [2 Quarts] [1 Quart]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through runoff. This product has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use this product in or on electrical equipment because of shock hazard.

[Pyrocyde® Concentrate 74406] contains the quick killing botanical insecticide, Pyrethrum, [an extract of Chrysanthemum flowers], which provides rapid effective control of insects upon direct contact with an added benefit of flushing [repelling] these insects to aid in a more complete control [kill]. This highly effective water-based formula can be used on flying and crawling insects indoors and outdoors.

[Pyrocyde Concentrate 74406] can be used to control fleas and ticks hidden in carpets, rugs, upholstery, draperies, floor cracks and cabinets. Also controls Deer Ticks and other Ixodid species that may carry and transmit Lyme disease.

[Pyrocyde Concentrate 74406] can be used in the home, schools, hospitals, restaurants and other listed areas to control [kill], on contact, listed crawling pests and is a fast, effective product for flying insect control. Pyrethrum has quick killing ability with no persistent residue.

[Pyrocyde Concentrate 74406] can be used to reduce annoyance of mosquitoes, including the Culex species that may carry and transmit the West Nile virus, outdoors around homes, in recreational areas, and other listed areas. Ideal for use before outdoor activities.

[Pyrocyde Concentrate 74406] can be used in food and non-food areas of food processing plants, canneries, industrial installations, bakeries, rice and wheat mills, tobacco warehouses, grain elevators, warehouses, storage sites and other listed areas to control [kill], on contact, listed crawling pests and is a fast, effective product for flying insect control.

[Pyrocyde Concentrate 74406] provides rapid kill of insects without objectionable residues.

To protect your pet against and to minimize reintroduction of adult fleas from outdoors, use EPA registered flea and tick products, ([such as] flea or flea and tick collar, pet powder, pet spray, pet dip, pet shampoo, flea shampoo or flea and tick pet shampoo), in conjunction with this application and prior to re-entry.

Remove pets and birds and cover fish aquariums before spraying.

FOR THE CONTROL OF INSECTS including: Almond Moths, Angoumois Grain Moths, Ants, Bed Bugs, Black Carpenter Ants, Clothes Moths, Clover Mites, Black Flies, Black Widow Spiders, Blood Sucking Lice, Boxelder Bugs, Brown Dog Ticks, Caddises, Carpet Beetles, Cat Fleas, Cellar Spiders, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Dark Mealworms, Darkling Beetles, Deer Flies, Deer Ticks, Dermestid Beetles, Dog Fleas, Driedfruit Beetles, Drugstore Beetles, Earwigs, Face Flies, Fannia Flies, Firebrats, Flat Grain Beetles, Fleas, Flies, Fruit Flies, Fungus Gnats, Gnats, Grain Mites, Granary Weevils, Hide Beetles, Horn Flies, Hornets, Horse Flies, House Flies, Indian Meal Moths, Khapra Beetles, Lesser Grain Borers, Lesser House Flies, Lice, Mediterranean Flour Moths, Merchant Grain Beetles, Midges, Miller Moths, Millipedes, Mosquitoes – including the Culex species that may carry and transmit the West Nile virus, Palmetto Bugs, Pillbugs, Poultry Lice, Red Flour Beetles, Rice Flour Beetles, Rice Weevils, Rusty Grain Beetles, Sawtoothed Grain Beetles, Sheep Ticks, Silverfish, Skipper Flies, Small Flying Moths, Sowbugs, Spiders, Spider Beetles, Stable Flies, Ticks that may carry and transmit Lyme disease, Tobacco Moths, Trogoderma Beetles, Warehouse Beetles, Wasps, Waterbugs, Yellow Mealworms and Yellowjackets.

INDOOR AREAS for use in: Animal Areas, Apartments, Attics, Automobiles, Basements, Bathrooms, Bedding, Bedrooms, Boats, Bookcases, Cabinets, Cabins, Camp Sites, Campers, Carpets, Cat Sleeping Quarters, Chests, Closed Porches, Closets, Clothes Storage, Club Houses, Condominiums, Craw Spaces, Credenzas, Decks, Dens, Dog Houses, Dog Sleeping Quarters, Dormitories, Draperies, Dressers, Dwellings, False Ceilings, Floors, Food Storage Areas, Furniture, Garages, Garbage Bins, Garbage Cans, Garbage Compactors, Gazebos, Homes, Household Contents, Houses, Kennels, Kitchens, Lanais, Lavatories, Libraries, Living Rooms, Mattresses, Mini Storage, Mobile Homes, Patios, Pet Bedding, Pet Carriers, Pet Sleeping Areas, Play Houses, Play Rooms, Porches, Public Storage, Recreational Vehicles, Rooms, Rugs, Solariums, Storage Areas, Sun Porches, Sun Rooms, Tents, Tool Sheds, Trailers, Tree Houses, Verandas and Wine Cellars.

OUTDOOR AREAS for uses in and around: Ant Hills, Back Yards, Balconies, Front Yards, Gazebos, Kennels, Patios, Rooftops, Sidewalks, Side Yards and Tennis Courts.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product
in a manner inconsistent with its labeling.

READ ALL DIRECTIONS COMPLETELY BEFORE USE. For best results, follow directions for specific use areas.

[For use in the cordless, manually activated [Brand Name], sold separately]

RESTRICTIONS:

- **ENTRY RESTRICTIONS:**
 - For Surface Spray Use: Do not allow adults, children, or pets to enter the treated area until sprays have dried.
 - For Space Spray Use: Do not allow adults, children or pets to enter the treated area until sprays have settled.
- **APPLICATION RESTRICTIONS:**
 - Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift.
 - Remove pets, birds, and cover fish aquariums before spraying.
 - Do not make applications during rain.
 - Do not remain in treated area. Exit area immediately and remain outside the treated area until aerosols, vapors, and/or mists have dispersed.
 - Do not wet plants to the point of runoff or drip
 - Do not water the treated area to the point of runoff.
 - Do not wet articles to point of runoff or drip.
 - Remove or cover exposed food and drinking water before application.
 - Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces or wash them before use.
 - Except for use with fogging devices, all outdoor applications must be limited to spot or crack and crevice treatments only, except for the following permitted uses:
 - Treatment to soil or vegetation around structures;
 - Applications to lawns, turf, and other vegetation;
 - Applications to building foundations, up to a maximum height of 3 feet.
 - Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack and crevice applications, only.

GENERAL: This concentrated product must be diluted with water. When making dilutions, add concentrate to the tank and add water. Agitate well while adding water and apply as directed.

- 0.04% Pyrethrins – add 1.6 fl. oz. of concentrate / gallon of water (13 ml/L)
- 0.05% Pyrethrins – add 2 fl. oz. concentrate / gallon of water (15.86 ml/L)
- 0.10% Pyrethrins – add 4.3 fl. oz. concentrate / gallon of water (34.10 ml/L)
- 0.30% Pyrethrins – add 13 fl. oz. concentrate / gallon of water (103.09 ml/L)
- 0.50% Pyrethrins – add 22 fl. oz. concentrate / gallon of water (174.46 ml/L)

For maximum effectiveness, following the dilution chart and using a combination of spot, surface treatment and fogging are recommended.

SURFACE SPRAYING: To kill Cockroaches, Black Carpenter Ants, Silverfish, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles and other listed insects, use a sprayer adjusted to a coarse wet spray. Mix well and apply as a coarse wet spray, using approximately 1 gallon of spray mixture per 1,000 sq. ft. of surface. To provide maximum penetration and impingement on the insects, direct the spray first into all cracks and crevices, behind all equipment, cupboards, obstructions or dark harborage areas, on walls and floors, underneath sinks, behind pipes, beneath stoves and refrigerators, around garbage cans and cabinets, along baseboards, door sills, window sills, door and window frames and floors, around and on drains, pipes, and plumbing, behind bookcases, storage and other utility installation areas, and in all places that harbor insects. Contact as many insects as possible. Do not allow spray to contact stored food. If surface application only is to be used, spray floor and other surfaces. To kill Ants, treat doors and around window frames and other areas of entry. Spray on Ants whenever possible. Apply at the rate of 28 fl. oz. of concentrate/gallon of water per 1,000 sq. ft. For surface spraying or crack and crevice treatments do not exceed maximum application rate of 28 fl. oz. of concentrate per 1,000 sq. ft.

CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply liberally to empty chests, closets, bureaus, and other storage areas, directing the spray into cracks, joints, and crevices. Apply at a rate of 28 fl. oz. of concentrate/gallon per 1000 sq. ft. Do not exceed the maximum application rate of 0.056 lbs. pyrethrins per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1000, sq. ft.

CARPET BEETLES: Directly spray as many insects as possible. Spray all surfaces of upholstered furniture. Where possible, thoroughly spray both top and bottom of rugs and carpets. Also treat shelves, closets, floors, walls, windowsills, baseboards, areas under furniture, and wherever these insects are seen or suspected. Apply at a rate of 28 fl. oz. of concentrate/gallon per 1,000 sq. ft. Do not exceed the maximum application rate of 0.056 lbs. pyrethrins per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1,000 sq. ft.

BED BUGS: Spray mattresses lightly, paying special attention to tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, floors, walls, furniture, and other infested areas. Apply at a rate of 28 fl. oz. of concentrate/gallon per 1000 sq. ft. Do not exceed the maximum application rate of 0.056 lbs. pyrethrins per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1000, sq. ft.

FOR USE IN LISTED PET AREAS: For best results remove, destroy, or wash pet's old bedding. To reduce infestation, thoroughly spray pet [dog] [cat] [puppy] [kitten] [animal] bedding, sleeping quarters, kennel, around baseboards, window and door frames, into wall cracks, on local areas of floors, and other animal resting places until the surface is slightly moist. Treat these areas at the rate of two seconds per linear foot at the distance of 12 to 18 inches (30.5 to 45.7 cm). Also, spray the entire inside surface of doghouses. Do not exceed the maximum application rate of 28 fl. oz. of concentrate per 1,000 sq. ft. Put fresh bedding in pet's quarters after spray has dried. Apply at a rate of 28 fl. oz. of concentrate/gallon of water per 1,000 sq.ft. Allow treated areas to dry thoroughly before allowing pets to re-enter.

Do not exceed the maximum application rate of 0.056 lbs. Pyrethrins/ 1,000 sq. ft. or 0.56 lbs. PBO per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1,000 sq. ft.

FLEA AND TICK INFESTATION ON RUGS AND CARPETS: Before use, vacuum all infested areas thoroughly and dispose of vacuum bag in outside trash. Spray rugs and carpets where fleas tend to hide and usually where animals frequent. Spray thoroughly into hiding places such as cracks and crevices, in walls, on floors, and along baseboards, window and door frames. Spray at the rate of one sq. ft. per second. To kill ticks and other labeled insects, apply directly to bugs. Apply at a rate of 28 fl. oz. of concentrate/gallon per 1000 sq. ft. Do not exceed the maximum application rate of 0.056 lbs. pyrethrins per 1,000 sq. ft. or 28 fl. oz. of concentrate per 1000, sq. ft Do not apply directly to pets. Treat pets with an EPA registered [brand name] product in conjunction with this application.

For best results, to protect your pet against and to minimize reintroduction of adult fleas from outdoors, use EPA registered flea and tick products, ([such as] [brand name] flea or flea and tick collar, [brand name] pet powder, [brand name] pet spray, [brand name] pet dip, [brand name] pet shampoo, flea shampoo or flea and tick pet shampoo), in conjunction with this application. Do not exceed maximum application rate of 28 fl. oz. of concentrated product per 1,000 sq. ft.

FOR KENNELS AND OTHER RESIDENTIAL OUTDOOR ANIMAL AREAS: Apply at the rate of 1 fl. oz / gallon of water per 1,000 sq. ft. . Apply liberally to surfaces, [paying] [with] special attention to cracks and crevices and other insect hiding places. Allow treated areas to dry thoroughly before allowing pets [animals] to re-enter. Do not exceed maximum application rate of 0.002 lbs. pyrethrin per 1,000 sq. ft. or 1 fl. oz. of concentrated product per 1,000 sq. ft.

DIRECTIONS FOR USE IN OUTDOOR RESIDENTIAL MISTING SYSTEMS [Unit / Misters]:

NOTE: NOT REGISTERED for use in Automatic ULV Spraying Systems in the State of New York.

Use to control or temporarily reduce annoyance from accessible stages of: Mosquitoes including the Culex species that may carry and transmit the West Nile virus, Flies, Gnats, Face Flies, Fannia Flies, Deer Flies, Horse Flies, Stable Flies, Horn Flies, Hornets, Wasps, Yellowjackets, Firebrats, Small Flying Moths, Earwigs, Fleas, Ticks, Clover Mites, Brown Dog Ticks, Crickets, Cockroaches, Silverfish, Spiders, Darkling Beetles, and other nuisance insects.

- Do not apply this pesticide when people, pets and food are present.
- Do not use in an evaporative cooling system.
- Do not set nozzles to direct mist near outside air conditioner systems or other home air intakes.
- Direct nozzles to spray towards the target area and away from swimming pools, water bodies, or eating and cooking areas.
- When using this product, installers and service technicians must comply with the license, certification, or registration requirements of the state(s), tribe(s), or local authority(ies) where they are installed.
- If the system works when a person operates a remote activation device, then application of this pesticide when people, pets and food are present is prohibited.
- If system works on an automatic timer, set the timing for application when people, pets and food are unlikely to be present.
- Do not use in misters located within 3 feet of air vents, air conditioner units, or windows.
- If used in a system with a reservoir tank for the end-use dilution, the system reservoir tank must be locked. Securely attach the end-use pesticide label and a dilution statement to the system reservoir tank in a weather protected area or plastic sleeve. The dilution statement must be phrased as follows: This container holds ___ parts [product name] to ___ parts water.
- If used in a direct injection system, the pesticide container must be locked. Securely attach the end-use label to the pesticide container in a weather protected area or plastic sleeve.
- Contact the system manufacturer for guidance or assistance to ensure compliance with the following requirement.
- This product must only be used in systems that have been calibrated to apply no more than the maximum application rate of 1 fl. oz. per minute.

Automatic Spraying System Dilution Table

A. System Tank size (in gallons)	B. Conversion Factor (fl. oz. of concentrate per gallon of water)	A. x B. = fl. oz. of concentrate for that tank size, to be mixed with water to achieve 0.044% pyrethrin solution.
30 (113.7 L)	1.88 (55.59 ml)	56.40 (1,668 ml)
55 (208.4 L)	1.88 (55.59 ml)	103.40 (3,058 ml)
125	1.88 (55.59 ml)	235 (6,949 mls)
250	1.88 (55.59 ml)	470 (13,898 mls)

When filling tank, partially fill the system [unit / mister] reservoir tank with water and add concentrate. Then fill the rest of the tank with water and agitate well.

Contact the system manufacturer for guidance or assistance to ensure compliance with the following requirement:

This product must only be used in systems [unit / misters] that have been calibrated to apply no more than the maximum application rate of 0.00011 lbs. (0.05 g) a.i. Pyrethrins per 1,000 cu. ft. per day or 0.00058 lbs. of Piperonyl Butoxide per 1,000 cu. ft. per day, whichever is lower (do not exceed more than 2 fl. oz. of diluted product per 1,000 cu. ft. per day).

For best performance, the spray solution should be in the pH range of 5.5 to 7.0.

This product will control insects directly contacted by its spray solution. Since many of the targeted insects can migrate in and out of treatment areas, it is recommended that this concentrate be used as part of an Integrated Pest Management (IPM) program. Control practices including eliminating breeding and harborage sites combined with appropriate use of contact insecticides, Insect Growth Regulators (IGRs) and / or residual insecticides will provide the most effective control of target insects.

OUTDOOR USE DILUTION TABLE: Follow dilution chart and apply as directed.

Solution percentage	Ratio of concentrate to water	Fl. oz. of concentrate per gallon of water
0.02% Pyrethrins	1 part concentrate to 128 parts water	1 fl. oz of concentrate / gallon of water (7.8 ml/L) Apply at the rate of 1 gallon / 1,000 sq. ft.

Take care to avoid spraying when wind conditions might create a mist that can blow back to applicator. Do not apply this product in or around bodies of water. Do not spray directly on plants. Spray surfaces and contact as many insects as possible.

SMALL FLYING INSECTS AND OTHER FLYING INSECTS including FLIES, GNATS, MOSQUITOES, SMALL FLYING MOTHS, and MIDGES: Apply at a rate of 1.0 fl. oz. of concentrate/gallon of water per 1,000 sq ft. of area. Hold sprayer nozzle 18 to 24 inches (45.72 to 60.96 cm) from shrubs, bushes, grassy areas, and other areas infested with these pests. Do not apply directly to plants. Spray with a slow sweeping motion while moving away from the treated area. To reduce annoyance from these pests, treat open porches, patios and under decks in a similar manner. Kills and repels Flies, Mosquitoes including the Culex species that may carry and transmit the West Nile virus, Midges, and Gnats when used intermittently during outdoor activities and will keep surrounding areas free from these pests. Allow spray to dissipate before occupying the treated area. Do not exceed 0.002 lb. pyrethrins or 1 fl. oz. of concentrate per 1000 sq. ft. of area.

ANTS: Spray ant trails, hills, and other areas of entry and infestation. Directly spray ants where possible.

SPIDERS: Directly spray spiders and webs.

SOWBUGS, MILLIPEDES, BEETLES, EARWIGS, Pillbugs, Centipedes, Scorpions, Cockroaches, Crickets, Waterbugs, and Palmetto Bugs: Thoroughly spray surfaces and into hiding places such as cracks and crevices, moist areas, openings around pipes and spigots, in storage areas and sheds, and other areas where these pests tend to congregate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store in a cool, dry area. Always store pesticides in the original container. Store away from food and pet food.

PESTICIDE DISPOSAL AND CONTAINER HANDLING

Non-refillable container. Do not reuse or refill this container.

If empty:

Place in trash or offer for recycling if available.

If partly filled:

Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.