

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

1021-1807

6/23/2009

1/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460-0001

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

June 23, 2009

Mr. Thomas A. Lennan, Registration Specialist  
McLaughlin Gormley King Company  
8810 Tenth Avenue North  
Minneapolis, MN 55427-4319

**Subject: Amended Labeling, First Food Use on All Food/Feed Crops Following Wide-Area Mosquito  
Adulticide Application (PP 7F7251)  
MGK® MULTICIDE® Fogging Concentrate 2807, EPA Reg. No. 1021-1807  
Your E-Mail Resubmission Dated June 18, 2009**

Dear Mr. Lennan:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

McLaughlin Gormley King Company should note that a notice of availability of EPA's RED for products containing piperonyl butoxide was published in the *Federal Register* on July 26, 2006 (Volume 71, Number 143, Pages 42385-42386). As a result, additional data and/or labeling changes may be required for products containing piperonyl butoxide.

A copy of your label stamped "Accepted" is enclosed for your records. If you have any questions about this label review, please contact Mr. Carmen Rodia at (703) 306-0327 or via e-mail at [Rodia.Carmen@epa.gov](mailto:Rodia.Carmen@epa.gov).

Sincerely Yours,

FOR

Richard J. Gebken  
Product Manager (10)  
Insecticide Branch  
Registration Division (7505P)

Enclosure: Copy of label stamped "Accepted"  
001021-01807 D415338



8810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4319 U.S.A. - Telephone (763) 544-0341

[ ]denotes alternate language

[A water based formulation]

For use only by federal, state, tribal or local government officials responsible for public health or vector control, or by persons certified in the appropriate category or otherwise authorized by the state or tribal lead pesticide regulatory agency to perform adult mosquito control applications, or by persons under their direct supervision.

ACTIVE INGREDIENTS:

Sumithrin®: 3-Phenoxybenzyl-(1RS, 3RS, 1RS, 3SR)-2,2-dimethyl-3-(2-methylprop-1-enyl) Cyclopropanecarboxylate .....	10.00%
* Piperonyl Butoxide, Technical .....	10.00%
OTHER INGREDIENTS .....	80.00%
	<u>100.00%</u>

\* Equivalent to 8.00% (butylcarbityl) (6-propylpiperonyl) ether and 2.00% related compounds  
 Contains 0.847 pounds of Technical SUMITHRIN®/Gallon and 0.847 pounds Technical Piperonyl Butoxide/Gallon.

MULTICIDE® - Registered trademark of McLaughlin Gormley King Company  
 Sumithrin™ - Trademark of Sumitomo Company Ltd.

**CAUTION**

KEEP OUT OF REACH OF CHILDREN  
 PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> <li>▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>▪ Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and aquatic invertebrates. Runoff from treated areas or deposition of spray droplets into a body of water may be hazardous to fish and aquatic invertebrates. Do not apply over bodies of water (lakes, rivers, permanent streams, natural ponds, commercial fish ponds, swamps, marshes or estuaries), except when necessary to target areas where adult mosquitoes are present, and weather conditions will facilitate movement of applied material away from the water in order to minimize incidental deposition into the water body. Do not contaminate bodies of water when disposing of equipment rinsate or washwaters.

BEE WARNING: This product is toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply to or allow drift onto blooming crops or weeds when bees are visiting the treatment area, except when applications are made to prevent or control a threat to public and/or animal health determined by a state, tribal or local health or vector control agency on the basis of documented evidence of disease causing agents in vector mosquitoes or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Before making the first application in a season, it is advisable to consult with the state or tribal agency with primary responsibility for pesticide regulation to determine if other regulatory requirements exist.

IN CALIFORNIA: This product is to be applied by County Health Department, State Department of Health Services, Mosquito and Vector Control or Mosquito Abatement District personnel only.

Net Contents \_\_\_\_\_

Manufactured by:  
 Mc LAUGHLIN GORMLEY KING COMPANY  
 8810 Tenth Avenue North  
 Minneapolis, MN 55427

EPA Reg. No. 1021-1807

ACCEPTED  
 EPA Est. No. 1021-MN-2

June 23, 2009  
 Under the Federal Insecticide, Fungicide,  
 and Rodenticide Act, as amended, for the  
 pesticide registered under:

EPA. Reg. No: 1021-1807

FW

Wear long sleeved shirt and long pants, socks and shoes.

MULTICIDE® Fogging Concentrate 2807 cannot be mixed with oil. Use only clean water.

SHAKE OR AGITATE THOROUGHLY BEFORE DILUTING. Diluted product must be thoroughly agitated before use. Avoid storing excess formulation in equipment.

**USE AREAS:** For use in mosquito adulticiding programs involving outdoor residential and recreational areas where adult mosquitoes are present in annoying numbers, and in vegetation surrounding parks, woodlands, swamps, marshes, overgrown areas and golf courses. For best results, apply when mosquitoes are most active and meteorological conditions are conducive to keeping the spray cloud close to the ground. Application in calm air conditions is to be avoided. Apply only when ground wind speed is greater than 1 mph. Air temperature should be greater than 50°F when conducting all types of applications.

MULTICIDE Fogging Concentrate 2807 may be applied over agricultural areas prior to harvest for the control of adult mosquitoes within or adjacent to these areas.

Do not treat a site with more than 0.0036 pounds of sumithrin per acre in a twenty-four hour period. Do not exceed 1.0 pounds of Sumithrin® per acre in any site in any year. More frequent applications may be made to prevent or control a threat to public and/or animal health determined by a state, tribal, or local health or vector control agency on the basis of documented evidence of disease causing agents in vector mosquitoes or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort.

Do not contaminate food, feed or drinking water. In treatment of corrals, feed lots, swine lots and zoos, cover any exposed drinking water, drinking water fountains and animal feed before application.

**SPRAY DROPLET SIZE DETERMINATION:**

**Ground-Based Application:** Spray equipment must be adjusted so that the volume median diameter (VMD) is 8 to 30 microns ( $8 \leq D_v 0.5 \leq 30 \mu m$ ) and that 90% of the spray is contained in droplets smaller than 50 microns ( $D_v 0.9 < 50 \mu m$ ). Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using a laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated.

**Aerial Application:** Spray equipment must be adjusted so that the volume median diameter produced is less than 60 microns ( $D_v 0.5 < 60 \mu m$ ) and that 90% of the spray is contained in droplets smaller than 115 microns ( $D_v 0.9 < 115 \mu m$ ). The effects of flight speed and, for non-rotary nozzles, nozzle angle on the droplet size spectrum must be considered. Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using a wind tunnel and laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated.

**GROUND U.L.V. APPLICATION:** Apply MULTICIDE® Fogging Concentrate 2807 undiluted at a flow rate of 1.1 to 3.3 fluid ounces per minute at an average vehicle speed of 10 mph using a swath width of 300 feet for acreage calculations (see chart below). Under normal residential conditions a flow rate of 2.2 fluid ounces per minute is recommended. If a different vehicle speed is used, adjust rate accordingly. These rates are equivalent to 0.0012 to 0.0036 pounds of Sumithrin® and Piperonyl butoxide per acre. Vary flow rate according to vegetation density and mosquito population. Use high flow rate in heavy vegetation or when populations are high. For proper application, mount the applicator so the nozzles is at least 4 ½ feet above ground

level and directed out the back of the vehicle. Failure to follow the above directions may result in reduced effectiveness.

MULTICIDE® Fogging Concentrate 2807 may also be diluted with water and applied so as not to exceed the maximum pounds of active ingredient per acre as shown in the tables below. If an alternative dilution rate is used, adjust the flow rate accordingly, provided the droplet size and spectrum remains within label specifications.

Rates to use MULTICIDE® Fogging Concentrate 2807 Undiluted for Mosquito Control:

Pounds a.i./Acre		MFC 2807 Fl. oz./Acre
Sumithrin®	PBO	
0.0036	0.0036	0.54
0.0024	0.0024	0.36
0.0012	0.0012	0.18

Flow Rates in fluid oz./minute at truck speeds of:			
5 MPH	10 MPH	15 MPH	20 MPH
1.6	3.3	4.9	6.6
1.1	2.2	3.3	4.4
0.5	1.1	1.6	2.2

Rates to use diluted (1 to 1 dilution ratio) MULTICIDE® Fogging Concentrate 2807

Pounds a.i./Acre		Finished Spray Fl. oz./Acre
Sumithrin®	PBO	
0.0036	0.0036	1.09
0.0024	0.0024	0.73
0.0012	0.0012	0.36

Flow Rates in fluid oz./minute at truck speeds of:			
5 MPH	10 MPH	15 MPH	20 MPH
3.3	6.6	9.9	13.2
2.2	4.4	6.6	8.8
1.1	2.2	3.3	4.4

Rates to use diluted (1 to 2 dilution ratio) MULTICIDE® Fogging Concentrate 2807 for Mosquito Control

Pounds a.i./Acre		Finished Spray Fl. oz./Acre
Sumithrin®	PBO	
0.0036	0.0036	1.63
0.0024	0.0024	1.09
0.0012	0.0012	0.54

Flow Rates in fluid oz./minute at truck speeds of:			
5 MPH	10 MPH	15 MPH	20 MPH
4.9	9.9	14.8	19.8
3.3	6.6	9.9	13.2
1.6	3.3	4.9	6.6

MULTICIDE® Fogging Concentrate 2807 may also be applied undiluted with non-thermal, portable, motorized backpack equipment adjusted to deliver ULV particles of 50 to 100 microns

h/c

VMD. Use 0.18 to 0.54 fl.oz. of the undiluted spray per acre (equal to 0.0012 to 0.0036 lb.ai./acre) as a 50 ft (15.2 m) swath while walking at a speed of 2 mph (3.2 kph).

MULTICIDE® Fogging Concentrate 2807 may be applied for urban ULV mosquito control. For control of resting or flying adult mosquitoes, biting flies and biting midges in areas such as utility tunnels, sewers, storm drains and catch basins, pipe chases, underground basements, underground passages, parking decks, crawl spaces or uninhabited buildings. MULTICIDE® Fogging Concentrate 2807 may be applied using mechanical foggers, hand-held or truck mounted ULV equipment, non-thermal foggers or other spray equipment suitable for this application. Apply MULTICIDE® Fogging Concentrate 2807 at rates up to but not exceeding 0.0036 pounds sumithrin per acre per twenty-four hour period (not to exceed 1.0 pounds Sumithrin per acre in any site in any year).

**AERIAL APPLICATION:**

FLORIDA: Do not apply by aircraft unless approved by the Florida Department of Agriculture and Consumer Services.

MULTICIDE® Fogging Concentrate 2807 may be applied at rates of 0.18 to 0.54 fluid ounces MULTICIDE® Fogging Concentrate 2807 per acre by fixed wing or rotary aircraft equipped with suitable ULV application equipment. MULTICIDE® Fogging Concentrate 2807 may also be diluted with water and applied by aerial ULV equipment so long as 0.54 fluid ounces per acre of MULTICIDE® Fogging Concentrate 2807 is not exceeded. Application shall be made at 75 to 300 feet for rotary aircraft and 100 to 300 feet for fixed-wing aircraft.

Reference the dilution charts on this label. If an alternative dilution rate is used, adjust the flow rate accordingly, provided the droplet size and spectrum remains within label specifications.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE:

Store in a cool, dry place. Keep container closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable container. Do not reuse or refill this container.

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a rinse tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available for reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

[Or for containers over 5 gallons:]

[Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.] Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

[Refillable Containers:]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into rinsate collection system. Repeat his rinsing procedure two more times. Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**NEXT**

**LABEL**

1021-1807

6/23/2009

1/2



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460-0001

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

June 23, 2009

Mr. Thomas A. Lennan, Registration Specialist  
McLaughlin Gormley King Company  
8810 Tenth Avenue North  
Minneapolis, MN 55427-4319

**Subject: Supplemental Labeling, First Food Use on All Food/Feed Crops Following Wide-Area Mosquito Adulticide Application (PP 7F7251) MGK® MULTICIDE® Fogging Concentrate 2807, EPA Reg. No. 1021-1807 Your E-Mail Submission Dated June 11, 2009**

Dear Mr. Lennan:

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable subject to the comment listed below.

**"This supplemental label is valid for a period of 18 months from the date of this letter."**

McLaughlin Gormley King Company should note that a notice of availability of EPA's RED for products containing piperonyl butoxide was published in the *Federal Register* on July 26, 2006 (Volume 71, Number 143, Pages 42385-42386). As a result, additional data and/or labeling changes may be required for products containing piperonyl butoxide.

A copy of your supplemental label stamped "Accepted" is enclosed for your records. If you have any questions, please contact Carmen Rodia at (703) 306-0327 or via e-mail at [Rodia.Carmen@epa.gov](mailto:Rodia.Carmen@epa.gov).

Sincerely Yours,

FOR

Richard J. Gebken  
Product Manager (10)  
Insecticide Branch  
Registration Division (7505P)

Enclosure: Copy of supplemental label stamped "Accepted"  
001021-01807 D415338

2  
/2

**MGK**<sup>®</sup>

**McLAUGHLIN  
GORMLEY  
KING COMPANY**

MULTICIDE<sup>®</sup> Fogging Concentrate 2807

8810 Tenth Avenue N./ Minneapolis, Minnesota 55427-4319 U.S.A. Telephone (763) 544-0341

**SUPPLEMENTAL LABEL**

READ THE LABEL AFFIXED TO THE CONTAINER FOR MULTICIDE<sup>®</sup> Fogging Concentrate 2807, EPA Reg. No. 1021-1807, BEFORE APPLYING. USE OF THIS PRODUCT ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR THIS PRODUCT. THE LABELLING MUST BE IN POSSESSION OF THE USER AT THE TIME OF APPLICATION.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MULTICIDE<sup>®</sup> Mosquito Adulticiding Concentrate 2706 may be applied over agricultural areas for the control of adult mosquitoes within or adjacent to the treatment areas.

This supplemental label supersedes the prohibition to apply this product over agricultural areas for the control of mosquitoes on the label attached to the container.

Manufactured by:  
Mc LAUGHLIN GORMLEY KING COMPANY  
8810 Tenth Avenue North

EPA Reg. No. 1021-1807

Minneapolis, MN 55427

EPA Est. No. 1021-MN-2

**ACCEPTED**

*June 23, 2009*

**Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended, for the  
pesticide registered under:**

**EPA. Reg. No:** 1021-1807