

1021-1815

06/13/2006

117



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

June 13, 2006

Ms. Julie B. Schlekau,
Registration Specialist
McLaughlin, Gormley, and King Company
8810 Tenth Avenue North
Minneapolis, MN 55427-4319

Subject: Amendment: Extend residual claim for cockroaches on label from 4 to 12 months
✓ Evercide Esfenvalerate 6.4% CS (EPA Reg. Nos. 1021-1815)
Evercide Esfenvalerate 0.025% RTU CS (EPA Reg. Nos. 1021-1816)
Applications dated January 18, 2006

Dear Ms. Schlekau:

The above referenced submissions, in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, are acceptable with the following comments:

- **Front Panel:** For both products, revise the statement "Kills Cockroaches for up to 12 month (1 year) on concrete" to read "***Kills for up to 12 months (1 year) ****"; and define or clarify with an asterisk directly below the claim(s) – "****For German Cockroaches on indoor non-porous surfaces only***".

Please make the above stated changes to the products' labeling and submit two copies of each revised final printed label to the Agency. A stamped copy of each label is enclosed for your record.

If you have any questions please contact Dr. B.A. Akinlosotu at (703) 605-0653.

Sincerely,

A handwritten signature in black ink, appearing to read "George T. LaRocca".

George T. LaRocca,
Product Manager 13
Insecticide Branch
Registration Division (7505C)

Enclosure



McLAUGHLIN
GORMLEY
KING COMPANY

EVERCIDE® Esfenvalerate 6.4% CS
(Microencapsulated Insecticide)

For Pest Management Professional Use

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4372 U.S.A. - Telephone (763) 544-0341

Kills Cockroaches for up to 12 months (1 year) ~~on concrete indoor non-porous surfaces.~~
*FA German Cockroaches, on ~~indoor non-porous surfaces only.~~
ACTIVE INGREDIENT:

(S)-cyano(3-phenoxyphenyl)methyl-(S)-4-chloro-alpha-(1-methylethyl)benzeneacetate	6.40%
*OTHER INGREDIENTS	93.60%
*Contains petroleum distillates	100.00%

EVERCIDE® - Registered trademark of McLaughlin Gormley King Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN: Contains petroleum distillate - vomiting may cause aspiration pneumonia. Have the product container or label with you when calling a poison control center or doctor or ng for treatment. For information regarding medical emergencies or pesticide incidents, call 888-740-8712.	

Net Contents _____
Manufactured by:
Mc LAUGHLIN GORMLEY KING COMPANY
8810 Tenth Avenue North
Minneapolis, MN 55427

EPA Reg. No. 1021-1815

EPA Est. No. 1021-MN-2

ACCEPTED
with COMMENTS
In EPA Letter Dated:

JUN 13 2006

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
1021-1815**

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear protective eyewear (goggles, face shield or safety glasses with side shields). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and other aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirement of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Remove pets and birds and cover fish aquariums before spraying.

For control of insects, indoors and outdoors, in food and non-food areas such as, but not limited to: Homes, Schools, Warehouses, Office Buildings, Apartment Buildings, Theatres, Hotels, Industrial Buildings, Motels, Kennels, Livestock Housing, Food Processing Plants, Food Service Establishments, Restaurants, Supermarkets and Grocery Stores, Transportation Equipment, Truck Trailers, Railroad Cars, and Food Manufacturing and Warehousing Establishments. Also for use on Backyards, Lawns, Trees, Ornamental Landscaping, Recreational Areas, Parks and Athletic Fields.

Insects Controlled:

Ants	Fire Ants	Lice	Leaf Rollers
Aphids	Fleas	Litter Beetles	Leaf Tiers
Asian Lady Beetles	Gnats	Mites	Leafhoppers
Bedbugs	Hornets	Mosquitoes	Lesser Appleworms
Beetles	Lice	Stable Flies	Lesser Peach Tree Borers
Boxelder Bugs	Mosquitoes		Loopers

217

Carpet Beetles	Scorpions	Wood-Destroying Pests including:	Lygus Bugs
Centipedes	Spiders	Carpenter Ants	Mexican Bean Beetles
Cockroaches	Ticks	Carpenter Bees	Mites
Crickets	Wasps	Deathwatch Beetles	Mole Crickets
Dog Ticks	Yellow Jackets	Furniture Beetles	Naval Orangeworms
Earwigs		Old House Borers	Northern Pine Weevils
Fire Ants	Stored Product Pests Including:	Powder Post Beetles	Oriental Fruit Moths
Firebrats	Angoumois Grain Moths	Round-headed House Borers	Painted Lady Caterpillars
Fleas	Ants Cadelles	Swarming Termites	Pea Weevils
Mealy Bugs	Cigarette Beetles		Peach Tree Borers
Midges	Confused Flour Beetles	Plant Pests including:	Peach Twig Borers
Millipedes	Dark Mealworms	Alfalfa Caterpillars	Pear Psyllid
Mites	Dried Fruit Beetles	Artichoke Plum Moth	Pear Slugs
Pillbugs	Drugstore Beetles	American Plum Borers	Pecan Leaf Phylloxera
Psocids	Flat Grain Beetles	Annual Bluegrass Weevils	Pecan Nut Casebearers
Scales	Fruit Flies	Aphids	Pecan Spittlebugs
Silverfish	Grain Mites	Apple Maggots	Pecan Stem Phylloxera
Spiders	Grain Moths	Armyworms	Pecan Weevils
Springtails	Granary Weevils	Bagworms	Pepper Weevils
Swarming Termites	Indian Meal Moths	Balsam Woolly Adelgids	Periodical Cicadas
Ticks	Lesser Grain Borers	Beet Armyworms	Pickleworms
Waterbugs	Maize Weevils	Billbugs	Pillbugs
	Meal Moth Larvae	Blueberry Spanworms	Pine Chafers
Flying Insects Including:	Mediterranean Flour Moths	Cherry Fruit Flies	Pine Coreid Bugs
Blow Flies	Merchant Grain Beetles	Cherry Fruitworms	Plant Bugs
Chiggers	Red Flour Beetles	Chinch Bugs	Plum Curculios
Clothes Moths	Rice Weevils	Codling Moths	Red Pine Sawflies
Crane Flies	Rusty Grain Beetles	Cowpea Curculios	Red-striped Fireworms
Deer Flies	Saw-toothed Grain Beetles	Cranberry Fruitworms	Red-headed Pine Sawflies
Face Flies	Skippers	Cranberry Weevils	Rindworms
Fruit Flies	Spider Beetles	Cucumber Beetles	Salt Marsh Caterpillars
Fungus Gnats	Spider Mites	Cutworms	Sap Beetles
Gnats	Square-necked Grain Beetles	Diamondback Moths	Scales
Horn Flies	Tobacco Moths	Elm Leaf Beetles	Sod Webworms
Houseflies	Yellow Mealworms	European Pine	Sowbugs

		Sawflies	
Mosquitoes		Fall Webworms	Spiders
Small Flying Moths	Livestock Premise Pests:	Filbert Worms	Spittlebugs
Whiteflies	Bedbugs	Flea Beetles	Stink Bugs
	Carrion Beetles	Grasshoppers	Tarnished Plant Bugs
Biting and Stinging Pests:	Chiggers	Green Cloverworms	Tent Caterpillars
Bedbugs	Darkling Beetles (lesser mealworm)	Green Fruitworms	Thrips
Bees	Deer Flies	Gypsy Moth Caterpillars	Tobacco Hornworms
Biting Flies	Face Flies	Hickory Shuckworms	Tufted Apple Bud Moths
Chiggers	Fleas	Imported Cabbageworms	Velvet Bean Caterpillars
Deer Flies	Flies	Japanese Beetles	Walnut Husk Flies
Deer Ticks	Hide Beetles	Lace Bugs	Western Bean Cutworms
Dog Ticks	Horn Flies	Leaf Feeding Caterpillars	
	Horse Flies	Leaf Miners	Whiteflies

GENERAL INFORMATION:

Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm. Do not apply to classrooms while in use. Do not apply in institutions (including libraries, schools, sports facilities, etc.) in the immediate area where occupants are present.

EVERCIDE® Esfenvalerate 6.4% CS is a microencapsulated suspension concentrate containing 0.54 pounds active ingredient per gallon.

EVERCIDE® Esfenvalerate 6.4% CS is designed to provide long residual activity against pests listed on this label when applied as surface or spot treatments, injected into wall voids, or as a crack and crevice spray in and around residential and commercial structures, as well as on turf and landscaping. EVERCIDE® Esfenvalerate 6.4% CS may be sprayed on any surface that will not be stained by water. DO NOT use this product as a fogger or apply it as a space spray. Do not apply in electrical conduits, junction and switch boxes, motor housings, or other electrical equipment due to shock hazard from water-based spray.

Use only in areas described. Remove pets and cover fish aquariums and pets' food and water dishes before spraying. Keep all people (especially children) and pets out of areas being treated and restrict access to these areas until all surfaces are dry. DO NOT leave product where children or pets can come into contact with it. DO NOT allow spray to contact food or food-containing surfaces, feed, or water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated with this product. Food utensils such as teaspoons and measuring cups must not be used for food purposes after use with pesticides.

APPLICATION IN FOOD PROCESSING/HANDLING ESTABLISHMENTS:

Application is Permitted Within Food and Non-Food Areas of Food Service and Handling Establishments (Places Other than Private Residences) Including But Not Limited to: Restaurants, Grocery Stores, Bakeries, Food Manufacturing and Processing Establishments,

and Food Warehousing Establishments. Do not treat establishments where livestock feed is present.

Food areas include areas for receiving, storing, packing, canning, bottling, wrapping, boxing, preparing, edible waste storage, and enclosed processing systems, mills, and dairies. Serving areas are places where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held. Non-food areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, garages, mop closets, and storage after canning or bottling.

General Surface Applications: Do not apply EVERCIDE® Esfenvalerate 6.4% CS as a general surface application in food/feed areas of the facility when the facility is in operation or food/feed is exposed. Do not apply directly to food. Cover or remove all food and food processing equipment prior to application. After spraying in meat packing plants, bakeries, and other food processing plants, wash with an effective cleaning compound, and then rinse water all equipment, benches, shelving, etc. where exposed food will be handled with potable water. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

Spot or Crack & Crevice Applications: Spot or crack and crevice applications may be made while facility is in operation, provided exposed food is covered or removed from the area being treated. Do not apply directly to food or food-handling surfaces. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

In the home, cover exposed food and do not allow spray to contact food/feed surfaces. If spray does contact these surfaces, clean surfaces with soap and water.

MIXING INSTRUCTIONS:

EVERCIDE® Esfenvalerate 6.4% CS should be mixed with water and applied with hand pressurized or power operated sprayers. Shake container of EVERCIDE® Esfenvalerate 6.4% CS before diluting. Clean spray equipment before use. For dilution, add approximately half the required water to spray tank and then add the appropriate amount of EVERCIDE® Esfenvalerate 6.4% CS. Agitate, and slowly add the remaining water. Agitate spray thoroughly before using and also occasionally during use to ensure dispersion. If spray filter screens are used, they should be 50 mesh or larger. Use 0.5 fluid ounces (15 cc.) of EVERCIDE® Esfenvalerate 6.4% CS in 1 gallon of water to make a spray mix containing 0.025% active ingredient. Use 1.0 fluid ounce (30 cc.) per gallon of water to make a spray containing 0.05% active ingredient. Use 0.025% solution for light infestations or as a maintenance control rate. Use 0.05% for heavy infestations or as an initial clean out rate. Apply two-second bursts of spray per square foot of area being treated. Avoid excessive application. Dampen surfaces but not to the point of saturation or run-off. Only dilute EVERCIDE® Esfenvalerate 6.4% CS with water.

Formula for determining the active ingredient content of the finished spray mixture: The following formula may be used to determine the percent active ingredient that is in the sprayer tank after mixing EVERCIDE® Esfenvalerate 6.4% CS:

$$\frac{(6.4) \times (\text{number of fl. oz. of EVERCIDE® Esfenvalerate 6.4\% CS added to tank})}{(\text{Gallons of finished spray mix}) \times (128)} = \text{Percent Active Ingredient of Spray Mix}$$

Tank Mixing:

EVERCIDE® Esfenvalerate 6.4% CS may be tank mixed with an insect growth regulator such as NyGuard® IGR Concentrate or pyrethrum-containing products or any other currently registered pesticides unless expressly prohibited by the product label. The resulting tank mix may be applied in areas where these products are allowed to be sprayed. Do not tank mix EVERCIDE® Esfenvalerate 6.4% CS with products containing dichlorvos (DDVP).

INDOOR USE:

Application Rates for Indoor Structural Pests:

Pests	Concentration of Active Ingredient	Dilution Rate In Water	Application Rate of Mixed Solution	Application Method(s)
Crawling Insects	0.025%	0.5 fl. oz. / gallon	1 gallon / 1,000 sq. ft	Apply as a course, wet spray * Broadcast Surface Spray * Crack & Crevice Treatment * Spot Treatment * Injected into Wall Voids
Light Infestation	0.050%	1.0 fl. oz. / gallon		
Heavy Infestation				
Flying Insects	0.025%	0.5 fl. oz. / gallon	1 gallon / 1,000 sq. ft	Apply as noted above, with a fine particle spray.
Stinging Insects	0.050%	1.0 fl. oz. / gallon	1 gallon / 1,000 sq. ft	
Ticks	0.025%	0.5 fl. oz. / gallon	1 gallon / 1,000 sq. ft.	
Fleas, Lice & Bedbugs	0.025%	0.5 fl. oz. / gallon	1 gallon / 1,000 sq. ft.	

Crawling Insect Pests Indoors:

Apply as a course, wet spray to surfaces where these pests are normally found. Treat floors, baseboards, around doors and windows, in attics, crawl spaces, eaves, corners, closets, walls, utility pipes, storage areas, and all cracks and crevices. Treat underneath sinks, dishwashers, refrigerators, stoves, the underside of shelves, drawers, cabinets, areas behind pipes, and in all places where these insects shelter. Contact as many insects as possible with direct spray. Repeat applications as needed, but do not exceed more than one (1) application every seven (7) days.

For Ant Control Indoors:

Apply to ant trails, around garbage receptacles, and near food sources. Also apply around doors and windows and wherever these pests may find entrance to the structure. Remove sources of food through basic hygiene practices whenever possible.

Flying Insect Pests Indoors:

Treat insect resting areas such as walls and ceilings, screens, around windows, doors, and light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season, and repeat treatments during periods of heavy infestation.

4/7

Stinging Insect Pests Indoors:

Apply spray to nests late in the evening when stinging insects are at rest. Thoroughly spray nest, nest entrance, and surrounding areas where insects land or walk. Treat around doors and windows, in attics, crawl spaces, and possible harborage sites or points of entry.

For Control of Brown Dog Ticks:

Thoroughly apply as a spot treatment to infested areas such as pet beds and resting areas, nearby cracks and crevices, along and behind baseboards, window and door frames, and localized areas of floor and floor coverings where these pests may be present. **DO NOT SPRAY PETS WITH THIS PRODUCT.** Treat dogs and cats with a product registered for use on animals

For Control of Fleas:

Thoroughly apply as a fine particle broadcast spray to infested rugs, carpets, and pet resting areas. Prior to treatment, aquariums and fish bowls should be covered, and pet animals should be removed from the area being treated. Do not permit humans or pets to contact treated surfaces until spray has dried. Old pet bedding should be removed and replaced with clean, fresh bedding after treatment. **DO NOT SPRAY PETS WITH THIS PRODUCT.** Treat dogs and cats with a product registered for use on animals.

To Control Bedbugs:

Thoroughly clean and sanitize mattresses and box springs. Treat mattresses and box springs with an approved pesticide, such as a pyrethrin aerosol. Apply EVERCIDE® Esfenvalerate 6.4% CS as a spot treatment to potential harborage sites and migration paths, and cracks and crevices, around baseboards, floorboards, headboards, and walls.

For Control of Stored Product Pests:

Spray thoroughly around and into floor drains, non-food conveyors, benches, pipes, pallets, moist areas, storage racks, pieces of equipment, and other areas where stored product pests may be found. Tank-mix or sequential use of an insect growth regulator, such as NyGuard® IGR Concentrate, is recommended to break the insect reproduction cycle. Do not apply this spray to surfaces or utensils that may come in contact with food, since excessive residues in food may result.

To Treat Voids in Equipment and Structures:

To kill insects harboring in wall voids and other inaccessible spaces in equipment and structures, use injection equipment designed for deep-void applications. Follow the injection equipment manufacturer's recommendations for proper set up and air pressures. Place the applicator tip at or into the void space to be treated. For inaccessible voids, it may be necessary to drill an access hole(s). Inject product into the void space in short bursts, allowing air pressure to push insecticide deep into the space. Avoid applying to the point of runoff or drip. Ventilate area thoroughly before re-entry. Do not reapply more than every fourteen (14) days.

USE IN AND AROUND LIVESTOCK HOUSING:

For use in and around unoccupied areas of livestock facilities, such as, but not limited to: Barns, Cow and Calf Pens & Hutches, Dairy Barns & Milk Rooms, Hog Barns, Horse Barns, Sheep Barns, Poultry Houses, and Rabbit Hutches. Do not contaminate milk, food, or drinking water. Remove animals from area being treated. Cover feeders and waterers. Do not apply or allow insecticide to drift onto animals. Do not allow animals to enter treated areas until surfaces are dry.

Pests	Concentration of Active Ingredient	Dilution Rate In Water	Application Method(s)
Crawling Insects * Including litter beetles	0.050%	1 fl. oz. / gallon or 1 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a course, wet spray * General Surface
Flying Insects	0.025%	0.5 fl. oz. / gallon or 0.5 fl. oz. in sufficient water to cover 1,000 sq. ft.	Spray * Crack & Crevice Treatment * Spot Treatment * Injected into Wall Voids

Crawling Insect Pests in Livestock Premises:

Apply as a general surface, spot treatment, and/or crack and crevice treatment. Apply to floors and vertical and overhead surfaces where crawling insects are or may be present. Treat stanchions, pipes, windows, doors, posts, cage framing, gates, under (but not in) feeders, and other areas where insects hide or congregate. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1 – 3 feet up the exterior foundation wall and 3 – 6 feet out from the foundation.

Litter Beetle Control in Livestock Premises:

To control litter beetles (darkling, hide, and carrion beetles) in poultry housing, apply EVERCIDE® Esfenvalerate 6.4% CS to walls and floors at cleanout and before reintroduction of animals. Treat areas where beetles frequently occur, such as walls, supports, cages, cage framing, stalls, and around feeders. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1 – 3 feet up the exterior foundation wall and 3 – 6 feet out from the foundation.

Flying Insect Pests in Livestock Premises:

For residual control of flying insects, treat insect resting areas such walls, ceilings, screens, around windows, doors, light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season and repeat treatments during periods of heavy infestation.

OUTDOOR USE:

Do not spray in or near fishponds or other bodies of water.

Not for use on plants being grown for sale or other commercial use. Not for use in nurseries, sod farms or golf courses.

517

Application Rates for Outdoor Pests:

	Concentration of Active Ingredient	Dilution Rate In Water	Application Method(s)
Treating Exterior Walls, Foundations and Structures	0.025 to 0.050%	0.5 to 1 fl. oz. / gallon or 0.5 to 1 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a course, wet spray * General
Treating nests and harborage areas of stinging insects	0.050%	1 fl. oz. / gallon or 1 fl. oz. in sufficient water to cover 1,000 sq. ft.	Surface Spray * Crack & Crevice
To Control Swarming termites and Wood Destroying Pests.	0.050%	1 fl. oz. / gallon or 1 fl. oz. in sufficient water to cover 1,000 sq. ft.	Treatment * Spot Treatment
Treating Ornamental Trees and Landscapes	0.005 to 0.050%	0.1 to 1.0 fl. oz. / gallon or 0.1 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a course, wet spray * Broadcast
Mosquito Breeding Sites	0.025%	0.5 fl. oz. / gallon or 0.5 fl. oz. in sufficient water to cover 1,000 sq. ft.	Spray Treatment * Foliar Spray Treatment
Treating Lawns and Turf	0.005 to 0.050%	0.1 to 1 fl. oz. / gallon or 0.1 to 1 fl. oz. in sufficient water to cover 1,000 sq. ft. 4.3 to 43 fl. oz. per acre	Apply as a course, wet spray * Broadcast Spray Treatment

Treating exterior walls, foundations and structures:

To control infestations of listed pests, treat exterior surfaces of buildings, walls, window frames, around garbage cans, eaves, cracks and crevices, porches, decks, gazebos, patios, carports, garages, fence lines, and other areas where pests are active or may be hiding.

To help prevent insect infestation of buildings: Treat a 2 – 6 foot band of soil or other substrate adjacent to buildings. Treat building foundation to a height of 2 – 3 feet. Treat exterior walls, eaves, cracks and crevices, and other areas where listed pests are active, and may find entrance into building.

Stinging Insect Control Outdoors:

For stinging insects, apply spray solution to nests and harborage areas late in the evening when stinging insects are commonly at rest. Thoroughly spray nest, nest entrance, harborage sites, and surrounding areas where insects land or walk. For nests inside walls, inject sufficient spray to treat nest. Do not use in areas where an electrical shock hazard exists. For applications made when pests are active, applicator should wear protective equipment as required. Repeat application when there are signs of renewed insect activity.

For Yellow Jacket control, EVERCIDE® Esfenvalerate 6.4% CS can be mixed with baits in traps. Follow trap instructions for preparation of bait.

For control of Fire Ants, combine broadcast application with mound drenches to control foraging workers and newly mated fly-in queens. Apply EVERCIDE® Esfenvalerate 6.4% CS as a broadcast application at a rate of 8 fl. oz. per 1,000 square feet. Treat mounds with a registered fire ant mound treatment.

Insect Control in Ornamental Trees and Landscapes:

For residential and commercial trees, shrubs, ground covers, and bedding and foliage plants that will not be harvested for food, apply EVERCIDE® Esfenvalerate 6.4% CS in appropriate volumes of water to obtain thorough coverage. Apply as a full-coverage foliar spray, applying to the point of drip but not runoff. Treat active soil pests with an application to surrounding soil. Repeat treatments as necessary to achieve control, using higher application rates as pest pressure and foliage area increases. Repeat applications as necessary, but no more than once every seven (7) days. Certain plants may be sensitive to the final spray solution. A small-scale test is recommended to verify safety to ornamental plants. Spray and observe for one (1) week prior to application of an entire planting.

Insect Control on Lawns, Turf Grass, and Turf:

For best results, lawn or turf grass should be mowed 1 – 2 days before spraying. Treat with spraying equipment or a hand sprayer. Use application volumes of up to 10 gallons per 1,000 square feet to get uniform coverage when treating dense grass foliage. For low volume applications using less than 2 gallons of spray solution per 1,000 square feet, immediate irrigation with at least 0.25 inches of water is recommended to improve effectiveness on sub-surface pests.

When hand spraying, spray using a slow, even sweeping motion, making sure to cover the entire lawn or turf grass area where pests are observed. Spray under ornamentals and trees. Repeat treatments may be necessary at 7 – 14-day intervals.

Rate Range:

Use lower rate range for pests that are commonly exposed and will be contacted by spray solution at the time of application. For pests that will not be contacted by spray solution at the time of application, use the upper rate range. Use 1 fl. oz per 1,000 sq. ft. when treating Mole Crickets and Chinch Bugs.

To Kill Swarming Termites: OUTDOORS ONLY.

Apply spray mix as a coarse, wetting spray when swarming termites are seen emerging from woodpiles, wooden fence posts, wooden structures, or from the ground. Swarming usually occurs in the spring or at other times when a termite colony becomes overcrowded and new reproductive termites with wings emerge and fly away to mate and establish new colonies. This treatment will control the sprayed termites and will not protect the structure from which the swarm is coming. Use only as a contact spray to kill emerging reproductive (winged) and worker termites emerging from infested wood. This treatment is not a substitute for a comprehensive termite control program.

6/7

To Kill Wood Destroying Pests: OUTDOORS ONLY.

Apply spray mix as a coarse, wetting spray to exposed pests and to the damaged areas of wood, spraying into galleries or tunnels in the exposed wood. Also, spray around doors, window and door frames, and other areas where these pests may hide or enter the house. Spray into cracks and crevices and, if necessary, drill small holes and spray into inaccessible wooden structural voids where these pests are suspected.

Mosquito Breeding sites:

Mosquito populations may be reduced by application of EVERCIDE® Esfenvalerate 6.4% CS to sites where mosquitoes rest, harbor, and breed. Apply spray solution into tall grass, shrubbery, and around backyards and lawns where these pests may hover or rest. Apply while air is still.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry, well-ventilated area. Keep container closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. In case of spill or leak, soak up with sand, earth, or synthetic absorbents. Do not use alkaline absorbents or clean spill area with alkaline detergents.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container in several layers of newspaper and discard in trash.