

1021-1815

03106/2009

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms Miriam Frugis
McLaughlin Gormley King Company
8810 10th Avenue North
Minneapolis, MN 55427

MAR 6 2009

Subject: Label Notification(s) for Pesticide Registration Notices 2007-4 and 98-10
Storage and Disposal language Revision and Other Changes

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 and 98-10 dated January 30, 2009 for:

EPA Registration 1021-1815 Evercide®Esfenvalerate 6.4% CS

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and 98-10 and finds that the label change(s) requested falls within the scope of PRN-2007-4 and 98-10. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Banza Djapao of my staff at 703-305-7269.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

3/11



January 30, 2009

Document Processing Desk (NOTIF)
U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs (H7504P)
Room S-4900, One Potomac Yard
2777 South Crystal Dr.
Arlington, VA 22202-4501

Via Courier

Subject: EVERCIDE® Esfenvalerate 6.4% CS
(Microencapsulated Insecticide)
EPA Reg. No. 1021-1815
Notifications Per PR-Notice 2007-4 and PR-Notice 98-10

Dear Sir or Madam:

By Notification we are changing the storage and disposal language to comply with the Final Rule "Pesticide Management and Disposal; Standards for Pesticide Containers and Containment". We are also adding metric units to the Directions for Use, without changes in application rates.

Enclosed are 2 copies of the label, one copy has changes highlighted.

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.


This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

If there are any questions or you need anything further, please contact this office.

Sincerely,

McLAUGHLIN GORMLEY KING COMPANY


Miriam Frugis
Registration Specialist



Enclosure

Quality Products Since 1902



McLAUGHLIN
GORMLEY
KING COMPANY

EVERCIDE® Esfenvalerate 6.4% CS
(Microencapsulated Insecticide)

For Pest Management Professional Use

8810 Tenth Avenue N. / Minneapolis, Minnesota 55427-4319 U.S.A. - Telephone (763) 544-0341

Kills for up to 12 months (1 year)*

*For German Cockroaches on indoor non-porous surfaces only

NOTIFICATION

MAR 6 2009

ACTIVE INGREDIENT:

(S)-cyano(3-phenoxyphenyl)methyl-(S)-4-chloro-alpha-(1-methylethyl)benzeneacetate.....	6.40%
*OTHER INGREDIENTS.....	93.60%
*Contains petroleum distillates	100.00%

EVERCIDE® - Registered trademark of McLaughlin Gormley King Company

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Do not give any liquid to the person. ▪ Do not induce vomiting unless told to do so by a poison control center or a doctor. ▪ Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. ▪ Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15-20 minutes. ▪ Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN: Contains petroleum distillate – vomiting may cause aspiration pneumonia.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	

Net Contents _____

Manufactured by:
Mc LAUGHLIN GORMLEY KING COMPANY
8810 Tenth Avenue North
Minneapolis, MN 55427

EPA Reg. No. 1021-1815

EPA Est. No. 1021-MN-2

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear protective eyewear (goggles, face shield or safety glasses with side shields). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and other aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirement of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product
in a manner inconsistent with its labeling.

Remove pets and birds and cover fish aquariums before spraying.

For control of insects, indoors and outdoors, in food and non-food areas such as, but not limited to: Homes, Schools, Warehouses, Office Buildings, Apartment Buildings, Theatres, Hotels, Industrial Buildings, Motels, Kennels, Livestock Housing, Food Processing Plants, Food Service Establishments, Restaurants, Supermarkets and Grocery Stores, Transportation Equipment, Truck Trailers, Railroad Cars, and Food Manufacturing and Warehousing Establishments. Also for use on Backyards, Lawns, Trees, Ornamental Landscaping, Recreational Areas, Parks and Athletic Fields.

Insects Controlled:

Ants	Face Flies	Skippers	Cowpea Curculios
Aphids	Fruit Flies	Spider Beetles	Cranberry Fruitworms
Asian Lady Beetles	Fungus Gnats	Spider Mites	Cranberry Weevils
Bedbugs	Gnats	Square-necked Grain Beetles	Cucumber Beetles
Beetles	Horn Flies	Tobacco Moths	Cutworms
Boxelder Bugs	Houseflies	Yellow Mealworms	Diamondback Moths
Carpet Beetles	Mosquitoes		Elm Leaf Beetles
Centipedes	Small Flying Moths	Wood-Destroying Pests including:	European Pine Sawflies
Cockroaches	Whiteflies	Carpenter Ants	Fall Webworms
Crickets		Carpenter Bees	Filbert Worms
Dog Ticks	Stored Product Pests including:	Deathwatch Beetles	Flea Beetles
Earwigs	Angoumois Grain Moths	Furniture Beetles	Grasshoppers
Fire Ants	Ants	Old House Borers	Green Cloverworms
Firebrats	Cadelles	Powder Post Beetles	Green Fruitworms
Fleas	Cigarette Beetles	Round-headed House Borers	Gypsy Moth Caterpillars
Mealy Bugs	Confused Flour Beetles	Swarming Termites	Hickory Shuckworms
Midges	Dark Mealworms		Imported Cabbageworms
Millipedes	Dried Fruit Beetles	Plant Pests including:	Japanese Beetles
Mites	Drugstore Beetles	Alfalfa Caterpillars	Lace Bugs
Pillbugs	Flat Grain Beetles	Artichoke Plum Moth	Leaf Feeding Caterpillars
Psocids	Fruit Flies	American Plum Borers	Leaf Miners
Scales	Grain Mites	Annual Bluegrass Weevils	Leaf Rollers
Silverfish	Grain Moths	Aphids	Leaf Tiers
Spiders	Granary Weevils	Apple Maggots	Leafhoppers
Springtails	Indian Meal Moths	Armyworms	Lesser Appleworms
Swarming Termites	Lesser Grain Borers	Bagworms	Lesser Peach Tree Borers
Ticks	Maize Weevils	Balsam Woolly Adelgids	Loopers
Waterbugs	Meal Moth Larvae	Beet Armyworms	Lygus Bugs
	Mediterranean Flour Moths	Billbugs	Mites
Flying Insects including:	Merchant Grain Beetles	Blueberry Spanworms	Mexican Bean Beetles
Blow Flies	Red Flour Beetles	Cherry Fruit Flies	Mole Crickets
Chiggers	Rice Weevils	Cherry Fruitworms	Naval Orangeworms
Clothes Moths	Rusty Grain Beetles	Chinch Bugs	Northern Pine Weevils
Crane Flies	Saw-toothed Grain Beetles	Codling Moths	Oriental Fruit Moths
Deer Flies			
Painted Lady Caterpillars	Red-striped Fireworms	Biting and Stinging Pests:	Livestock Premise Pests:
Pea Weevils	Red-headed Pine Sawflies	Bedbugs	Bedbugs
Peach Tree Borers	Rindworms	Bees	Carrion Beetles
Peach Twig Borers	Salt Marsh Caterpillars	Biting Flies	Chiggers
Pear Psyllid	Sap Beetles	Chiggers	Darkling Beetles (lesser mealworm)
Pear Slugs	Scales	Deer Flies	Deer Flies

6/11

Pecan Leaf Phylloxera	Sod Webworms	Deer Ticks	Face Flies
Pecan Nut Casebearers	Sowbugs	Dog Ticks	Fleas
Pecan Spittlebugs	Spiders	Fire Ants	Flies
Pecan Stem Phylloxera	Spittlebugs	Fleas	Hide Beetles
Pecan Weevils	Stink Bugs	Gnats	Horn Flies
Pepper Weevils	Tarnished Plant Bugs	Hornets	Horse Flies
Periodical Cicadas	Tent Caterpillars	Lice	Lice
Pickleworms	Thrips	Mosquitoes	Litter Beetles
Pillbugs	Tobacco Hornworms	Scorpions	Mites
Pine Chagers	Tufted Apple Bud Moths	Spiders	Mosquitoes
Pine Coreid Bugs	Velvet Bean Caterpillars	Ticks	Stable Flies
Plant Bugs	Walnut Husk Flies	Wasps	
Plum Curculios	Western Bean Cutworms	Yellow Jackets	
Red Pine Sawflies	Whiteflies		

GENERAL INFORMATION:

Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm. Do not apply to classrooms while in use. Do not apply in institutions (including libraries, schools, sports facilities, etc.) in the immediate area where occupants are present.

EVERCIDE® Esfenvalerate 6.4% CS is a microencapsulated suspension concentrate containing 0.54 pounds active ingredient per gallon (64 mg of a.i./L).

EVERCIDE® Esfenvalerate 6.4% CS is designed to provide long residual activity against pests listed on this label when applied as surface or spot treatments, injected into wall voids, or as a crack and crevice spray in and around residential and commercial structures, as well as on turf and landscaping. EVERCIDE® Esfenvalerate 6.4% CS may be sprayed on any surface that will not be stained by water. DO NOT use this product as a fogger or apply it as a space spray. Do not apply in electrical conduits, junction and switch boxes, motor housings, or other electrical equipment due to shock hazard from water-based spray.

Use only in areas described. Remove pets and cover fish aquariums and pets' food and water dishes before spraying. Keep all people (especially children) and pets out of areas being treated and restrict access to these areas until all surfaces are dry. DO NOT leave product where children or pets can come into contact with it. DO NOT allow spray to contact food or food-containing surfaces, feed, or water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated with this product. Food utensils such as teaspoons and measuring cups must not be used for food purposes after use with pesticides.

APPLICATION IN FOOD PROCESSING/HANDLING ESTABLISHMENTS:

Application is Permitted Within Food and Non-Food Areas of Food Service and Handling Establishments (Places Other than Private Residences) Including But Not Limited to: Restaurants, Meat Processing Plants, Grocery Stores, Bakeries, Food Manufacturing and Processing Establishments, and Food Warehousing Establishments. Do not treat establishments where livestock feed is present.

Food areas include areas for receiving, storing, packing, canning, bottling, wrapping, boxing, preparing, edible waste storage, and enclosed processing systems, mills, and dairies. Serving areas are places where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held. Non-food areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, garages, mop closets, and storage after canning or bottling.

General Surface Applications: Do not apply EVERCIDE® Esfenvalerate 6.4% CS as a general surface application in food areas of the facility when the facility is in operation or food is exposed. Do not apply directly to food. Cover or remove all food and food processing equipment prior to application. After spraying in meat packing plants, bakeries, and other food processing plants, wash with an effective cleaning compound, and then rinse all equipment, benches, shelving, etc. where exposed food will be handled with potable water. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

Spot or Crack & Crevice Applications: Spot or crack and crevice applications may be made while facility is in operation, provided exposed food is covered or removed from the area being treated. Do not apply directly to food or food-handling surfaces. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

In the home, cover exposed food and do not allow spray to contact food surfaces. If spray does contact these surfaces, clean surfaces with soap and water.

MIXING INSTRUCTIONS:

EVERCIDE® Esfenvalerate 6.4% CS should be mixed with water and applied with hand pressurized or power operated sprayers. Shake container of EVERCIDE® Esfenvalerate 6.4% CS before diluting. Clean spray equipment before use. For dilution, add approximately half the required water to spray tank and then add the appropriate amount of EVERCIDE® Esfenvalerate 6.4% CS. Agitate, and slowly add the remaining water. Agitate spray thoroughly before using and also occasionally during use to ensure dispersion. If spray filter screens are used, they should be 50 mesh or larger. Use 0.5 fluid ounces (15 cc.) of EVERCIDE® Esfenvalerate 6.4% CS in 1 gallon of water to make a spray mix containing 0.025% active ingredient. Use 1.0 fluid ounce (30 cc.) per gallon of water to make a spray containing 0.05% active ingredient. Use 0.025% solution for light infestations or as a maintenance control rate. Use 0.05% for heavy infestations or as an initial clean out rate. Apply two-second bursts of spray per square foot of area being treated. Avoid excessive application. Dampen surfaces but not to the point of saturation or run-off. Only dilute EVERCIDE® Esfenvalerate 6.4% CS with water.

Formula for determining the active ingredient content of the finished spray mixture: The following formula may be used to determine the percent active ingredient that is in the sprayer tank after mixing EVERCIDE® Esfenvalerate 6.4% CS:

$$\frac{(6.4) \times (\text{number of fl. oz. of EVERCIDE® Esfenvalerate 6.4\% CS added to tank})}{(\text{Gallons of finished spray mix}) \times (128)} = \text{Percent Active Ingredient of Spray Mix}$$

Tank Mixing:

EVERCIDE® Esfenvalerate 6.4% CS may be tank mixed with an insect growth regulator such as NyGuard® IGR Concentrate or pyrethrum-containing products or any other currently registered pesticides unless expressly prohibited by the product label. The resulting tank mix may be applied in areas where these products are allowed to be sprayed. Do not tank mix EVERCIDE® Esfenvalerate 6.4% CS with products containing dichlorvos (DDVP).

INDOOR USE:

Application Rates for Indoor Structural Pests:

Pests	Concentration of Active Ingredient	Dilution Rate In Water	Application Rate of Mixed Solution	Application Method(s)
Crawling Insects Light Infestation	0.025%	0.5 fl. oz. / gallon (3.97cc/L)	1 gallon / 1,000 sq. ft (40 ml/m ²)	Apply as a coarse, wet spray * Broadcast Surface Spray * Crack & Crevice Treatment
Heavy Infestation	0.050%	1.0 fl. oz. / gallon (7.93 cc/L)		
Flying Insects	0.025%	0.5 fl. oz. / gallon (3.97cc/L)	1 gallon / 1,000 sq. ft (40 ml/m ²)	* Spot Treatment * Injected into Wall Voids
Stinging Insects	0.050%	1.0 fl. oz. / gallon (7.93 cc/L)	1 gallon / 1,000 sq. ft (40 ml/m ²)	
Ticks	0.025%	0.5 fl. oz. / gallon (3.97cc/L)	1 gallon / 1,000 sq. ft. (40 ml/m ²)	
Fleas, Lice & Bedbugs	0.025%	0.5 fl. oz. / gallon (3.97cc/L)	1 gallon / 1,000 sq. ft. (40 ml/m ²)	Apply as noted above, with a fine particle spray.

Crawling Insect Pests Indoors:

Apply as a coarse, wet spray to surfaces where these pests are normally found. Treat floors, baseboards, around doors and windows, in attics, crawl spaces, eaves, corners, closets, walls, utility pipes, storage areas, and all cracks and crevices. Treat underneath sinks, dishwashers, refrigerators, stoves, the underside of shelves, drawers, cabinets, areas behind pipes, and in all places where these insects shelter. Contact as many insects as possible with direct spray. Repeat applications as needed, but do not exceed more than one (1) application every seven (7) days.

For Ant Control Indoors:

Apply to ant trails, around garbage receptacles, and near food sources. Also apply around doors and windows and wherever these pests may find entrance to the structure. Remove sources of food through basic hygiene practices whenever possible.

Flying Insect Pests Indoors:

Treat insect resting areas such as walls and ceilings, screens, around windows, doors, and light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season, and repeat treatments during periods of heavy infestation.

Stinging Insect Pests Indoors:

Apply spray to nests late in the evening when stinging insects are at rest. Thoroughly spray nest, nest entrance, and surrounding areas where insects land or walk. Treat around doors and windows, in attics, crawl spaces, and possible harborage sites or points of entry.

For Control of Brown Dog Ticks:

Thoroughly apply as a spot treatment to infested areas such as pet beds and resting areas, nearby cracks and crevices, along and behind baseboards, window and door frames, and localized areas of floor and floor coverings where these pests may be present. **DO NOT SPRAY PETS WITH THIS PRODUCT.** Treat dogs and cats with a product registered for use on animals

For Control of Fleas:

Thoroughly apply as a fine particle broadcast spray to infested rugs, carpets, and pet resting areas. Prior to treatment, aquariums and fish bowls should be covered, and pet animals should be removed from the area being treated. Do not permit humans or pets to contact treated surfaces until spray has dried. Old pet bedding should be removed and replaced with clean, fresh bedding after treatment. **DO NOT SPRAY PETS WITH THIS PRODUCT.** Treat dogs and cats with a product registered for use on animals.

To Control Bedbugs:

Thoroughly clean and sanitize mattresses and box springs. Treat mattresses and box springs with an approved pesticide, such as a pyrethrin aerosol. Apply EVERCIDE® Esfenvalerate 6.4% CS as a spot treatment to potential harborage sites and migration paths, and cracks and crevices, around baseboards, floorboards, headboards, and walls.

For Control of Stored Product Pests:

Spray thoroughly around and into floor drains, non-food conveyors, benches, pipes, pallets, moist areas, storage racks, pieces of equipment, and other areas where stored product pests may be found. Tank-mix or sequential use of an insect growth regulator, such as NyGuard® IGR Concentrate, is recommended to break the insect reproduction cycle. Do not apply this spray to surfaces or utensils that may come in contact with food, since excessive residues in food may result.

To Treat Voids in Equipment and Structures:

To kill insects harboring in wall voids and other inaccessible spaces in equipment and structures, use injection equipment designed for deep void applications. Follow the injection equipment manufacturer's recommendations for proper set up and air pressures. Place the applicator tip at or into the void space to be treated. For inaccessible voids, it may be necessary to drill an access hole(s). Inject product into the void space in short bursts, allowing air pressure to push insecticide deep into the space. Avoid applying to the point of runoff or drip. Ventilate area thoroughly before re-entry. Do not reapply more than every fourteen (14) days.

To Kill the Accessible Stages of Listed Granary Insects: Pest Management Professionals and grain producers may use this concentrate to treat grain storage facilities, and other listed areas, for stored product pest control. For control of exposed adult and immature stages of stored product pests, apply to cracks, crevices, and other surfaces where the pests have been seen or have harborage. Treat areas where products are stored before filling with the product. Apply at the rates listed above, using one gallon of spray mix per 1000 square feet of surface area to be treated. Cleaning all areas prior to use will increase levels of control. Any foodstuffs infested with pests should be removed and destroyed. Do not apply when food-processing facility is in operation or foods are exposed. Do not apply this spray to surfaces or utensils that may come into contact with food.

Repeat Application as Needed, but Do Not Exceed More Than One Application Every Fourteen (14) Days.

USE IN AND AROUND LIVESTOCK HOUSING:

For use in and around unoccupied areas of livestock facilities, such as, but not limited to: Barns, Cow and Calf Pens & Hutches, Dairy Barns & Milk Rooms, Hog Barns, Horse Barns, Sheep Barns, Poultry Houses, and Rabbit Hutches. Do not contaminate milk, food, or drinking water. Remove animals from area being treated. Cover feeders and waterers. Do not apply or allow insecticide to drift onto animals. Do not allow animals to enter treated areas until spray solution has dried.

9/11

Pests	Concentration of Active Ingredient	Dilution Rate In Water	Application Method(s)
Crawling Insects * Including litter beetles	0.050%	1 fl. oz. / gallon (7.93 cc/L) or 1 fl. oz. in sufficient water to cover 1,000 sq. ft. (0.32 cc/m ²)	Apply as a coarse, wet spray * General Surface
Flying Insects	0.025%	0.5 fl. oz. / gallon (3.97 cc/L) or 0.5 fl. oz. in sufficient water to cover 1,000 sq. ft. (0.16 cc/m ²)	Spray * Crack & Crevice Treatment * Spot Treatment * Injected into Wall Voids

Crawling Insect Pests in Livestock and Poultry Premises:

Apply as a general surface, spot treatment, and/or crack and crevice treatment. Apply to floors and vertical and overhead surfaces where crawling insects are or may be present. Treat stanchions, pipes, windows, doors, posts, cage framing, gates, under (but not in) feeders, and other areas where insects hide or congregate. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1 – 3 feet (0.3 -0.9 m) up the exterior foundation wall and 3 – 6 (0.3 -1.8 m) feet out from the foundation.

Litter Beetle Control in Livestock and Poultry Premises:

To control litter beetles (darkling, hide, and carrion beetles), apply EVERCIDE® Esfenvalerate 6.4% CS to walls and floors at cleanout and before reintroduction of animals. Treat areas where beetles frequently occur, such as walls, supports, cages, cage framing, stalls, and around feeders. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1 – 3 feet (0.3 – 0.9 m) up the exterior foundation wall and 3 – 6 feet (0.3 – 1.8 m) out from the foundation.

Flying Insect Pests in Livestock and Poultry Premises:

For residual control of flying insects, treat insect resting areas such as walls, ceilings, screens, around windows, doors, light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season and repeat treatments during periods of heavy infestation.

OUTDOOR USE:

Do not spray in or near fishponds or other bodies of water.

Not for use on plants being grown for sale or other commercial use. Not for use in nurseries, sod farms or golf courses.

Application Rates for Outdoor Pests:

	Concentration of Active Ingredient	Dilution Rate In Water	Application Method(s)
Treating Exterior Walls, Foundations and Structures	0.025 to 0.050%	0.5 to 1 fl. oz. / gallon (3.97 cc to 7.93 cc/L) or 0.5 to 1 fl. oz. in sufficient water to cover 1,000 sq. ft. (0.16 cc to 0.32 cc/m ²)	Apply as a coarse, wet spray * General Surface
Treating nests and harborage areas of stinging insects	0.050%	1 fl. oz. / gallon (7.93 cc/L)or 1 fl. oz. in sufficient water to cover 1,000 sq. ft.(0.32 cc/m ²)	Spray * Crack & Crevice Treatment
To Control Swarming Termites and Wood Destroying Pests.	0.050%	1 fl. oz. / gallon (7.93 cc/L) or 1 fl. oz. in sufficient water to cover 1,000 sq. ft. .(0.32 cc/m ²)	* Spot Treatment
Treating Ornamental Trees and Landscapes	0.005 to 0.050%	0.1 to 1.0 fl. oz. / gallon (0.793 cc to 7.93 cc/L) or 0.1 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.(0.032cc to 0.32 cc/m ²)	Apply as a coarse, wet spray * Broadcast Spray Treatment

<p>Mosquito Breeding Sites</p>	<p>0.025%</p>	<p>0.5 fl. oz. / gallon (3.97 cc/L) or 0.5 fl. oz. in sufficient water to cover 1,000 sq. ft. (0.16 cc/m²)</p>	<p>* Foliar Spray Treatment</p>
<p>Treating Lawns and Turf</p>	<p>0.005 to 0.050%</p>	<p>0.1 to 1 fl. oz. / gallon (0.793 cc to 7.93 cc/L) or 0.1 to 1 fl. oz. in sufficient water to cover 1,000 sq. ft. (0.032cc to 0.32 cc/m²) 4.3 to 43 fl. oz. per acre (127.17 cc to 1,271.66 cc/acre)</p>	<p>Apply as a coarse, wet spray * Broadcast Spray Treatment</p>

Treating exterior walls, foundations and structures:

To control infestations of listed pests, treat exterior surfaces of buildings, walls, window frames, around garbage cans, eaves, cracks and crevices, porches, decks, gazebos, patios, carports, garages, fence lines, and other areas where pests are active or may be hiding.

To help prevent insect infestation of buildings: Treat a 2 – 6 foot (0.6 – 1.8 m) band of soil or other substrate adjacent to buildings. Treat building foundation to a height of 2 – 3 feet (0.6 – 0.9 m). Treat exterior walls, eaves, cracks and crevices, and other areas where listed pests are active, and may find entrance into building.

Apply with sufficient water to adequately cover the area being treated, but do not allow dripping or run-off to occur. Alternate mixing directions are to use 2 to 4 fluid ounces of concentrate per 50 gallons of water.

Stinging Insect Control Outdoors:

For stinging insects, apply spray solution to nests and harborage areas late in the evening when stinging insects are commonly at rest. Thoroughly spray nest, nest entrance, harborage sites, and surrounding areas where insects land or walk. For nests inside walls, inject sufficient spray to treat nest. Do not use in areas where an electrical shock hazard exists. For applications made when pests are active, applicator should wear protective equipment as required. Repeat application when there are signs of renewed insect activity.

For Yellow Jacket control, EVERCIDE® Esfenvalerate 6.4% CS can be mixed with baits in traps. Follow trap instructions for preparation of bait.

For control of Fire Ants, combine broadcast application with mound drenches to control foraging workers and newly mated fly-in queens. Apply EVERCIDE® Esfenvalerate 6.4% CS as a broadcast application at a rate of 8 fl. oz. of concentrate per 1,000 square feet (63.44 cc/L). Treat mounds with a registered fire ant mound treatment.

Insect Control in Ornamental Trees and Landscapes:

For residential and commercial trees, shrubs, ground covers, and bedding and foliage plants that will not be harvested for food, apply EVERCIDE® Esfenvalerate 6.4% CS in appropriate volumes of water to obtain thorough coverage. Apply as a full-coverage foliar spray, applying to the point of drip but not runoff. Treat active soil pests with an application to surrounding soil. Repeat treatments as necessary to achieve control, using higher application rates as pest pressure and foliage area increases. Repeat applications as necessary, but no more than once every seven (7) days. Certain plants may be sensitive to the final spray solution. A small-scale test is recommended to verify safety to ornamental plants. Spray and observe for one (1) week prior to application of an entire planting.

Insect Control on Lawns, Turf Grass, and Turf:

For best results, lawn or turf grass should be mowed 1 – 2 days before spraying. Treat with spraying equipment or a hand sprayer. Use application volumes of up to 10 gallons per 1,000 square feet (400 ml/m²) to get uniform coverage when treating dense grass foliage. For low volume applications using less than 2 gallons of spray solution per 1,000 square feet (80 ml/m²), immediate irrigation with at least 0.25 inches (6.35 cm) of water is recommended to improve effectiveness on sub-surface pests.

When hand spraying, spray using a slow, even sweeping motion, making sure to cover the entire lawn or turf grass area where pests are observed. Spray under ornamentals and trees. Repeat treatments may be necessary at 7 – 14-day intervals.

Rate Range:

Use lower rate range for pests that are commonly exposed and will be contacted by spray solution at the time of application. For pests that will not be contacted by spray solution at the time of application, use the upper rate range. Use 1 fl. oz per 1,000 sq. ft. (7.93 cc/L) when treating Mole Crickets and Chinch Bugs.

11/11

To Kill Swarming Termites: OUTDOORS ONLY.

Apply spray mix as a coarse, wetting spray when swarming termites are seen emerging from woodpiles, wooden fence posts, wooden structures, or from the ground. Swarming usually occurs in the spring or at other times when a termite colony becomes overcrowded and new reproductive termites with wings emerge and fly away to mate and establish new colonies. This treatment will control the sprayed termites and will not protect the structure from which the swarm is coming. Use only as a contact spray to kill emerging reproductive (winged) and worker termites emerging from infested wood. This treatment is not a substitute for a comprehensive termite control program.

To Kill Wood Destroying Pests: OUTDOORS ONLY.

Apply spray mix as a coarse, wetting spray to exposed pests and to the damaged areas of wood, spraying into galleries or tunnels in the exposed wood. Also, spray around doors, window and door frames, and other areas where these pests may hide or enter the house. Spray into cracks and crevices and, if necessary, drill small holes and spray into inaccessible wooden structural voids where these pests are suspected.

Mosquito Breeding sites:

Mosquito populations may be reduced by application of EVERCIDE® Esfenvalerate 6.4% CS to sites where mosquitoes rest, harbor, and breed. Apply spray solution into tall grass, shrubbery, and around backyards and lawns where these pests may hover or rest. Apply while air is still.

Not for wide area mosquito use. Do not apply with hand held or truck mounted cold aerosol ULV sprayers and thermal fogging devices.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry, well-ventilated area. Always store pesticides in the original container. Store away from food and pet food.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container.

If empty:

Place in trash or offer for recycling if available.

If partly filled:

Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTIFICATION

MAR 06 2009