



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

November 13, 2025

Thomas A. Lennan
Senior Regulatory Affairs Specialist
McLaughlin Gormley King Company D/B/A MGK
7325 Aspen Lane N
Minneapolis, MN 55428

Subject: Label Amendment - Registration Review Mitigation for Zeta-Cypermethrin
Product Name: EVERCIDE® 28511
EPA Registration Number: 1021-1859
Case Number: 476065
Application Dates: June 2, 2021

Dear Thomas A. Lennan:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Cypermethrins Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Tiffany Green by phone at 919-541-2446, or via email at green.tiffany@epa.gov.

Sincerely,

A handwritten signature in dark ink, appearing to read "Kevin Costello". The signature is fluid and cursive, with the first name "Kevin" and last name "Costello" clearly distinguishable.

Kevin Costello, Branch Chief
Risk Management and Implementation Branch 2
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

EVERCIDE® 28511

ACCEPTED

Nov 13, 2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 1021-1859

[] Denotes alternate language

For use outdoors and in homes and non-food areas of commercial buildings, institutions, warehouses, theaters, office buildings, schools, athletic fields, motels, hotels, restaurants, food/feed processing establishments, hospitals, kennels, farms, ranches

[One quart makes up to 25 gallons of spray]

[Makes up to 25 gallons]

[Controls [Kills] German Cockroaches for up to 5 weeks on non-porous surfaces]

[Kills German Cockroaches on Contact and Keeps Killing for up to 5 weeks on non-porous surfaces]

[Kills Carpenter Ants for up to 8 weeks on non-porous surfaces]

[Water Based Formula]

[Kills Bugs on Your Lawns, Trees and Shrubs, Roses and Flowers and Vegetables]

[Aids in Reducing Annoyance from Mosquitoes and Other Flying Insects]

[Kills Mosquitoes that transmit West Nile Virus]

[Kills Deer Ticks and other Crawling insects]

[May be used on indoor and outdoor plants] [Where to use: On lawns, listed fruits and vegetables, trees and shrubs, roses and flowers and outside surfaces of buildings]

[Controls over 50 common insect pests]

[Controls insects on: roses & ornamentals; trees, shrubs; lawns]

[Kills nuisance pests: ants, deer ticks, crickets & earwigs]

[16 oz. Treats up to 2,650 sq. ft.][Treats up to 2,650 sq. ft.]

[Protects horses, ponies, and other livestock from stable flies, mosquitoes, and gnats]

[Contains Zeta-Cypermethrin, a long lasting insecticide that provides contact and residual control of many hard-to-kill sucking and chewing insects]

ACTIVE INGREDIENT:

* Zeta-Cypermethrin [* (3-phenoxyphenyl)methyl (+) cis/trans 3-(2,2-dichloroethenyl)-2,2 dimethylcyclopropanecarboxylate].....

0.75%

OTHER INGREDIENTS 99.25%

100.00%

* Cis,trans ratio: Max 75% (±) cis and Min. 25% (±) trans

EVERCIDE® - Registered trademark of McLaughlin Gormley King Company

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none">Immediately call a poison control center or doctor.Do not induce vomiting unless told to do so by a poison control center or doctor.Do not give any liquid to the person.Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	

Manufactured by



7325 Aspen Lane North
Minneapolis, MN 55428

EPA Reg. No. 1021-1859

EPA Est. No. 1021-MN-2

Net Contents: _____

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not apply this product around electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product
in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USE. READ ENTIRE LABEL BEFORE EACH USE.

For both indoor and outdoor use

[Read Directions on Box for additional application instructions before using this product]

USE RESTRICTIONS:

- Not for use on plants being grown for sale or other commercial seed production or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, ornamental gardens or parks or lawns or grounds.
- Do not apply this product in patient rooms, or in any rooms while occupied by the elderly or infirm.
- Do not apply to classrooms when in use.
- Do not allow to enter indoor or outdoor drains No permita la entrada a desagües internos o externos.
-



{Note to reviewer: Pictogram must be no smaller than 1.5 cm² or 0.25 sq. in. unless size is greater than 10% of label size}

- All outdoor spray applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
 2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
 3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
 4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
 5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
 6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.
- Do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Do not apply directly to sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- Do not allow the product to enter any drain during or after application.
- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- Do not apply or irrigate to the point of runoff.
- Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application.
- Spot treatments must not exceed two square feet in size. (for example 2 ft by 1. Ft. or 4 ft. by 0.5 ft).
- Do not apply when the wind speed is greater than 15 mph.

THIS PRODUCT MUST BE DILUTED PRIOR TO USE. To prepare dilutions, add the required amount of concentrate to water and blend thoroughly.

Dilution rates on this label will yield the following ready-to-use spray mixes:

RATE	OZ. CONC./GAL	Zeta-Cypermethrin CONC.
1:5	25.6	0.150%
1:10	12.8	0.075%
4:100	5.12	0.030%
3:128	3.00	0.0176%
2:100	2.56	0.015%
2:128	2.00	0.012%
1.5:128	1.50	0.009%
1:100	1.28	0.0075%
1:128	1.00	0.006%

This product helps reduce annoying and troublesome outdoor insects. People and pets should not be allowed in treated areas until spray has dried.

This product kills most [troublesome] outdoor insects, which destroy lawns and gardens and are an annoyance on patios and at picnics.

This product can be applied using a pump up pressure sprayer or by watering can. Carefully dilute the recommended amount of concentrate and water according to the directions found under the desired insect to control. Use a clean sprayer. Apply insecticide when insects first appear and repeat not more frequently than every 3 days. Flush sprayer or watering can with clean water after each use.

Use on Lawns: Bent, Bermuda, Bluegrass, Dichondra, Fescue, Irish Moss, Merion and St. Augustine.

Pests: Ants Armyworms, Chinch bugs, Crickets, Cutworms, Deer Ticks, Grasshoppers, Japanese beetle grubs, Mole crickets and Sod webworms. Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For heavy infestations, repeat application after 2 weeks.

For ants, crickets and grasshoppers use [4 tsp in 1 gallon to cover 333 sq. ft.] 2 fl. oz. in 3 gallons of water to cover 1,000 sq. ft. For other listed insects, use 6 fl. oz. in 10 gallons of water to cover 1,000 sq. ft. [For listed insects, 16 fl. oz. treats 2,650 sq. ft.]

Use on Trees and Shrubs: Arizona Cypress, Azalea, Birch, Non-Bearing Citrus, Conifers, Elm, English Ivy, Euonymus, Weeping Fig, Fir, Honeysuckle, Lilac, Mock-Orange, Oak, Palm, Philodendron, Pine, Poinsettia, Tulip, Poplar, Rhododendron, Taxus.

Pests: Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Inchworms, Leafminers, Mealybugs, Spider mites, Whiteflies. Fall webworms on Cherry and Honeysuckle, Budworms on Conifer, Elm leaf beetles, Elm spanworms and Fall webworms on Elms, Tussock moths on Douglas and Grand Firs, Citrus black flies on Non-Bearing Citrus, Oakworms on Oaks, Pine beetles, Pine moths, Pine needleminers, and Needle scale on Pines.

Use 1 fl. oz. in 1 gallon of water. Apply as a thorough spray, wetting leaves and branches but do not let runoff occur. Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature range is from 50 to 75°F and when there is little or no wind. Spray at the first sign of insects. To maintain effectiveness, spray as reinfestation occurs, but use intervals of 4-8 days.

Use on Roses and Flowers: Ageratum, Aster, Azalea, Begonia, Coleus, Common Ninebark and Snowberry, Exacum, Gladiolus, Gold Bells, Hypoestes, Ivy, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Pansy, Pea Shrub, Petunia, Poinsettia, Rhododendron, Rose, Snapdragon, Zinnia.

Pests: Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Fall webworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider mites, and Whiteflies.

Use 1 fl. oz. in 1 gallon of water. Apply as a thorough spray, wetting the plants thoroughly, but do not let runoff occur. Try to spray underside of leaves and penetrate dense foliage. Spray in the late afternoon or evening, when the temperature range is from 50 to 75°F and when there is little or no wind. Spray at the first sign of insects. To maintain effectiveness, spray as reinfestation occurs, but use intervals of 4-8 days.

To kill Gypsy Moth Caterpillars and Adults, Adult Japanese Beetles, Eastern and Forest Tent Caterpillars: Dilute 4 parts water (12.8 fl. oz. per gallon of water) (0.075%). Apply to insure thorough coverage of upper and lower surfaces.

On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches, and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers, making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

Head and Stem Brassica Vegetables: Broccoli, Chinese Broccoli, Brussels Sprouts, Cauliflower, Cabbage, Rape Greens.

Pests: Corn earworms, Beet armyworms, Loopers (Alfalfa, Cabbage), Tobacco budworms.

How to Use: Use 2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 7 days as needed. A maximum of 0.15 lbs. of active ingredient may be applied per acre per season. Can be applied up to 1 day prior to harvest.

Head Lettuce:

Pests: Corn earworms, Beet armyworms, Loopers (Alfalfa, Cabbage), Tobacco budworms.

How to Use: Use 2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 5-10 days as needed. A maximum of 0.15 lbs. of active ingredient may be applied per acre per season. Can be applied up to 5 day prior to harvest.

Peppers: Includes: Bell Pepper, Chili Pepper, Sweet Peppers.

Pests: Cabbage loopers, Corn earworms, European corn borers, Flea beetles, Pepper weevils, Vegetable leafminers.

How to Use: Use 2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 7 days as needed. Do not apply more than 0.15 lbs. of active ingredient per acre per season. Can be applied up to 1 day prior to harvest.

Eggplant:

Pests: Colorado Potato beetles, Vegetable leafminers.

How to Use: Use 3 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 7 days as needed. Do not apply more than 0.15 lbs. of active ingredient per acre per season. Can be applied up to 1 day prior to harvest.

Bulb Vegetables; Onions, Leeks, Garlic.

Pests: Cutworms, Armyworms, Onion Thrips, Aphids.

How to Use: Use 2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 7 days as needed. Do not apply more than 0.125 lbs. of active ingredient per acre per season. Can be applied up to 7 day prior to harvest.

Sweet Corn:

Pests: Armyworms, (Fall, Southern), Corn earworms, European corn borers.

How to Use: Use 1.5 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 3 to 5 days and when needed. Do not exceed 0.15 lbs. of active ingredient per acre per season. Can be applied up to 3 day prior to harvest.

Tomatoes:

Pests: Beet and Southern Armyworm, Tomato Pinworm, Tomato Fruitworm, Horn Worms, Granulate Cutworms, Colorado Potato Beetle, Cabbage Looper.

How to Use: Use 1.5 fl. oz. in 1 gallon of water per 1,000 sq. ft. Apply every 7 days as needed. Do not apply more than 0.15 lbs. of active ingredient per acre per season. Can be applied up to 1 day prior to harvest.

House Plants Indoors: To protect plants in non-commercial greenhouses and lath houses, and for indoor landscaping. Ageratum, Ardisia, Aster, Baby's Breath, Begonia, Birds nest Fern, Bleeding Heart, Carnation, Chrysanthemum, Coleus, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumb cane, English Ivy, Fuchsia, Gladiolus, Grape Ivy, Hypoestes Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia.

Pests: Aphids, Armyworms, Corn earworms, Cutworms, Cabbageworms, Exposed thrips, Fall cankerworms, Fungus gnats, Inchworms, Japanese beetles, Leafminers, Leafrollers, Mealybugs, Scales, Spider mites, Spring cankerworms, Whiteflies.

How to Use on Indoor House Plants Listed Above: Use 1 fl. oz. in 1 gallon of water. Spray will provide contact and residual control. Use sufficient spray to obtain full coverage of all foliage. Adjust spray volume used to plant size. Spray directly on insects. If application is to be made indoors, remove infested plants from living areas prior to application. Spray plants in an area not likely to be occupied by people or pets (particularly fish bowls or tanks). When leaves dry, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury. Use with care on Ferns, Palms and Fuchsia. As with all plants listed above, especially carnations, chrysanthemums, and roses, there exists a wide variety of susceptibility to injury. To ascertain possible toxicity to plants, spray a few plants as described above, wait 24 hours, and check for possible injury. Repeat application may be necessary to control citrus black flies on non-bearing citrus and adult Japanese beetles on roses.

USE IN AND AROUND STRUCTURES:

Do not use in food areas of food handling establishments, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms, but excluding areas where foods may be prepared or held. In the home, cover all food processing surfaces and utensils during treatment or thoroughly wash before use following treatment. Cover or remove exposed food. NOT FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS.

Non-food/feed areas include (but are not limited to) garbage rooms, lavatories, entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

AS A SURFACE SPRAY TO CONTROL: German Cockroaches, Waterbugs, Palmetto Bugs, Ants, Asian Lady Beetles, Silverfish, Crickets, Millipedes, Sowbugs, Pillbugs, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drug Store Beetles, Darkling Beetles, Mealworms, Grain Mites and Cadelles.

For initial cleanup of severe insect infestation, dilute at a rate of 4 parts concentrate in 100 parts water (5.12 fl. oz. per gallon) (32 teaspoons per gallon) (0.03%).

For normal infestations, dilute 2 parts concentrate in 100 parts water (2.56 ounces per gallon) (0.015%).

For maintenance use, dilute 1 part concentrate in 100 parts water (1.28 ounce per gallon) (4 teaspoonfuls per gallon) (0.0075%).

Use a good sprayer and adjust to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery and behind shelves and drawers. Spray bookcases for Silverfish. Spray Ant trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at a rate of 1 gallon to 750 sq. ft. of surface.

TO CONTROL Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving.

TO KILL Deer Ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor and floor covering. Place fresh bedding in animal quarters following treatment.

GENERAL DIRECTIONS - OUTDOOR USE:

- For crack and crevice treatments: Treat surfaces to ensure thorough coverage but avoid runoff. To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

SPRAY DRIFT ADVISORIES

- Take precautions to minimize spray drift.

Barrier Spray for Nuisance Pests: Such as Ants, Clover mites, Cluster flies, Crickets, Earwigs, Flying moths, Millipedes, Mosquitoes, Sowbugs (pillbugs), and Tent caterpillars.

For mosquitoes, dilute to a rate of 0.01% and greater to achieve up to two (2) week's residual control when applied to non-porous items, such as screens and siding. Before treating siding of house, spray in an inconspicuous place and observe for staining.

Use 6.4 fl. oz. in 1 quart of water (0.15%), treating 400 linear feet with a 6-inch spray band. [Use 26 2/3 fl. oz. in 1 gallon of water to treat 1,600 linear feet with a 6-inch spray band.]

Outside Surfaces of Buildings:

Remove all exposed food and cooking utensils. Cover all food handling surfaces or wash thoroughly after treatment and before use. Do not use on edible crops.

Hold sprayer two (2) to three (3) feet from surface to be treated and spray infested surfaces of patio or picnic areas, hitting as many insects as possible. Spray to slightly moisten the surface.

Pests: Ants, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs).

How to Use: To help prevent infestation of buildings [such as Homes, Commercial buildings, Institutions, Warehouses, Theaters, Office buildings, Schools, Motels, Hotels, Restaurants, Hospitals, Cattle barns, Horse barns, Swine houses, Kennels, Milking parlors, Milk rooms, Dairies and Poultry houses], treat the building foundation to a height of 2 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, picnic areas, garages, refuse piles, and other areas where these pests congregate or have been seen, hitting as many insects as possible. Spray to slightly moisten the surfaces. [Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, under patio furniture, in garages and other places where these insects alight or congregate.] To maintain effectiveness, reapply as reinfestation occurs.

Other Outside Areas:

Pests: Ants, Crickets, Earwigs, Deer Ticks.

How to Use: For treatment of localized infestations of these insects in areas where there are weeds, shrubs or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as reinfestations occur.

Pests: Cluster flies, Flying moths, Mosquitoes.

How to Use: For use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, under patio furniture, in garages and other places where these insects alight or congregate.

Pests: Tent caterpillars.

How to Use: Application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent, however, they should die within several hours.

FOR HORSES: Do not apply on foals under 3 months of age.

THIS PRODUCT MUST BE DILUTED PRIOR TO USE. To prepare dilutions, add the required amount of concentrate to water and blend thoroughly. Dilution rates on this label will yield the following ready-to-use spray mixes:

RATE	OZ. CONC/GAL.	Zeta-Cypermethrin CONC.
1:10	12.8	0.075%

TO CONTROL Stable Flies, Mosquitoes, and Gnats: Apply as a wipe or spray to wet the hair thoroughly, with particular attention to topline, underline, flanks, withers, and other infested areas. Repeat treatment at intervals of 5 to 12 days for small insect populations or as needed when flies are emerging in large numbers.

TO CONTROL Stable Flies and Mosquitoes: Apply a quart per adult animal as a wipe or spray to wet the hair thoroughly with particular attention to the legs, flanks, barrel, topline and other body areas commonly attacked by those flies.

Sensitivities may occur after using ANY pesticide product for animals. If signs of sensitivity occur, bathe your animal with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing animals.

STORAGE AND DISPOSAL

Follow proper disposal procedures on this label. Siga las indicaciones del etiquetado para el desecho apropiado del producto.

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE:

Store in a cool, dry place, preferably a locked storage area. Store above freezing. Do not store diluted spray. Store with container tightly closed. Always store pesticide in the original container; do not transfer to food or drink container. Store away from food and pet food. Avoid contamination of food, feed, and domestic water supplies.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Do not pour or dispose down-the-drain or sewer. Call your local solid waste agency for local disposal options.