

1021-2560

2/5/2014

1110



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

FEB - 5 2014

Thomas A. Lennan  
McLaughlin Gormley King Company  
8810 Tenth Avenue North  
Minneapolis, MN 55427

Subject: Notification to Add a Placeholder for Company Logo and Associated Marketing Phrase  
EPA Reg. #: 1021-2560

Dear Mr. Lennan:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated January 21, 2014 for the product 1021-2560. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the request to add a placeholder for the company logo and associated marketing phrase falls within the scope of PRN 98-10.


If you have any questions, please call me directly at 703-305-6701 or Carlyn Petrella of my staff at 703-347-0439.

Sincerely,

A handwritten signature in black ink, appearing to read "Richard Gebken".

Richard Gebken  
Insecticide Branch PM 10 Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs

2/10

	United States Environmental Protection Agency Washington, DC 20460	X	Registration Amendment Other	OPP Identifier Number <b>NOTIFICATION</b>
	<b>Application for Pesticide - Section I</b>			<b>FEB 05 2014</b>
	1. Company/Product Number 1021-2560		2. EPA Product Manager Richard Gebken	
4. Company/Product (Name) EVERGREEN Pyrethrum Concentrate		PM# 10		<input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
5. Name and Address of Applicant (Include Zip Code) McLAUGHLIN GORMLEY KING COMPANY D/B/A MGK 8810 TENTH AVENUE NORTH MINNEAPOLIS, MN 55427  <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to:  EPA Reg. No. _____  Product Name _____		

**Section - II**

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification pursuant to PR-Notice 98-10

**Section - III**

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
<i>*Certification must be submitted</i>	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
			No. per container	<input type="checkbox"/> Other (Specify)	
3. Location of Net Contents information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 1 Pint, 1 Gallon		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Thomas A. Lennan	Title Regulatory Affairs Specialist	Telephone No. (Include Area Code) (763) 544-0341
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received  (Stamped)
2. Signature 		
3. Title Regulatory Affairs Specialist		
4. Typed Name Thomas A. Lennan		5. Date 1/21/2014



January 21, 2014

Document Processing Desk (NOTIF)  
U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs (H7504P)  
Room S-4900, One Potomac Yard  
2777 South Crystal Dr.  
Arlington, VA 22202-4501

Via Courier

Subject: EVERGREEN® Pyrethrum Concentrate  
EPA Reg. No. 1021-2560  
Notification to add place holder for MGK's graphic logo and a marketing phrase.

Dear Sir or Madam:

By Notification we are adding a place holder for MGK's graphic logo and the marketing phrase "MGK® Insect control solutions."

Enclosed are 2 copies of the revised label. One copy has changes highlighted.

Notification of label change per PR Notice 98-10. This notification is consistent with the guidance in PR Notice 98-10 and the requirements of EPA's regulations at 40 CFR 152.46. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C Sec. 1001 to willfully make any false statement to EPA. I further understand that if the notification is not consistent with the requirements of PR Notice 98-10 and 40 CFR 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

If there are any questions or you need anything further, please contact this office.

Sincerely,

McLAUGHLIN GORMLEY KING COMPANY

Thomas A. Lennan  
Regulatory Affairs Specialist

Enclosure

8810 Tenth Avenue North  
Minneapolis, MN 55427  
TOLL FREE 800.645.6466  
TEL 763.544.0341  
FAX 763.544.6437  
WWW.MGK.COM

4110

GROUP	3A	INSECTICIDE
-------	----	-------------

# EVERGREEN® Pyrethrum Concentrate



[ ] denotes optional wording. [Graphic: MGK® Logo] [MGK® Insect Control Solutions] Kills Biting Flies Kills Fleas Kill Brown Dog Ticks Kills Spiders (excluding Brown Recluse Spiders) Kills Stored Products Pests Kills Listed Flying Insects Kills Listed Crawling Insects Kills Cockroaches	Kills Ants (excluding fire and Pharaoh ants) Quick Knockdown Fast Acting Effective ULV Spray Contact Spray Space Spray Area Spray In Poultry Houses On Poultry Provides rapid knockdown & kill of plant pests Kills more than 140 insects For use in food handling, processing and storage areas	Non-flammable Low odor Non-corrosive to spray equipment  One gallon treats up to 28 acres  For Indoor and Outdoor production  For use in food handling, processing and storage areas	For use in Barns, Dairies and Poultry Houses  Contains pyrethrins – a botanical insecticide derived from chrysanthemums  Can be tank mixed with other insect control products  For use in Restaurants
--	--	--	---

**NOTIFICATION**  
**FEB 05 2014**

ACTIVE INGREDIENT:	Pyrethrins, a botanical insecticide .....	5.00%
OTHER INGREDIENTS .....		95.00%
		100.00%

EVERGREEN® - Registered trademark of McLaughlin Gormley King Company

## KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> <li>Immediately call a poison control center or doctor.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
IF IN EYES:	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-740-8712 for emergency medical treatment information.

Manufactured by



8810 Tenth Avenue North  
Minneapolis, MN 55427

EPA Reg. No. 1021-2560

EPA Est. No. 1021-MN-2

Net Contents: \_\_\_\_\_

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Harmful if absorbed through skin. Harmful if swallowed. Causes moderate eye irritation. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using toilet. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants; chemical-resistant gloves, such as, Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton; shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep PPE and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS:**

Users should:

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic organisms, including fish and invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA. Do not contaminate water when disposing of equivalent wash-waters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS**

- Apply this product only as specified on this label.
- Do not contaminate food or feedstuffs.
- Do not use product directly on domestic (companion) animals (dogs, cats, puppies or kittens).
- Not for use in outdoor residential misting systems.
- Do not make applications during rain.
- Do not use in aircraft cabins.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not enter or allow others to enter until sprays have dried.
- Do not enter or allow to enter until vapors, and mists, and/or aerosols have dispersed, and the treated area has been thoroughly ventilated.
- Do not remain in treated area. Exit area immediately and remain outside the treated area until aerosols, vapors, and/or mists have dispersed.
- When used indoors as a crack-and-crevice, surface, or space spray applications. Remove or cover exposed food and drinking water before application. Remove or cover dishes, utensils, food processing equipment and food preparation surfaces or wash them before use.
- When used in indoor food handling/ processing facilities. Do not make space spray applications when the facility is in operation. Prior to space spray applications, cover or remove exposed food. Prior to space spray applications, cover food processing surfaces or clean after treatment and before use.
- When used in dairy barns or facilities: Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios. Except as a spot or crack and crevice treatment. During application do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. All outdoor applications, if permitted elsewhere on the label, must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses, if allowed elsewhere on this label:
  1. Applications to soil or vegetation, as listed on this label, around structures;
  2. Applications to lawn, turf, and other vegetation, as listed on this label;
  3. Applications to the side of a building, up to a maximum height of 3 feet above grade;
  4. Applications to the underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, or overhang, awning or other structure;
  5. Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width;
  6. Applications to vertical or horizontal surfaces when surrounded by or directly above soil, lawn, turf, mulch, vegetation, or other porous surface, where runoff to surface water is unlikely to occur.
- Do not wet plants to the point of runoff or drip.
- Remove pets, birds and cover fish aquarium before application.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas (that is permitted under Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Long-sleeved shirt and long pants;
- Chemical-resistant gloves, such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton;
- Shoes plus socks.

For any requirements specific to your State or Tribe, consult the agency for pesticide regulation.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR, Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

Wear protective clothing when using or handling this product to help avoid exposure to eyes and skin. Eye protection, gloves, a long-sleeved shirt and long pants are recommended.

Allow spray to dry before allowing adults, children or pets on treated areas.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**PESTICIDE RESISTANCE MANAGEMENT:**

EVERGREEN® Pyrethrum Concentrate contains a Group 3A insecticide. Insect biotypes with acquired resistance to Group 3A may eventually dominate the insect population if Group 3A insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by EVERGREEN® Pyrethrum Concentrate or other Group 3A insecticides.

To delay insecticide resistance consider:

- Avoiding the consecutive use of EVERGREEN® Pyrethrum Concentrate or other group 3A insecticides that have a similar target site of action, on the same insect species.
- Using tank mixtures or premixes with insecticides from a different target site of action Group as long as the involved products are all registered for the same use and have different sites of action.
- Basing insecticide use on a comprehensive IPM program.
- Monitoring treated insect population for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for insecticide resistance management and or IPM recommendations for the specific site and resistance pest problems.

INDOOR AREAS for use in and around: Aircraft (not aircraft cabins), Animal Areas and Quarters, Animal Control Centers, Animal Shelters, Apartments, Arcades, Assisted Living Facilities, Attics, Auditoriums, Bakeries, Barns, Bars, Basements, Bathrooms, Beverage Plants, Boats, Bottling Plants, Breweries, Brewery Warehouses, Buildings, Buses, Cabinets, Cabins, Cafeterias, Campers, Canneries, Casinos, Cattle Barns, Cattle Stalls, Churches, Circus Tents, City Missions, Classrooms, Closets, Club Houses, Coffee Shops, Colleges, Commercial Buildings, Communication Centers, Concession Stands, Condominiums, Court Houses, Crawl Spaces, Crematoriums, Dairies, Dairy Stalls and holding areas, Day Care Centers, Delicatessens, Department Stores, Diners, Dog Houses, Storage or Processing Areas that contain Dried Fruit Products, Drugstores, Dumpsters, Dwellings, Egg Processing Plants, Factories, False-Ceilings, Food Processing Plants, Food Storage Areas, Freight containers, Frozen food plants, Fruit Packing Sheds, Funeral parlors, Garages, Garbage Bins, Garbage Cans, Garbage Compactors, Garbage Trucks, Gas Stations, Grain Elevators, Grain Harvesting and Handling Equipment, Grain Mills, Granaries, Grocery Stores, Health Care Facilities, Herb drying and processing facilities, Herb warehouses, Homes, Horse Stables and Barns, Horse Trailers, Hospices, Hospitals, Hotels, Indoor Eating Establishments, Industrial Installations, Institution Dining Areas, Institutions, Jails, Kennels, Laboratories, Laundromats, Lavatories, Libraries, Liquor Stores, Livestock Barns, Livestock Hauling Equipment, Loafing Sheds, Locker Rooms, Mausoleums, Meat Packing Plants, Mess Halls, Milk Rooms, Milking Parlors, Mills, Mini Storage, Mobile Canteens, Mobile Homes, Morgues, Mortuaries, Motels, Movie Theaters, Museums, Mushroom Processing Areas, Nursing Homes, Office buildings, Other Public Buildings, Parking Garages, Passenger Railroad Cars, Peanut Warehouses, Pool Halls, Poultry Houses, Poultry Plants, Pounds, Prisons, Public Storages, Rabbit Processing Plants, Railroad Cars, Recreational Vehicles, Research Animal Quarters, Restaurants, Retail Stores, Rice Mills, Schools, Seed production facilities, Seed warehouses, Ship Holds, Ships, Smoke Shops, Spice plants, Spice production facilities, Sports Stadiums, Stables, Storage Bins (areas), Stores, Supermarkets, Swine Houses, Taverns, Textile Mills and Warehouses, Theaters, Tobacco Factories, Tobacco Warehouses, Tool Sheds, Trains, Transportation Equipment, Truck Trailers, Trucks, Tunnels, Universities, USDA Inspected Facilities, Utilities, Utility Rooms, Vending Machines, Veterinary Hospitals, Wall and Equipment voids, Warehouses, Washrooms, Wheat Mills, Wholesale Stores, Wine Cellars, Wineries and Zoos.

OUTDOOR AREAS for use in and around: Alleys, Amphitheaters, Amusement parks, Athletic Fields (baseball, football, soccer), Back yards, Balconies, Baseball Fields, Bleachers, Bridges, Campgrounds, Car Washes, Circuses, Construction Sites, Corrals, Courtyards, Curbs, Drive-In Restaurants, Drive-In theaters, Dumpsters, Exterior of Homes and other Buildings, Fair Grounds, Feedlots, Front Yards, Game Fields, Garbage Dumps, Gardens, Gazebos, Golf Courses, Grandstands, Greenhouses, Junk Yards, Kennels, Levees, Livestock Pasture, Manure Piles, Nurseries, Outdoor Eating Establishments, Outdoor Growing Crops, Parking Lots, Parks, Patios, Playgrounds, Racetracks, Ranges, Recreation Areas, Rest Areas, Roadsides, Rooftops, Sanitary Sewers (Manholes), School Yards, Sidewalks, Side yards, Stadiums, Tennis Courts, Trees / Shrubs / Flower and Foliage Plants, Truck Stops, Underpasses, Urban Areas, Utility Rooms Outdoors, Weigh Stations, Wild Animal Parks, Wood Piles, Zoos and other similar areas.

FOR USE ON: Beefalo, Birds, Buffalo, Calves, Cattle – beef and milking, Chickens, Donkeys, Ducks, Elk, Emus, Exotics, Geese, Goats, Hogs, Horses, Lambs, Livestock, Llamas, Mules, Ostrich, Oxen, Pheasants, Ponies, Poultry, Pullets, Quail, Ratites, Roosters, Sheep, Swine and Turkeys.

TO KILL THE FOLLOWING INSECTS: Angoumois Grain Moths, Ants (excluding fire and Pharaoh ants), Aphids, Apple Maggots, Armored Scales, Armyworms, Artichoke Plume Moths, Asparagus Beetles, Bagworms, Bean Beetles, Beet Armyworms, Beetles, Bermudagrass Mirids, Billbugs, Blister Beetles, Blow Flies, Boll Weevils, Box Elder Bugs, Branch and Twig Borers, Brown Dog Ticks, Cabbage Loopers, Cabbage Maggots, Caddisflies, Cankerworms, Carrot Weevils, Caterpillars, Chalcids, Cheese Skippers, Cherry Fruit Flies, Cigarette Beetles, Clover Mites, Clover Weevils, Cockroaches, Codling Moths, Colorado Potato Beetles, Confused Flour Beetles, Corn Earworms, Crane Flies, Crickets, Cross-striped Cabbageworms, Cucumber Beetles, Cucumber Beetles (12-spotted), Dark Mealworms, Darkling Beetles (lesser meal worm), Diamondback Larvae and Moths, Dried Fruit Beetles, Drugstore Beetles, Earwigs, Eastern Tent Caterpillars, Elm Leaf Beetles, Eriophyid Mites, European Pine Tip Moths, Fire worms, Flea Beetles, Fleas, Forest Tent Caterpillars, Fruit Flies, Fruit tree Leaf rollers, Fruitworms, Fungus Gnats, Garden Symphylan, Glassy Winged Sharpshooters, Gnats, Grain Mites, Grape Leaf Skeletonizers, Grape Leafhoppers, Grape Mealy bugs, Grasshoppers, Green bugs, Green Fruit Worms, Green Peach Aphids, Greenhouse Thrips, Gypsy Moths (adults & larvae), Harlequin Bugs, Heliothis spp., Horn Flies, Hornworms, House Flies,

Imported Cabbageworms, Indian Meal Moths, Japanese Beetles, Katydid, Lace Bugs, Leaf-footed plant bugs, Leafhoppers, Leafminers, Leafrollers, Leafliers, Lice, Loopers, Lygus, Maggots, Mealy Bugs, Mediterranean Flour Moths, Mexican Bean Beetles, Midges, Millipedes, Mites, Mole crickets, Moths, Mushroom Flies, Navel Orangeworms, Olive Fruit Flies, Onion Maggots, Orange Tortrix, Pacific flatheaded borers, Pear Psyllids, Pepper Weevils, Pink bollworms, Potato Leafhoppers, Potato Tuberworms, Poultry Lice, Proba bugs, Psyllids, Red Flour Beetles, Rice Weevils, Saltmarsh caterpillars, Saw Toothed Grain Beetles, Scales, Sharpshooters, Sheep "Tick" or Ked, Shore flies, Shot hole borers, Silverfish, Skippers, Small Flying Moths, Soft Scales, Southern chinch bugs, Sow bugs, Spider Beetles, Spiders (excluding Brown Recluse Spiders), Springtails, Squash bugs, Stable Flies, Stink Bugs, Tarnished Plant Bugs, Thrips, Tobacco Moths, Tomato budworms, Tomato bugs, Tomato Fruit worms, Tomato Hornworms, Tomato pinworms, Tomato Russet Mite, Tussock moths, Vine mealy bugs, Vinegar Flies, Webworms, Weevils, Western Yellow-striped armyworms, Whiteflies and Yellow Mealworms.

**IMPORTANT NOTE:** It is recommended that the final spray mix be buffered to a pH of 5.5-7.0. Outside of this range pyrethrins can degrade and the product will lose effectiveness.

To avoid possible harm to honey bees and other beneficial insects, it is advisable to apply when the non-target insects are least active.

Pyrethrins degrade rapidly in sunlight. To ensure the best performance, coverage is key. Use spray equipment that will provide the best coverage and direct contact with as many insects as possible.

**TANK MIX:** This product may be tank mixed with other insecticides, acaricides, fungicides, adjuvants, and wetting agents. This application should conform to accepted use precautions and directions for both products. Tank mix applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

Prior to tank mixing, a compatibility test should be conducted using the proper proportions of products and water to ensure the physical compatibility of the mixture.

**PHYTOTOXICITY NOTE:** Plant safety is an important consideration when using insecticides in a greenhouse. However, it is not possible to evaluate the phytotoxicity of EVERGREEN Pyrethrum Concentrate on numerous plant varieties that may react differently to insecticides in different growth stages or under varying environmental conditions. Before making widespread applications of EVERGREEN Pyrethrum Concentrate or tank mix combinations, treat a limited number of plants and observe for phytotoxicity over a 10-day period.

**GROWING CROPS IN GREENHOUSES AND NURSERIES:**

Apply 4.5 to 18 fluid ounces per acre (333 ml to 1,331 ml/Ha) by ground equipment in sufficient water for thorough coverage. This product may be applied by air at the rate of 4.5 to 18 fluid ounces per acre (333 ml to 1,331 ml/Ha) in a minimum of 20 gallons (75.6L) of water. Mix only enough for immediate use. Spraying should begin when insects first appear. Do not wait until plants are heavily infested. Do not apply more than 1 time per day.

**ORNAMENTALS:** Including: African Violet, Ageratum, Aster, Azalea, Begonia, Calceolaria, Calendula, Calla, Camellia, Carnation, Ceanothus, Cineraria, Chrysanthemum, Coleum, Cyclamen, Cypress, Daffodil, Dahlia, Delphinium, Dogwood, Elm, Eucalyptus, Fern, Ficus, Foliage Plants, Fuchsia, Gardenia, Geranium, Gladiolus, Gloxinia, Gypsophila, Holly, Hyacinth, Hydrangea, Iris, Lily, Maidenhair Fern, Marigold, Narcissus, Palm, Pansy, Peony, Pelargonium, Petunia, Philodendron, Phlox, Pine, Pyracantha, Rhododendron, Roses, Rubber Plant, Snapdragon, Sweet Pea, Tulips, Viburnum, Wandering Jew, Yew, Zinnia and Andromeda, Arborvitae, Ash, Beech, Birch, Boxwood, Cotoneaster, Crabapple, Euonymus, Fir, Firethorn, Forsythia, Hawthorn, Hemlock, Hickory, Honey Locust, Horse Chestnut, Juniper, Larch, Laurel, Lilac, Linden, Mimosa, Myrtle, Oak, Pine, Privet, Tulip Tree, Viburnum, Willow.

**USE ON GREENHOUSE FRUIT, VEGETABLE, FLOWER AND FOLIAGE PLANTS:** Combine 16 to 32 fluid ounces with 100 gallons (1.25 ml to 2.50 ml/L) of water for applications with conventional sprayers or (7 to 15 ml) ¼-½ of a fluid ounce per gallon (2 ml to 4 ml/L) of water for applications with handheld or backpack sprayers.

**FOR USE OUTDOORS ON TREES, SHRUBS, FLOWERS AND FOLIAGE PLANTS:** Combine 16 to 32 fluid ounces of EVERGREEN Pyrethrum Concentrate with 100 gallons (1.25 ml to 2.50 ml/L) of water for applications with conventional sprayers or 16 to 32 fluid ounces of EVERGREEN Pyrethrum Concentrate with 10 gallons (12.5 ml to 25.0 ml/L) of water for applications with low volume mist blowers or (7 to 15 ml) ¼-½ of a fluid ounce per gallon (2 ml to 4 ml/L) water for applications with handheld or backpack sprayers.

**FOR USE ON AND AROUND LISTED OUTDOOR AREAS:** To kill insects on Turf, Ornamental Plants, Gardens, Landscaping, Foundation Walls and Perimeters mix (7 to 15 ml) ¼ to ½ of a fluid ounce of EVERGREEN Pyrethrum Concentrate per gallon (2 ml to 4 ml/L) of water for applications with compressed air sprayers or pump driven sprayers.

**FOR USE IN VEGETABLE AND TRUCK GARDENS:** For commercial production of listed crops to kill listed insects dilute ¼ to ½ oz. of EVERGREEN Pyrethrum Concentrate per gallon of water (2 ml/ Liter of water to 4 ml/L of water). Apply diluted solution with all types of agricultural sprayers, including compressed air sprayers or pump driven sprayers.

**USE INDOORS ON TREES, SHRUBS, FLOWERS AND FOLIAGE PLANTS:** Combine 16 to 32 fluid ounces of EVERGREEN Pyrethrum Concentrate with 100 gallons (1.25 ml to 2.50 ml/L) of water for applications with conventional sprayers or (7 to 15 ml) ¼ to ½ of a fluid ounce of EVERGREEN Pyrethrum Concentrate per gallon (2 ml to 4 ml/L) of water for applications with handheld or backpack sprayers.

**FOR USE ON HARVESTED FRUITS AND VEGETABLES:** Apples, Blackberries, Blueberries, Boysenberries, Cherries, Crabapples, Currants, Dewberries, Figs, Gooseberries, Grapes, Guavas, Loganberries, Mangoes, Muskmelons, Oranges, Peaches, Pears, Peas, Pineapples, Plums, Raspberries, Tomatoes.

**DIRECT SPRAY TO FRUITS IN BASKETS, ON TRUCKS OR IN PROCESSING PLANTS:** To kill *Drosophila* spp., *Tephritid* spp., Fruit Flies, Vinegar Flies and other nuisance pests, dilute this concentrate at the rate of 1 part with 900 parts water (1 pint (0.5L) with 125 gallons (472.5L) of water or 5 ml with 10 pints (4.7L) of water). Thoroughly mix the emulsion in the spray tank and treat as follows:

- 1) Apply liberally to fruits and vegetables in baskets, on trucks and in plants. Use sprayers at a high pressure for applying at the rate of five or six pints (2.4L to 2.9L) of diluted spray to a 2-ton load of produce. Direct the spray for maximum coverage of the baskets or hampers. It is important to spray between and beneath the containers.

- 2) Spray the raw stock stacked in the yard.
- 3) Dip baskets in the diluted spray, after dumping the produce to kill adhering larvae and pupae.

**USE DIRECTIONS FOR INDOOR APPLICATIONS:**

Remove pets and birds and cover fish aquariums before spraying.

Kills only at time of application. Listed insect pests must be directly contacted to be effective.

Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm. Patients should be removed prior to treatment. If application is done as a surface spray, patients may re-enter once the spray has dried. If application is done as a space spray, patient rooms should be thoroughly ventilated for 2 hours following spraying, and before returning patients to treated areas.

Do not apply in institutions (including libraries, schools, sports facilities etc.) in the immediate area where occupants are present. Do not apply to classrooms while in use. Vacate rooms prior to treatment. If application is done as a surface spray, rooms should be thoroughly ventilated for 2 hours following spraying, and before re-entering treated areas.

**FOOD AREA USE:** Avoid contamination of food or feedstuffs. Do not apply directly to food. In the home all food processing surfaces, dishes and utensils should be covered during treatment or thoroughly washed before use.

**SURFACE SPRAY:**

**FOR USE AS A SURFACE SPRAY IN LISTED INDOOR AREAS:** to kill accessible, exposed stages of listed crawling insects, dilute 1-3 fluid ounces of concentrate with one gallon (8 ml to 24 ml/L) of water, mix well and apply as a coarse wetting spray at the rate of 1 gallon of spray mixture per 750 square feet (54 ml/m<sup>2</sup>) of surface. To ensure maximum kill of listed crawling insects, spray thoroughly into all cracks, moldings, crevices, walls, floors and surfaces of bins, storage and handling areas. Particular attention should be paid to areas where moisture collects such as drains, sinks, and water pipes and around the base of heavy machinery and equipment, contacting as many insects as possible. To kill ANTS (excluding fire and Pharaoh ants), spray directly into nests, along trails, and directly on all ants observed.

**Dilution Table for Surface Spray**

Fluid ounces of Concentrate Per Gallon of Solution	Pounds of Pyrethrum Per Gallon of Solution.	Percent Pyrethrins in Solution
1	0.0032 (0.38g/L)	0.039%
2	0.0064 (0.77g/L)	0.078%
3	0.0096 (1.15 g/L)	0.118%
4	0.0128 (1.53 g/L)	0.159%
5	0.0160 (1.92 g/L)	0.200%
6	0.0192 (2.30 g/L)	0.242%
7	0.0224 (2.68 g/L)	0.284%
8	0.0256 (3.07 g/L)	0.328%
12	0.038 (4.56 g/L)	0.508%
16	0.051 (6.12 g/L)	0.702%
22	0.070 (8.4 g/L)	1.020%

**FOR USE IN MEAT AND POULTRY PROCESSING FACILITIES:** To kill accessible, exposed stages of listed crawling insects, dilute 9 fluid ounces with one gallon (71.4 ml/L) of water and apply at the rate of 1 gallon to 750 square feet (54 ml/m<sup>2</sup>), paying special attention to force the spray into all cracks and crevices.

**SPACE SPRAY IN FOOD AND NONFOOD AREAS OF LISTED INDOOR AREAS:** For use in mechanical fogging or spraying equipment to kill accessible stages of listed FLYING AND CRAWLING INSECTS. Close room and shut off all air conditioners or ventilating equipment. Dilute at the rate of 1 part concentrate into 1-8 parts water (see table below) and mix well. Apply at the rate of 1 fluid ounce of diluted solution per 1000 cubic feet (1.04 ml/m<sup>3</sup>) of space, filling the room with mist. Direct the space treatment upward and whenever practical, keep doors and windows closed for at least 30 minutes after application. Vacate treated area and thoroughly ventilate before re-entry.

This concentrate may also be diluted with oil. If the product is used in a certified organic facility, check with certification regulations regarding allowable oils. Any oil should also meet Federal Food Additive Regulation requirements as defined by CFR Title 21. Conduct a small jar compatibility test by mixing a small amount of oil and concentrate and observe the physical compatibility of the mixture. Dilute at the rate of 1 part of concentrate with up to 8 parts oil. Apply at the rate of 1 fluid ounce of diluted solution per 1000 cubic feet (1.04 ml/m<sup>3</sup>) of space.

When this product is applied as a space spray in food processing or food handling establishment applications should be confined to time periods when the plant is not in operation.

**Dilution Table for ULV Foggers**

Dilution Ratio:		Ratio of Concentrate to solution by volume.	Ratio of water to solution by volume.	Pounds of Pyrethrum Per 1,000 cu. ft. when applied at 1 fl oz per 1000 cu. ft.	Percent Pyrethrins in Solution
Concentrate	Water				
Undiluted		1.00	0.00	0.0032 (0.051 g/m <sup>3</sup> )	5.00%
1	1	0.50	0.50	0.0016 (0.025 g/m <sup>3</sup> )	2.47%
1	2	0.33	0.67	0.0011 (0.018 g/m <sup>3</sup> )	1.65%
1	3	0.25	0.75	0.00101 (0.016 g/m <sup>3</sup> )	1.23%
1	4	0.20	0.80	0.00064 (0.010g/m <sup>3</sup> )	0.99%
1	8	0.11	0.89	0.00035 (0.0056 g/m <sup>3</sup> )	0.547%



**FOR USE ON SWEET POTATOES IN STORAGE:** To kill Fruit Flies and Vinegar Flies, dilute this concentrate at 1 part to 14 parts water (9 fluid ounces with one gallon of water (71.37 ml/L)). Apply as a space fog with a mechanical fogger capable of producing particles of aerosol size at the rate of 1 gallon diluted spray per 100,000 cubic feet of space (40 ml/L). Apply only when flying insects are present. Several applications may be necessary during periods of heavy infestation, but do not make more than 10 applications.

**FOR USE ON STORED PRODUCTS:** The product can be used to kill accessible stages of listed insects on the following stored commodities: Almond Nutmeat and Shells, Barley, Beans, Birdseed, Buckwheat, Cocoa Beans, Corn, Cottonseed, Dried Apricots, Dried Fruit, Dried Prunes, Figs, Flax, Grain Nuts, Oats, Peanut, Pistachio, Raisins, Rice, Rye, Sorghum, Tobacco, Wheat, and Walnut Nutmeat and Shells held in storage.

**DRIED FRUIT PRODUCTS IN STORAGE OR BEING PROCESSED:** To kill the following stored product pests: Dried Fruit Beetle and Saw-Toothed Grain Beetle. Remove and destroy infested products. Dilute concentrate at the rate of 1 part concentrate to 8 parts water and mix well. Disperse the solution as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. If the solution is applied as a space spray, use one ounce diluted spray per 1000 cubic feet of space (1 ml diluted spray / m<sup>3</sup> of space). Immediately following application, leave the room closed off for 1 hour. Thoroughly ventilate the treated area before re-entry. Do not apply more than 1 time per day. Sweep up and destroy fallen insects.

**ON ALMONDS, PEANUTS, PISTACIOS AND WALNUTS IN BULK OR IN BAGS:** To kill the following stored product insects: Almond Moths, Angoumois Grain Moths, Ants (excluding fire and Pharaoh ants), Cadelles, Cigarette Beetles, Confused Flour Beetles, Drugstore Beetles, Flat Grain Beetles, Granary Weevils, Indian Meal Moths, Lesser Grain Borers, Maize Weevils, Mediterranean Flour Moths, Merchant Grain Beetles, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Saw-toothed Grain Beetles and Square-necked Grain Beetles, dilute 2.0 fluid ounces of EVERGREEN Pyrethrum Concentrate per gallon (15.86 ml/L) of water and apply as a coarse wet spray over the top of stored nuts or the outside surface of stacked bagged nuts at the rate of 4 gallons per 1000 square feet (165 ml/m<sup>2</sup>). Apply at weekly intervals for about 6 weeks and then at 15-day intervals. The first two applications should be applied at the rate of 4 gallons per 1000 square feet (165 ml/m<sup>2</sup>), and subsequent treatments should be applied at the rate of 2 gallons per 1000 square feet (80 ml/m<sup>2</sup>).

**TREATMENT OF STORED GRAIN AND SEED AREAS:** For best results to kill stored product pests in grain and seed storage areas and stored grain and seed, treat the area when it is empty of product, as a protectant applied as commodity is put into storage, and as a surface spray once it is in storage.

**FOR USE IN STORAGE SITES:** This concentrate can be used to treat grain and seed in warehouse bins and trucks, cargo ships, mills, bin hoppers, elevators and conveying equipment as a clean up prior to using them for storage. In mills and elevators, all grain-infested accumulations should be removed from the bin hoppers. All storage areas and conveying equipment should be thoroughly cleaned by sweeping out the waste grain, cobwebs and other debris from the walls and rafters as well as on the floor and door frames with special attention to material lodged in the cracks and crevices. All debris should be removed and burned to kill eggs and insects that might be present.

For farms, particular attention should be given to cleaning up around the used feed and grain bags, grain residues from wagons, harvesting equipment and feed troughs. Newly harvested grain should not be placed in the same bin with carry-over grain, and all carry-over grain stocks that are not treated with grain protectant should be fumigated. These cleaning operations should be done within two or three weeks before harvest.

After above sanitation measures have been employed, spray all areas prior to use for storage with 1 part to 22 parts water (6 fluid ounces with one gallon of water) up to 1 part to 8 parts (1 pint (0.5 L) with one gallon (3.78L) of water). Apply at the rate of one gallon per 750 square feet (54 ml/m<sup>2</sup>) on walls, floors, ceilings and partition boards of bins, paying particular attention to forcing the spray into all cracks and crevices.

**FOR USE AS A GRAIN PROTECTANT:** This concentrate when diluted with water and sprayed directly on grains will effectively protect the grain against grain storage insects for a full season or approximately 8 months. Dilute at the rate of 1 part to 22 parts water (6 fluid ounces with one gallon (48 ml/L) of water). Thoroughly mix the emulsion and apply at the rate of 5 gallons (18.9L) per 1000 bushels of grain as it is carried along a belt or as it enters the auger or elevator. This concentrate may be used in combination with a registered fumigant for use on heavily infested stored products.

Monthly inspections should be made. If the top 2 or 3 inches (0.6 -0.9 m) are infested, re-treat applying at the rate of 2 gallons (7.56 L) of diluted material per 1000 bushels of stored product.

**SURFACE TREATMENT OF STORED GRAIN AND SEED:** To kill Indian Meal Moths, Angoumois Grain Moths and Mediterranean Flour Moths, monthly inspections should be made after the grain is placed in storage. If the top two or three inches are infested, dilute 1 part EVERGREEN Pyrethrum Concentrate with 14 parts of water and apply at the rate of 2 gallons per 1000 square feet (80ml/m<sup>2</sup>) of grain. Rake the mixture into the grain to a depth of 4 inches (10 cm).

**FOR USE ON LISTED ANIMALS:**

- 1) To kill Horn Flies, House Flies and Gnats, dilute at the rate of 1.5 to 3 fluid ounces per gallon (12 ml to 24 ml/L) of water and apply to wet hair thoroughly, paying particular attention to topline, underline, flanks, withers and other infested areas. Repeat treatment at intervals of 5 to 12 days for small insect populations or as needed when flies are emerging in large numbers.
- 2) To kill stable flies, dilute at the rate of 2.5 to 4 fluid ounces per gallon (20 ml to 32 ml/L) of water and apply at a quart per adult animal to wet hair thoroughly, paying particular attention to legs, flanks, barrel, topline and other body areas commonly attacked by these flies. Repeat treatment each week as needed.
- 3) To kill face flies, dilute at the rate of 2.5 fluid ounces per gallon (20 ml/L) of water and apply using spray which produces large wetting droplets. Apply to face of animal in the morning before releasing to pasture. Apply sufficiently to wet the face but not more than 1-1.5 ounces (28.35 g-42.52 g) of spray solution per animal. Repeat daily as needed.
- 4) To kill biting and sucking lice on livestock, dilute at the rate of 1 quart with 60 gallons (226.8L) of water (1.5 fluid ounces with 2 gallons (6 ml/L)) and spray to thoroughly wet hair of animal, including the head and brush of the tail. Repeat treatment in 10 days to kill newly hatched lice.
- 5) To kill poultry lice, using a dilution of 2.5 to 4 fluid ounces of concentrate per gallon (20 ml to 32 ml/L) of water. Spray roosts, walls and nests or cages thoroughly. It is not necessary to remove poultry from the housing unit during treatment. This should be followed by spraying over the birds with a fine mist.

- 6) To kill mites on poultry and in poultry houses, dilute at the rate of 2.5 to 4 fluid ounces per gallon (20 ml to 32 ml/L) of water and spray crevices of roost poles, cracks in walls and cracks in nests where the mites hide. This should be followed by spraying over the birds with a fine mist.
- 7) To kill sheep "Tick" or Ked, dilute at the rate of 1.5 to 3 fluid ounces per 4 gallons (3 ml/L to 6 ml/L) of water and thoroughly wet all portions of the body by dipping or spraying with sufficient pressure and with a nozzle adjustment to give penetration of the wool. Treat at a rate sufficient to wet the animal.
- 8) To kill fleas and brown dog ticks on livestock and to obtain protection against reinfestation, dilute at the rate of 9 fluid ounces per gallon (71 ml/L) of water and wet the animal by dipping or spraying. This product may also be used for livestock/pet quarters. Do not spray pets. For best results against fleas and brown dog ticks on dogs and cats the kennels and/or animal quarters and bedding should be treated. Remove and destroy pet's old bedding. To reduce infestation, thoroughly spray the animal's bedding and sleeping quarters. Also spray the entire inside surface of dog houses. Put fresh bedding in pet's quarters after spray has dried. Also treat dogs and cats [pets] with a registered flea and tick control product before allowing them to reenter treated area.
- 9) To kill adult darkling beetle (lesser meal worm) in poultry houses, dilute 2.0-5.0 teaspoons (10-25 ml) of EVERGREEN® Pyrethrum Concentrate per gallon of water. Using a power or proportioner-type sprayer, apply a uniform spray until wet at pressures of 80 to 100 pounds per square inch to surfaces such as interior walls, roosting posts, structural support pillars, and other surfaces where adult beetles are observed. One gallon of spray mixture covers 250-500 square feet (23 m<sup>2</sup>-46 m<sup>2</sup>). Do not contaminate food/feed or water. For maximum effectiveness, treatments should coincide with each growout or sanitation procedure. Indoor control can be enhanced by making perimeter treatments around the outside of buildings. To prevent immigrating adult beetles, apply a uniform band of spray 2 feet (0.6 m) up and 1-4 feet (0.3-1.2 m) out from the foundation.

**FOR USE IN BARN, DAIRIES, MILKING PARLORS, MILKING ROOMS AND POULTRY HOUSES:** To kill the following flying insects: Fruit Flies, Gnats and Small Flying Moths, dilute at the rate of 2.5 fluid ounces per gallon (19.82 ml/L) of water. Apply as a fog or fine mist (at approximately 2 fluid ounces per 1000 cubic feet of space (2.08 ml/m<sup>3</sup>), directing the nozzle for maximum coverage and above livestock and poultry toward the ceiling and upper corners of the area being treated. For best results, close doors and windows before spraying, and keep them closed for 15 minutes immediately following application. Applicator should exit the treated area immediately, and thoroughly ventilate it before re-entry. Sweep up and destroy fallen insects.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep this product in its tightly closed original container, when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals.

**PESTICIDE DISPOSAL:**  
Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:**  
Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

[For containers with capacities equal to or less than 5 Gallons (18.9 L)]:  
Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Pressure rinse as follows [for containers greater than 5 Gallons (18.9L) too large to shake]: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.