



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

June 12, 2025

Chantha Coursolle
Regulatory Affairs Analyst
McLaughlin Gormley King Company d/b/a MGK
7325 Aspen Lane N
Minneapolis, MN 55428

Subject: Notification per PRN 98-10 – Updating referral statement, adding Wasp and Bees language and Ground Wasp language to “Outdoor Use” section.
Product Name: ULTRATEC DUST INSECTICIDE
EPA Registration Number: 1021-2617
Application Date: 05/23/2025
Case Number: 658707

Dear Chantha Coursolle:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “NOTIFICATION” and placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, please contact Laura Rademacher at Rademacher.Laura@epa.gov.

Sincerely,

Jacquelyn Herrick, Product Manager 03
Invertebrate & Vertebrate Branch 1
Registration Division (7505T)
Office of Pesticide Programs

UltraTec® Dust Insecticide

For Indoor/Outdoor Residential Use only

- [For Household Use
- Household Ornamental & Vegetable Garden Insect Control
- Controls Ants, Fire Ants, Fleas, Ticks, Termites (Localized Control only) and More
- Kills Organophosphate & Carbamate Resistant and Non-Resistant Cockroach Strains
- Odorless
- Long-Lasting Residual Action
- Kills Insects for up to 8 Months
- Kills Carpenter Ants invading treated wall voids for up to 1 year (12 months)
- Does not Clump or Absorb Moisture
- Water-Proof Dust
- Water-Proof - Does Not Absorb Moisture
- Use on Roses, Flowers, and Ornamentals
- 1 Lb. Treats up to 45 Fire Ant Mounds
- Easy to Use
- Convenient
- Kills Fire Ants on Contact - No Watering-in Required
- Kills the Queen
- Ready-to-Use Dry Treatment of Fire Ant Mounds
- Kills the Mound
- Works Fast]

NOTIFICATION

1021-2617

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

06/12/2025

ACTIVE INGREDIENT:

Deltamethrin 0.05%

OTHER INGREDIENTS:..... 99.95%
100.00%

UltraTec® is a registered trademark of McLaughlin Gormley King Company.

PRECAUCIÓN AL CONSUMIDOR: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

KEEP OUT OF REACH OF CHILDREN

CAUTION

[Peel label before use:] See Side/Back/Inside Panel For Additional Precautions [Precautionary Statements] [and] First Aid [and Complete Use Instructions]

FIRST AID	
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of Medical emergencies or health and safety inquiries or in case of fire, leaking or damaged containers, information may be obtained by calling 1-888-740-8712.	

Manufactured by:



7325 Aspen Lane North
Minneapolis, MN 55428

EPA Reg. No. 1021-2617

EPA Est. No. 1021-MN-2

Net Weight: _____

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May be harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product
in a manner inconsistent with its labeling.

IMPORTANT: Read the entire Directions for Use, Precautions and Disclaimer of Warranties before using this product.

UltraTec® Dust Insecticide is a ready-to-use insecticide dust which exhibits effective knockdown and long-lasting control against the pests listed when used according to label directions.

UltraTec Dust Insecticide can be used for long-lasting pest control in and around homes and their immediate surroundings. Permitted areas of use include but are not limited to:

animal quarters	houses
apartments	mobile homes and recreational vehicles
household greenhouses	ornamentals
	vegetables

This product is intended for applications with a hand duster or from the shaker can, with a paint brush or by other suitable means, to hiding and runway areas and other places where insects are found.

Application is prohibited directly into sewer drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

INDOOR USE

In living areas, make applications in such a manner as to avoid deposits on exposed surfaces or introducing the material into the air. To avoid the possibility of staining, do not use on carpets, upholstery and other fabrics. Do not apply as a broadcast application indoors.

To apply insecticide directly into cracks and crevices, use a bulbous duster, or other suitable equipment. Apply lightly and uniformly to infested area. Pay particular attention to cracks and crevices; service ducts; false floors and ceilings; wall void; around electrical and telephone fittings and equipment; around water and sewer pipes; under and behind cabinets; refrigerators and sinks; around windows and door frames; along baseboards; in attics and crawl spaces. The amount to be applied will vary with the site but usually be in the range of 2-3 grams of product per square yard (or 0.5 lbs. per 1,000 sq. ft.).

Ants, Boxelder Bugs, Cockroaches, (including Both Resistant and Non-resistant Strains), Crickets, Fleas, Firebrats and Silverfish:

Apply UltraTec Dust Insecticide thoroughly to all areas where these pests crawl and hide, especially in those areas described above. For ***Ants***, apply to ant trails, around doors and windows and other places where ants enter premises.

Termites, Carpenter Ants and Carpenter Bees: Apply to voids or channels in damaged wooden members of a structure; to cracks, spaces or bearing joints between wooden members of a structure; or between such members and the foundation; and in locations vulnerable to termite attack. Drilling may be necessary prior to applying dust in these locations, but it must be recognized that such drilling and treating of timber in a particular location may not result in complete coverage of infested and vulnerable voids or galleries. This dust can be applied in attics and crawl spaces to prevent entry of drywood and Formosan termites at a rate of 0.5 lb. per 1,000 sq. ft. of floor area. In these treatments, all vents and blowers should be closed. Inspect at 6 months and one year to ensure that colony has been killed.

The purpose of such applications of UltraTec Dust Insecticide is to kill workers or winged forms of subterranean termites, which may be present in the treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement. For active termite infestations, get a professional inspection.

Wall Void Treatments: [To control carpenter ants in wall voids for up to one year [12 months],] apply dust into wall voids through existing openings or by drilling a small hole. Insert end of puffer tip into opening and squeeze container [Insert tip of hand-duster into opening and squeeze] to apply approximately 0.2 grams (0.07 lbs./100 sq. ft.) of dust per wall void [see markings on side of container]. Reseal hole after treatment. Nests in wall voids can be located by listening for insect activity against the wall.

Centipedes, Ground Beetles, Millipedes, Scorpions, Spiders and Sowbugs: Apply dust along and behind baseboards, to window and door frames, corners, pipes, storage locations, attics, crawl spaces, and other areas where these pests may enter or crawl.

Wasps and Bees: It is generally advisable to treat wasp and bee nests in the evening when insects are less active and have returned to the nest. Wear protective clothing if deemed necessary to avoid stings.

Using hand or power duster or other suitable means, with extension tubes if necessary, thoroughly dust nest, entrance and surrounding areas where insects alight. Nests in wall voids can be located by listening with your ear against the wall. Drill a hole in the area, blow dust in and reseal. ~~For ***Ground Wasps***, dust entrance and surrounding areas. For best results, check nests carefully one or two days after treatment to ensure complete kill, then remove and destroy nest to prevent emergence of newly hatched insects. If removal is not feasible, retreat nest if necessary. Do not use this product to treat overhead nests.~~

Webbing Clothes Moths: Use UltraTec Dust Insecticide as a crack and crevice treatment in closets and other storage areas where these pests are found. This is an adjunct treatment and will not control moth larvae already on the clothes. DO NOT apply this product directly to clothing.

Carpet Beetles (Black, Furniture, Common and Varied): Dust where these insects are found, for example: under edges of floor coverings, behind baseboards, under rugs and furniture, and in closets. Do not apply directly to carpeting.

Bed Bugs: Apply to crack and crevices where evidence of bed bugs occurs. This includes bed frames, box springs, inside empty dressers and clothes closets, and carpet edges, high and low wall moldings, and wall paper edges. Do NOT use this product on mattresses, pillows, bed linens, or clothes. Remove all clothes and other articles from dressers or clothes closets before applications. Not recommended for use as sole protection against bed bugs, if evidence of bed bugs is found in/on mattresses use products approved for this use.

STORED PRODUCT PESTS:

To aid in the control of the exposed stages of:

cadellles	merchant grain beetles
cheese mites	red flour beetles
cigarette beetles	rice weevils
confused flour beetles	rusty grain beetles
dark mealworms	sawtoothed grain beetles
drugstore beetles	spider beetles
grain mites	tobacco moths
granary weevils	yellow mealworms
lesser grain borers	

In the home, cover all food processing surfaces and utensils during treatment or thoroughly wash before use. Remove or cover exposed food.

Clean up waste materials, dust, dirt, and other debris. Thoroughly treat floors, walls and other surfaces in storage and handling areas. Apply dust in cracks and crevices, behind and under cupboards, and other difficult to reach areas where insects may hide. DO NOT apply directly on food, utensils or other areas where food is stored.

OUTDOOR USE

Do not water the treated area to the point of runoff. Do not make applications during rain.

ORNAMENTAL PLANT PESTS

For the control of:

aphids	leafrollers
beet armyworms	orange tortrix
blister beetles	periodical cicadas
Boxelder bugs	plant bugs
certain leafminers	rose chafers
(Boxwood, Holly,	rose midges
Oak, Spruce Needle)	rose slugs
elm leaf beetles	tent caterpillars
flea beetles	thrips
Japanese beetles	weevils
leafhoppers	

SHRUBS AND ORNAMENTALS: Apply when insects or their damage first appear and repeat at weekly intervals if necessary. Apply 8 oz. per 1,000 sq. ft.; or thoroughly apply to place a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

ROSES, PETUNIAS, CHRYSANTHEMUMS, GERANIUMS, CARNATIONS AND OTHER RELATED FLOWERS: To control many leaf and flower chewing insects, apply as soon as the insect appears and repeat at 7 day intervals or as long as insects continue to appear. Apply a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

Wasps and Bees: It is generally advisable to treat wasp and bee nests in the evening when insects are less active and have returned to the nest. Wear protective clothing if deemed necessary to avoid stings.

For Ground Wasps, dust entrance and surrounding areas. For best results, check nests carefully one or two days after treatment to ensure complete kill, then remove and destroy nest to prevent emergence of newly-hatched insects. If removal is not feasible, retreat nest if necessary. Do not use this product to treat overhead nests.

PERIMETER TREATMENT: Use UltraTec Dust Insecticide as a residual treatment for control of **Ants, Bees, Centipedes, Cockroaches, Crickets, Firebrats, Fleas, Ground Beetles, Millipedes, Scorpions, Silverfish, Slugs, Sowbugs, Spiders, Ticks and Wasps**. Make a residual treatment around windows and doors, porches, screens, eaves, patios, garages, under stairways, in crawl spaces and other areas where pests hide. To help prevent the invasion of buildings and structures by these pests, apply thoroughly and uniformly to the foundation and crawl spaces where pests are active and may find entrance.

Fleas and Ticks in Rodent Burrows: To Control Fleas and Ticks in and around rodent burrows around homes and other buildings, apply dust with a shaker can. Thoroughly apply dust to adequately cover mound and around the perimeter (18" from the outer edge). Be sure to treat abandoned rodent burrows which are suspected of containing fleas and ticks. Treat for a minimum of 5 days for light rodent populations, 10 days when populations are dense.

HOME VEGETABLE GARDENS: Use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate available duster. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.

VEGETABLE	TARGET PESTS	REMARKS	Application Rate: Ounce of Dust / 100 ft ²	Max Applications per growing season	[Active Ingredient/Acre] (This column is alternate text)	[Packages/ 100 ft ²] (This column is alternate text)
Artichoke, Globe	Aphids, armyworms, cabbage loopers, cribrate weevils, leafminers, lygus bugs (plant bugs), pale striped flea beetles, plume moths, proba bugs	Do not apply more than recommend amount [0.14 lb. a.i./A (active ingredient/acre)] in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest.	2.13	5	0.028 lb.	X packages /100 ft ²
Bulb Vegetables – garlic, leeks, onions, shallots	Cutworms, stink bugs, thrips	Do not apply more than recommend amount [0.112 lb. a.i./A] in one growing season. Allow 5 days between applications. Do not apply within one day of harvest.	2.13	4	0.028 lb.	X packages /100 ft ²
Corn, Sweet	Aphids, armyworms, chinch bugs, click beetles, corn earworms, corn rootworms (adult), cutworms, European corn borers, flea beetles, grasshoppers, Japanese beetles (adult), June beetles (adult), Masked chafers (adult), southern corn borers, southwestern corn borers, stalk borers, stink bugs, webworms	Do not apply more than recommend amount [0.45 lb. a.i./A] one growing season. Do not apply within 24 hours of harvest.	2.13	16	0.028 lb.	X packages /100 ft ²
Corn, Pop	Aphids, armyworms, chinch bugs, click beetles, corn earworms, corn rootworms (adult), cutworms, European corn borers, flea beetles, grasshoppers, Japanese beetles (adult), June beetles (adult), Masked chafers (adult), southern corn borers, southwestern corn borers, stalk borers, stink bugs, webworms	Do not apply more than recommend amount [0.095 lb. a.i./A] in one growing season. Allow 21 days between applications.	1.68	4	0.022 lb.	X packages /100 ft ²
Cucurbit Vegetables – cantaloupes, chayote, Chinese waxgourd, citron melon, cucumber, gherkin, gourds, Momordica species (balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelon, pumpkin, squash, watermelon	Armyworms, cabbage loopers, Colorado potato beetles, corn earworms, cucumber beetles, cutworms, diamondback moth (larvae), flea beetles, grasshoppers, green peach aphids, leafhoppers, leafminers, melonworms, pickleworms, stink bugs, tarnished plant bugs, tobacco budworms, tomato pinworms, vegetable weevils (adult), western plant bugs, whiteflies (adult)	Do not apply more than recommend amount [0.168 lb. a.i./A] in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest.	2.13	6	0.028 lb.	X packages /100 ft ²

Fruiting Vegetables - eggplant, ground cherry, pepinos, peppers, tomatillo, tomato	Aphids, armyworms, cabbage loopers, Colorado potato beetles, cucumber beetles, cutworms, diamondback moths, European corn borers, flea beetles, imported cabbageworms, leafminers, pepper weevils, stink bugs, tarnished plant bugs, thrips, tomato fruitworms (corn earworms), tomato pinworms, variegated cutworms, whiteflies	Do not apply more than recommend amount [0.168 lb. a.i./A] in one growing season. Allow 5 days between applications. Do not apply within 1 day of harvest.	2.13	6	0.028 lb.	X packages /100 ft ²
Root Vegetables - carrot, celeriac, chervil(turnip-rooted), chicory, edible burdock, garden beet, ginseng, horseradish, parsley (turnip-rooted), parsnip, radish, oriental radish, rutabaga, salsify, black salsify, Spanish salsify, skirret, turnip	Armyworms, cabbage loopers, Colorado potato beetles, cutworms, European corn borers, flea beetles, green peach aphids, leafhoppers, lygus bugs (plant bugs), pea aphids, Tarnished plant bugs	Do not apply more than recommend amount [0.14 a.i./A] in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest. Other than radishes, applications may not be made to root vegetables whose tops or leaves are to be harvested for human consumption.	2.13	5	0.028 lb.	X packages /100 ft ²
Tuberous & Corm Vegetables - potato, sweet potato, arracacia, arrowroot, Chinese artichoke, Jerusalem artichoke, edible Canna, bitter and sweet cassava, chayote (root), chufa, dasheen, ginger, leren, tanier, turmeric, yarn bean, true yam	Armyworms, cabbage loopers, Colorado potato beetles, cutworms, European corn borers, flea beetles, green peach aphids, leafhoppers, lygus bugs (plant bugs), pea aphids, Tarnished plant bugs	Do not apply more than recommend amount [0.14 lb. a.i./A] in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest. Applications may not be made to tuberous or corm vegetables whose tops or leaves are to be harvested for human consumption	2.13	5	0.028 lb.	X packages /100 ft ²)

USE DIRECTIONS FOR FIRE ANT TREATMENT

Imported Fire Ant Mound Treatment: Apply UltraTec Dust Insecticide with a shaker can. Thoroughly apply dust to adequately cover mound and around the perimeter (18" from the outer edge of the mound). A second treatment may be necessary if complete control is not obtained. DO NOT DISTURB THE MOUND PRIOR TO APPLICATION. If possible, treat all colonies in the vicinity. Also, treat colonies which have not yet constructed a mound.

[Alternate directions - Evenly sprinkle one tablespoon (approximately 10 grams) of UltraTec Dust Insecticide over the top of each mound. Do not water in or disturb the mound. Apply when temperatures are between 65 and 80°F. Allow 3 to 4 days for maximum control, then repeat as necessary. Do not use food utensils, such as a teaspoon or measuring cups, for food after use with pesticides.]

STORAGE AND DISPOSAL

STORAGE: Store product in original container in a cool, dry, locked place out of the reach of children.

DISPOSAL: Non-refillable container. **If empty:** Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency [or 1-888-740-8712] for disposal instructions. Never place unused product down any indoor or outdoor drain.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of MGK. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MGK MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of MGK is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, MGK disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT MGK'S ELECTION, THE REPLACEMENT OF PRODUCT.

[Alternate Front Panel Bullets:

- *Long-Lasting*
- *For Use on Ornamentals*
- *Controls Many Different Insect Pests*
- *To Control Imported Fire Ants on Lawns and other Residential areas*
- *8 oz. Treats over 20 Fire Ant Mounds*
- *2 Lbs. Treats over 80 Fire Ant Mounds*
- *Fast, Effective Kill of Fire Ants*
- *Fast, Effective Kill of Stinging Ants*
- *Contains Deltamethrin*
- *Contains UltraTec®]*

[] Denotes alternate text in addition to any subset of the directions for use.