



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION
WASHINGTON, D.C. 20460

February 3, 2026

Thomas A. Lennan
Regulatory Affairs Specialist
McLaughlin Gormley King Company, D/B/A MGK
7325 Aspen Lane N
Minneapolis, MN 55428

Subject: Label Amendment – Approval of Addition of Directions for Use to Accommodate Smaller Containers for Product Distribution in Smaller Amounts to Existing Treatment Sites Holding Smaller Volumes of Water
Product Name: Sumilarv® 0.5G
EPA Registration Number: 1021-2819
Application Date: April 3, 2025
Case Number: 652175

Dear Thomas A. Lennan:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 C.F.R. § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 C.F.R. § 152.3.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 C.F.R. § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ

from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Carmen Rodia via e-mail at Rodia.Carmen@epa.gov.

Sincerely,



Tamica L. Cain, Product Manager 10
Invertebrate & Vertebrate Branch 2
Registration Division (7505T)

Enclosure: Stamped "Accepted" Master Label, February 3, 2026

Sumilarv® 0.5G

{Note to reviewer: [] or indicate optional wording}

Controlled release insect growth regulator formulation to break mosquito life-cycle and prevent their emergence

- Prevents mosquitoes from becoming breeding, biting adults
- Novel class of chemistry for public health
- Easy to apply sand granule
- Non-flushing formulation
- Works well even in areas that dry out and rewet
- [Provides up to 150 days (21 weeks) residual control of mosquitoes with an application of (75 g) in storm water catch basins]
- [Provides up to 28 days (4 weeks) residual control of mosquitoes with an application of (25 g) in storm water catch basins]
- [Sumilarv will provide up to 150 days] residual control of mosquitoes with an application of (75g) in storm water catch basins]
- [Sumilarv will provide up to 28 days (4 weeks) residual control of mosquitoes with an application of (25 g) in storm water catch basins]
- [Extended-release insect growth regulator formulation to stop mosquito life-cycle]
- Dosing flexibility allows the user to apply varying amounts of product

- for their desired duration of mosquito control
- For resistance management, Sumilarv 0.5G can be rotated with Bacillus species Group 11 insecticides for mosquito control
- [Prevents adult mosquito emergence, including those which may transmit [West Nile], [Zika], [Yellow Fever], [Dengue], [Eastern Equine Encephalitis], [Western Equine Encephalitis], [Chikungunya], [St. Louis Encephalitis], and [La Crosse Encephalitis] virus(es) and [Dog Heartworm parasites]]
- [Remains active after large rain events]
- [Sumilarv's sand granule formulation stays-put and will not flush from application site]
- [Sumilarv's sand granule formulation is heavy enough to penetrate algae or weed choked sites]
- Flexible dosing options
- For backyard water sources [containers]
- Season long mosquito control [up to 150 days]
- Controls *Aedes*, *Anopheles*, and *Culex* mosquitoes

ACTIVE INGREDIENTS:

Pyriproxyfen	0.50%
OTHER INGREDIENTS	99.50%
	100.00%

Sumilarv® is a registered trademark of Sumitomo Chemical Co., Ltd.

MGK® is a registered trademark of McLaughlin Gormley King Company.

[See inside booklet for Precautionary Statements, Directions for Use and Storage and Disposal]

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to treated, finished drinking water reservoirs or drinking water receptacles when the water is intended for human consumption. Do not apply to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Manufactured by:



7325 Aspen Lane North
Minneapolis MN 55428

EPA Reg. No. 1021-2819

EPA Est. No. 1021-MN-2

Net Weight: _____

ACCEPTED

02/03/2026

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under
EPA Reg. No. 1021-2819

DIRECTIONS FOR USE

It is a violation of Federal law to use this product
in a manner inconsistent with its labeling.

READ ALL DIRECTIONS COMPLETELY BEFORE USE

USE RESTRICTIONS:

- Apply this product only as specified by the EPA-accepted label and any supplemental labeling.
- Do not apply more than 0.1 lbs. pyriproxyfen per acre per year or (20 lbs. of Sumilarv® 0.5G/acre/year).
- Do not apply or allow drift onto pastureland or potable water supplies.
- In the treatment of corral lots, feedlots, swine lots and zoos, cover any exposed drinking water, drinking water fountains, and animal feed before application.
- For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

PESTICIDE RESISTANCE MANAGEMENT

For resistance-management, Sumilarv 0.5G contains a Group 7 insecticide. Any insect population may contain individuals naturally resistant to Sumilarv 0.5G and other Group 7 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed. To reduce the potential for developing insect resistance, rotate to an insecticide with a different mode of action. Monitor treated pest populations for resistance development. Read product label before applying any insecticide and follow label directions.

To delay insecticide resistance, take the following steps:

- Rotate the use of Sumilarv 0.5G or other Group 7 insecticides with different groups that control the same pests. Avoid application of more than the maximum seasonal use rate Sumilarv 0.5G.
- Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance-management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance-management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, record keeping, and which considers cultural, biological and other chemical control practices.

Monitor after application for unexpected target pest survival. Sumilarv 0.5G only targets juvenile insects. If the level of pest emergence suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.

- Contact your local extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance contact MGK Company representatives at 1-800-645-6466.

MOSQUITO GENERA CONTROLLED AND PRODUCT INFORMATION:

Pyriproxyfen effectively controls several mosquito species which can vector Zika virus, West Nile virus, Yellow Fever, Dengue, Eastern equine encephalitis (EEE), Western equine encephalitis (WEE), St. Louis encephalitis (SLE), and dog heartworm in the USA. These include mosquitoes in the following genera:

Aedes
Culex
Anopheles

Introduction of Sumilarv 0.5G into larval habitats where mosquitoes breed and develop will prevent adult mosquito emergence and affects reproductive viability of exposed mosquito eggs and adult female mosquitoes. Frequency of application will depend on three factors: dose rate, water quality (e.g. pollution) and dilution caused by rainfall or removal and replenishment of water within the treatment site. Note: presence of actively swimming larvae after a Sumilarv 0.5G treatment is normal. Sumilarv 0.5G does not act until the pupal stage of a mosquito.

Sumilarv 0.5G is a novel mode of action in Vector control and enhances vector control treatment protocols while providing additional

resistance management solutions to their integrated mosquito management programs.

USE SITES

Temporary or permanent water holding sites, including:

Abandoned swimming pools, animal waste lagoons, catch basins, cesspools, construction site depressions, flooded basements, fountains, gutters, junk yards, landfills, livestock runoff lagoons, manmade depressions, ornamental ponds, salvage yards, septic tanks, sewers, tire dumps, tire tracks, vegetation-choked phosphate pits, vehicle impounds, waste settling ponds, and wastewater impoundments associated with organic pollutants and industrial runoffs.

Natural and artificial water holding containers, including:

Abandoned vehicles, bird baths, buckets, flooded roof tops, flowerpots, hollow trees and tree holes, potted plants, rain barrels, and tires.

APPLICATION TO CATCH BASINS

Individual catch basins are one of the main breeding sites for mosquitoes that can transmit West Nile virus and cause biting nuisance in urban, suburban and rural environments. Sumilarv 0.5G may be applied as a pre-treatment to mosquito breeding catch basins or during the season to catch basins containing mosquito larvae. Once applied, the granules will be unaffected by dry, hot, cold or sunny conditions and will begin working when the catch basin is flooded during rain events. During dry periods, Sumilarv 0.5G remains active and will continue to deliver results once water returns to the catch basin. This non-flushing formulation stays in the catch basin, providing slow and consistent release of the active ingredient evenly within the basin.

For urban or suburban catch basin applications, Sumilarv 0.5G provides up to 21 weeks or 150 days of continuous control with a 75g (2.64 oz.) application, [or up to 4 weeks or 28 days with a 25g (0.88 oz.) application], even when exposed to significant rain events. Simply apply 75g of Sumilarv 0.5G through the grate of the catch basin. Application rates can be adjusted for desired duration of mosquito control according to the table below:

Recommended Application per Catch Basin	Residual Performance Expectations
25 grams (0.88 oz.)	Up to 28 days of control
75 grams (2.64 oz.)	Up to 150 days of control

APPLICATION TO CONTAINERS

Sumilarv 0.5G may be applied to natural and artificial water holding containers. Determine the volume of water in the container to be treated and apply a level scoop of Sumilarv 0.5G according to the rate table below

Treatment site (Volume of Water in Gallons)	Amount of Sumilarv 0.5G (grams)	Amount of Sumilarv 0.5G (ounces)	Number of Scoops of Sumilarv 0.5G (0.25g scoop)
0-5	0.25	0.009	1
5-25	1.25	0.044	5
25-100	5	0.176	20

APPLICATION TO WATER RETENTION or WATER DETENTION PONDS

To determine the volume of water in gallons for a specific treatment site, use the following calculation:

$$\text{Total cu. ft. of target treatment site (ft.}^3\text{)} = \text{surface area (ft.}^2\text{)} \times \text{average depth (ft.)}$$

Note: One cu. ft. contains 7.5 gallons of water

Total water volume = Total cu. ft. of treatment site x 7.5 gallons

Treatment site (Volume of Water in Gallons)	Water Depth Less than or Equal to (ft.)	Amount of Sumilarv 0.5G (grams)	Amount of Sumilarv 0.5G wt. ounces	Lb. Al
0 – 500	1	25	0.88 oz.	0.00028
500 – 1500	2	50	1.75 oz.	0.00056
1500 – 3000	4	75	2.65 oz.	0.00083
3000 – 4500	6	100	3.53 oz.	0.0011
4500 – 6000	8	125	4.41 oz.	0.00138

Treatments should be made every 4-5 weeks for water retention or water detention ponds depending on the volume of water or

frequency of rain events.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container. Store in a cool dry (preferably locked) area that is away from water, other pesticides fertilizers, veterinary supplies, food or feed, in a place that is inaccessible to children and animals.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of empty container in a sanitary landfill, or by other procedures approved by state and local authorities.