



### DIRECTIONS FOR USE

#### COTTON

DABAN 1 must be shaken well before each use. Use DABAN 1 for post-emergence control of certain annual and perennial grasses, weeds and vines such as watergrass, goosegrass, nutgrass, cocklebur and Johnsongrass. DABAN 1 should be used from the time cotton plants are 3 to 4 inches high until first bloom and not before or after this period. Apply DABAN 1 when grasses and weeds are growing and preferably before they are 3 inches high. Best results are obtained when the air temperature is above 70°F.

Add 1 gal. of DABAN 1 to 40 gallons of water and mix thoroughly. This solution should be applied at the rate of 40 gallons per acre of soil surface actually sprayed. When making directed application on the drill, use an amount of spray solution in proportion to the amount of land surface actually being sprayed. Thus, if spraying a band 9 inches wide on either side of the drill for a total band width of 18 inches with rows spaced 40 inches apart, use approximately 13 gallons of spray solution per acre of crop treated. When weed and grass

USDA Reg. No. 12-402

control is desired on the middles, an extra nozzle may be mounted to the side to apply a separate directed spray on these areas as well. Spray nozzles should be positioned to give complete coverage of the weeds and grasses to be controlled without contacting the cotton foliage.

#### GENERAL WEED AND GRASS CONTROL

Use DABAN 1 for post emergence control of certain annual and perennial grasses, weeds and vines such as watergrass, goosegrass, nutgrass, cocklebur and Johnsongrass.

Add 1 to 1½ gallons of DABAN 1 to 40 gallons of water. The unwanted vegetation should be thoroughly covered with spray to the point of runoff. Spot retreatments may be necessary and should be made where regrowth occurs. Any spray equipment which will give good spray coverage may be used with power sprayers being particularly effective for such areas as rights-of-way, drainage ditch banks, fence rows, implement storage yards and similar non-crop areas where control of the grasses and weeds mentioned above is desired.



# DABAN 1

NET CONTENTS 1 GALLON

(DSMA, Herban\* & Surfactant)

For Selective Post-Emergence Weed Control

### INGREDIENT STATEMENT

Active:	
Disodium methanearsonate.....	18.09%
Norea (3-(hexahydro-4(7-methanandian-5-yl)-1,1-dimethyl)urea).....	1.79%
Related reaction products.....	0.41%
Inerts:.....	80.71%
Total.....	100%

Total Arsenic, all in water soluble form, expressed as arsenic trioxide is 5.6 lbs. Contains 2.8 pounds Disodium Methanearsonate Herban\* active ingredient and 1 pound Herban\* per gallon.

### WARNING

**KEEP OUT OF THE REACH OF CHILDREN!**

Harmful if swallowed. Do not get in eyes. Avoid breathing spray mist. Wash hands after using. Avoid contact with skin. Avoid storage near food and food products. **KEEP AWAY FROM CHILDREN AND DOMESTIC ANIMALS.** Keep children and animals off treated areas until washed into soil.

**CHAPMAN CHEMICAL COMPANY**

MEMPHIS TENNESSEE U.S.A.

ANTIDOTE: If taken internally, induce vomiting and CALL A PHYSICIAN AT ONCE.

Do not contaminate water used by wildlife and aquatic life for domestic and irrigation purposes.

DO NOT FEED ANY PART OF TREATED COTTON PLANTS FORAGE OR OTHER TREATED VEGETATION TO LIVESTOCK. DO NOT GRAZE TREATED AREAS.

Applications which result in some of the spray solution getting into the cotton may result in injury to some of the cotton and a subsequent loss of yield.

Do not use on other crops or treated soil within 30 days after application. Do not reuse empty containers. Destroy by burning in a safe place.