


NET CONTENTS GALLONS



Contains Four Pounds of Ethyl Parathion per Gallon

COTTON

11.75 gal

INGREDIENTS

Dilute 2 pints of Ethyl Parathion-4 with 1 1/2 gallons of water. Calibrate spray nozzles so that 12 gallons of diluted spray are applied to one acre of cotton at the spray pressure and speed employed.

For early season control use one spray nozzle per row. Apply at time of first cultivation and again at time of second cultivation.

For mid-season control use two spray nozzles per row and make two to three applications 7 to 10 days apart, starting when squares first form.

For late season control use three spray nozzles per row and make applications whenever insects are noticed in damaging numbers. Make frequent inspections (2 to 4 days apart).

APPLICATION BY AIRPLANES: Dilute 1/2 gallon of Ethyl Parathion-4 with a minimum of 10 gallons or more of water and apply to 2 acres.

ETHYL PARATHION HAS BEEN FOUND TO CONTROL THE FOLLOWING MAJOR COTTON PESTS:

- Brown stink bug
- Green stink bug
- Leaf hoppers
- Whiteflies
- Armyworm
- Garden webworm
- Colorado potato beetle
- Colorado beetle
- Colorado chrysalis
- Colorado pupa
- Colorado larva
- Colorado nymph
- Colorado adult
- Colorado egg
- Colorado larva
- Colorado pupa
- Colorado nymph
- Colorado adult
- Colorado egg

For use only by operators experienced or adequately trained in the application of toxic insecticides, and only to be used on the following systems. Do not get in eyes or on skin. If on skin, wash with plenty of water. If on face, wash with soap and water. If on clothing, wash with soap and water. If on hands, wash with soap and water. If on shoes, wash with soap and water. If on car, wash with soap and water. If on house, wash with soap and water. If on furniture, wash with soap and water. If on food, wash with soap and water. If on dishes, wash with soap and water. If on glassware, wash with soap and water. If on linens, wash with soap and water. If on clothing, wash with soap and water. If on shoes, wash with soap and water. If on car, wash with soap and water. If on house, wash with soap and water. If on furniture, wash with soap and water. If on food, wash with soap and water. If on dishes, wash with soap and water. If on glassware, wash with soap and water. If on linens, wash with soap and water.

ACTIVE INGREDIENTS:

Parathion* 46.31%


Xylene 48.22%

INERT INGREDIENTS: 5.89%

TOTAL 100.00%

* O, O-diethyl O-p-nitrophenyl thiophosphate

DANGER



POISON

SEE SIDE PANELS FOR ADDITIONAL WARNINGS.

KEEP OUT OF REACH OF CHILDREN

POISONOUS BY SKIN CONTACT, INHALATION

OR SWALLOWING

RAPIDLY ABSORBED THROUGH SKIN

CALL A PHYSICIAN IMMEDIATELY

AVOIDANCE

INTERNAL: Give a teaspoonful of salt in a glass of warm water; repeat until vomit fluid is clear. Have patient lie down and keep quiet. EXTERNAL: If on skin, wash thoroughly with soap and water. If on clothing, wash with soap and water.

TOXICITY: Ethyl parathion exerts muscarinic-like effects, i.e., stimulation of the parasympathetic nervous system. Reported use of physostigmine as an antidote, such as atropine to the relief of patient's tremor. Be advised to relieve symptoms. The administration of oxygen is helpful.

If symptoms of poisoning include blurred vision, abnormal constriction of the pupil, do not wait for a reaction but give 2 drops of 1% atropine solution (1 drop per eye). Repeat until patient is satisfactory from the area where poisoning is present. Remove contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of parathion. If parathion gets in eyes, wash immediately with plenty of water for 15 minutes and seek medical attention. If on clothing, wash with soap and water. If on shoes, wash with soap and water. If on car, wash with soap and water. If on house, wash with soap and water. If on furniture, wash with soap and water. If on food, wash with soap and water. If on dishes, wash with soap and water. If on glassware, wash with soap and water. If on linens, wash with soap and water.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. PHYSICIAN'S NOTE: Warning symptoms include weakness, headache, lightness in the chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps. Treatment: Give atropine, preferably by injection, grains 1/100 2 or 3 tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

This product is toxic to fish and wildlife. Do not pollute streams or ponds with water by direct application, cleaning of equipment, or disposal of wastes and containers.

Bury spillage, decontaminate area with alkali solution and flush with water. Burn or bury containers; never reuse. Stay out of smoke or fumes.

Oct. 9, 1968 1022-445

Neither the Manufacturer nor the Seller makes any representation or warranty, expressed or implied, concerning the use of this product. The Manufacturer and Seller shall not be liable for any injury, property damage or loss resulting from the use of this product, except in the event of a defect. The user must read and understand the product label and instructions.