



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 23, 2024

Gail Watson
Registrations Representative
IBC Manufacturing Company
416 East Brooks Road
Memphis, TN 38109

Subject: Registration Review Label Mitigation for Permethrin
Product Name: Bee Gone
EPA Registration Number: 1022-586
Application Date: 3/18/2021
Decision Number: 572321

Dear Gail Watson:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Permethrin Interim Decision and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently

approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Carolyn Smith by phone at (202) 566-2273, or via email at smith.carolyn@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Jaclyn Pyne". The signature is written in a cursive, flowing style.

Jaclyn Pyne, Team Leader
Risk Management and Implementation Branch 3
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure

BEE Gone[®]

CONCENTRATE
FOR OUTDOOR USE ONLY


(Control Statement)

(Picture of Insects)

Active Ingredient
 Permethrin* 36.8%
Other Ingredients** 63.2%
TOTAL..... 100.0%

*cis/trans ratio: Max. 42% (\pm) cis and min. 58% (\pm) trans

** Contains petroleum distillates



KEEP OUT OF REACH OF CHILDREN

CAUTION See back panel booklet for Additional Precautionary Statements

EPA Reg. No. 1022-586
EPA Est. No. 1022-TN-001

Net Contents: 8 ounces

BEE Gone[®] is a registered trademark of ISK Americas Incorporated.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Failure to follow the directions for use and precautions on this label may result in poor pest control or plant injury.

For use as a surface spray. Do not allow adults, children, or pets to enter the treated area until sprays have dried. Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift.

USE INFORMATION

This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding, in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.

This product may be used to control insect pests outdoors, on ornamentals, lawns, trees, shrubs, and vines in landscape areas, as a perimeter treatment around residential buildings. BEE Gone may also be used to control home-invading insects as a residual spray on outside surfaces, such as walks, garbage areas, window frames, doorways, porches, and patios. Not for use on plants being grown for commercial sale, or plants grown for seed production or sod farms and golf course turf, or in greenhouses or nurseries.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: 1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation; 2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g. soil, mulch, or lawn); 3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure; 4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch; 5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level; 6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.

Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application. Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment. Treat surfaces to ensure thorough coverage but avoid runoff. Do not apply or irrigate to the point of runoff. To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

Not for use in outdoor residential misting systems. Do not water the treated area to the point of runoff. Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Application Instructions: BEE Gone mixes readily with water and is specifically packaged for residential use as a diluted spray. Use a clean sprayer, such as a tank type or hose-end sprayer to mix and apply the proper amount of concentrate with the desired volume of water. For perimeter and lawn applications, measure the area to be treated to determine the application volume. For trees, shrubs and flowers, apply as a thorough cover spray until foliage is wet. Do not apply when the wind speed is greater than 15 mph. Refer to Specific Application Instructions in the recommended rate tables for additional instructions.

HOME-INVADING PESTS (OUTDOOR USE ONLY)

MIXING INSTRUCTIONS

Mix 1.6 fluid oz. (3 ¼ Tablespoons) of BEE Gone Concentrate per 1 gallon of spray solution to make a 0.5% spray emulsion [1 fluid oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)]. Do not use utensils such as spoons and measuring cups after use with pesticides.

HOW TO APPLY

Spot Treatment

- Apply a 0.5% emulsion as a pin stream low-pressure spray or as a paint-on application with a paintbrush.
- Treat where pests are found or entry points of the structure such as window frames, doorways, porches, patios, and foundation.
- Spot-treatment may also be made to garages, garbage cans, and garbage areas.
- DO NOT apply indoors (including crawl spaces).

Perimeter Application

- Spray a 6 to 8 foot wide band of soil around the perimeter of the house. Treat the foundation wall to a height of 2 to 3 feet.
- Use a spray volume of 2 to 10 gallons of 0.5% emulsion per 1000 square feet. If mulch or debris is present, a higher volume application rate may be needed to ensure adequate coverage.
- Note: residual control will vary depending on exposure of the treated area to weathering conditions (temperature, sunlight, moisture). Retreat on a monthly schedule while insects are likely to be active.

Wood Destroying Pests

- Localized Areas Outdoors: for the control of insects such as termites, bees, carpenter ants and wood infesting beetles in localized above-ground areas of infested wood around structures, apply a 0.5% emulsion to voids and galleries in damaged wood and in spaces between wooden members of a structure and between wood and foundations where wood is vulnerable. Apply either as a spray or paint-on application. For termites, the purpose of application is to destroy workers or winged reproductive forms, which may be present at the time of treatment. This product is not designed for preventative use and should only be applied when pests appear. This application is not a substitute for soil or foundation treatment.
- Use a hose-end sprayer or sprinkling can to deliver a coarse drenching spray to treat stored lumber or wood piles. Firewood is not to be treated. Stored wood can be used for lumber one month after treatment. Do not apply inside structures.
- Repeat treatment as necessary but not more frequently than once a month. For active termite infestation, consult a professional Pest Control Applicator.

HOME-INVADING PESTS (OUTDOOR USE ONLY)	
PEST	SPECIFIC APPLICATION INSTRUCTIONS
Ants (including mounds) Armyworm Bark Beetles Bees Beetles Borers Boxelder Bugs Carpenter Ants Carpenter Bees Centipedes Chinchbugs Cockroaches Cockroaches (Asian) Crickets Earwigs Elm Leaf Fire Ants (Incl. mounds) Fleas Flies Firebrats Ground Beetles Gypsy Moths (adults & caterpillars) Millipedes Mole Crickets Pill Bugs Scorpions Silverfish Sod Webworm Sowbugs Spiders Ticks (including Deer Tick, Western Black-legged Tick known to transport Lyme disease and Rocky Mountain Spotted Fever) Wasps	<p>Boxelder Bugs; Elm Leaf Beetles and Gypsy Moth Caterpillars: Spray tree trunks, building siding, or wherever pests congregate. Avoid excessive application. Dampen surface thoroughly, but do not let runoff occur.</p> <p>Borers and Bark Beetles: To prevent infestation of ornamental trees and woody shrubs, spray the bark to the point of saturation, but do not let runoff occur. Repeat treatment every 21 days during early spring until mid-summer.</p> <p>Carpenter Bees: Apply as a pin stream, as a fine/coarse low pressure spray (20 psi or less) as a spot treatment.</p> <p>Fire Ants (mound drench): Apply 1 or 2 gallons of emulsion to each mound area by sprinkling the mound until it is wet and treat a 4-foot diameter circle around the mound. Use higher volume for mounds larger than 12 inches. For best results, apply in cool weather, such as in early morning or late evening hours, but not in the heat of the day.</p> <p>Scorpions: Remove accumulations of lumber, firewood, and other materials which serve as harborage sites and treat beneath and adjacent to infested sites.</p>

LAWN PESTS

BEE Gone can be applied to lawns adjacent to or around private homes, duplexes, townhouses, condominiums, house trailers, apartment complexes, carports, garages, fence lines, storage sheds, barns, and other residential areas where pests congregate.

MIXING INSTRUCTIONS

Prepare an emulsion by mixing 0.4 to 0.8 fluid oz. (2 ½ to 5 teaspoons) of BEE Gone Concentrate in sufficient volume of water to cover 1000 sq. ft. (approximately 4 to 25 gallons of water) [1 fluid oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)]. Do not use utensils such as spoons and measuring cups for food purposes after use with pesticides.

HOW TO APPLY

Mound Drench Treatment (Ant and Fire Ant Mounds in Lawns)

Use a sprinkling can. Mix 1.6 fluid oz. (3 ¼ Tablespoons) of BEE Gone Concentrate per 1 gallon of spray solution to make a 0.5% spray emulsion [1 fluid oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)]. For large ant mounds, such as Fire Ant mounds, apply 1 – 2 gallons of a 0.5% emulsion to each mound area including a 4 foot diameter circle around the mound. Apply to smaller mounds until the mound is completely saturated. Use higher water volume for mounds larger than 12”.

Lawn Broadcast Application

- Apply using a hose-end or pressure type sprayer to deliver a coarse spray of between 4 to 25 gallons of emulsion per 1,000 sq. ft. of lawn.
- Use the higher recommended label rate for heavy pest infestations.
- Use sufficient volume to uniformly cover plant surface.

Note: Higher volume of water should be used when situations of thick thatch or dense sod exist, or in arid or drought conditions.

WHEN TO APPLY

- For best results, treat in early morning or late evening hours, but not in the heat of the day.
- Spray evenly over the lawn when insects first appear.
- Repeat applications as necessary but not more frequently than every 10 – 14 days. Do not apply more than 1.6 fl. oz. / 1000 sq. ft. per year.

RECOMMENDED APPLICATION RATES FOR LAWN			
SITE	PEST	RATE	SPECIFIC APPLICATION INSTRUCTIONS
Lawns, residential including but not limited to St. Augustine, Bentgrass, Fescues, Buffalograss, Bahiagrass, Bluegrass, Perennial Ryegrass, and Centipedegrass	Ants Armyworms Bilbugs Chiggers Chinchbugs Crickets Cutworms Earwigs Fruit Flies Fleas Leafhoppers Millipedes Mole Crickets Sod Webworm Sowbugs Ticks (including Brown Dog Tick and Deer Tick, Western Black-legged Ticks known to transport Lyme disease and Rocky Mountain Spotted Fever) For additional pests consult the list of pests under Outdoor Pests Controlled by BEE Gone.	0.4 to 0.8 Fluid Ounces (2 ½ to 5 tsp) per 1000 sq. ft.	Apply using 4 – 25 gallons of spray volume / 1000 sq. ft. of lawn. Use higher volumes on St. Augustine, Buffalograss, Bermuda and Bahiagrass lawns. Use the higher rate for fast knockdown and increased residual control. Dense and excessive (greater than 3 inches) lawn height and arid conditions may require higher volume application rates. Repeat applications may be necessary (10 to 14 day interval) to control heavy infestations during peak periods of insect feeding. Do not exceed a total of 1.6 fl. oz. / 1000 sq. ft. per year (i.e. four (4) applications at the lower rate or two (2) applications at the higher rate.

ORNAMENTAL PESTS

MIXING INSTRUCTIONS

Mix 0.04 to 0.08 fluid oz. (=1/8 to ¼ teaspoons = 1-2 ml) of BEE Gone Concentrate in 1 gallon of water for general pest control. Refer to the Ornamental Application Rate Table for additional mixing instructions and for specific directions for stem-infesting beetles. (1 fluid oz. = 2 Tablespoons (T) = 6 teaspoons (tsp) = 29.6 ml]. Do not use utensils such as spoons and measuring cups for food purposes after use with pesticides.

HOW TO APPLY

Ornamentals (Foliage and Flowering Plants)

- Apply using a hose-end or pressure type sprayer to deliver a coarse spray. Apply thoroughly to stem and foliage.
- Use the higher recommended label rate (1/4 tsp. per gal. of water) for heavy pest infestations. Note: Higher volumes of water should be used when situations of dense foliage exists, or in arid or drought conditions.

WHEN TO APPLY

- For best results treat in early morning or late evening hours, but not in the heat of the day.
- Spray evenly over the lawn when insects first appear.
- To prevent insect reinfestation repeat treatment as necessary but not more frequently than once every 14 days.

RECOMMENDED APPLICATION RATES FOR USE IN ORNAMENTALS

SITE	PEST	RATE	SPECIFIC APPLICATION INSTRUCTIONS
Ornamental foliage and flowering plants (including Conifers, woody and herbaceous non-edible ornamentals.	<p>Ants</p> <p>Aphids</p> <p>Bagworm</p> <p>Beet Armyworm</p> <p>Cabbage Looper</p> <p>Cankerworms</p> <p>Citrus Thrips</p> <p>Coneworm (incl. web)¹</p> <p>and Seed bugs</p> <p>Fungus Gnat</p> <p>Gypsy Moth</p> <p>Caterpillars</p> <p>Heliothis spp</p> <p>Japanese Beetles</p> <p>Lace Bug</p> <p>Leaf Feeding</p> <p>Caterpillars</p> <p>Leaffooted Bugs</p> <p>Leafminers</p> <p>Leafhoppers</p> <p>Leaf rollers</p> <p>Lygus Bugs</p> <p>Mealybugs</p> <p>Mosquitoes</p> <p>Nantucket Pine Tip</p> <p>Moth</p> <p>Oblique Leafroller</p> <p>Pine Sawflies</p> <p>Plant Bugs</p> <p>Root Weevils (Adult)</p> <p>Rosy Aphid</p> <p>Rose Chafer</p> <p>Spotted Leafminer</p> <p>Stink Bugs</p> <p>Tarnished Plant Bug</p> <p>Tent Caterpillars</p> <p>Webworms</p> <p>Whiteflies</p> <p>Zimmerman Pine</p> <p>Moth¹</p>	<p>0.04 to 0.08</p> <p>Fluid Ounces</p> <p>(1/8 to ¼ tsp.)</p> <p>(1 to 2 ml)</p> <p>Per gal.</p>	<p>Note: Direct application to blooms may cause browning of petals. Marginal leaf burn may occur on Salvia, Dieffenbachia and Pteris Fern</p> <p>Mosquitoes coming to rest on treated plants will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.</p> <p>¹To control Webbing Coneworm make first application just prior to peak pollen flight. To control other Coneworms and Seed Bugs, make application 30 days prior to flower closure.</p>
	<p>Bark Beetles and Boring</p> <p>Insects, (including but</p> <p>not limited to: Ash</p> <p>Borer, Bronze Birch</p> <p>Borer, Rhododendron</p> <p>Borer, Elm Bark Beetles</p> <p>and Turpentine Beetles</p>	<p>0.3 to 0.6 fluid</p> <p>ounces</p> <p>(2 to 4 tsp)</p> <p>per gal.</p>	<p>Stem-Infesting Beetles: Treat lower branches and trunk prior to adult emergency. Complete, uniform coverage is needed for best control. A repeat application may be necessary on a 3 to 4 week interval to prevent severe tree stem damage.</p>

IMPORTANT

DO NOT use on vegetables or food crops. DO NOT treat pets with this product. DO NOT spray inside dwellings. DO NOT treat soil that is water saturated or frozen or in any conditions where run-off or movement from the treatment area (site) is likely to occur. DO NOT apply when wind velocity favors drift.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep out of the reach of children or animals. Store in original containers only. Store in a cool, dry place, preferably in a locked storage area and avoid excessive heat. Carefully open containers. After partial use, replace lid and close tightly. Do not put concentrate or dilute material into food or drink containers.

PESTICIDE DISPOSAL: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available.

If empty: Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Never pour unused product on the ground. Do not pour or dispose down-the-drain or sewer. Call your local solid waste Agency for local disposal options.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID:	
IF SWALLOWED:	<ul style="list-style-type: none"> ▪ Immediately call a poison control center or doctor. ▪ Do not induce vomiting unless told to do so by a poison control center or doctor. ▪ Do not give any liquid to the person. ▪ Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 – 20 minutes. ▪ Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p> <p>You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.</p> <p>NOTE TO PHYSICIAN: This product contains petroleum distillates – vomiting may cause aspiration pneumonia. Consideration should be given to gastric lavage with an endotracheal tube in place. Treatment is controlled removal of exposure followed by symptomatic and supportive care.</p>	

ENVIRONMENTAL HAZARDS:


This product is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

PHYSICAL OR CHEMICAL HAZARDS: Combustible. Do not use or store near heat or open flame.

CONTAINER USE DIRECTIONS:



1. Remove the measuring chamber cap and induction seal. Replace cap and securely tighten. Tip container until liquid fills measuring chamber.
2. Return container to level position. No adjustment is needed.
3. Removing measuring chamber cap and dispense into proper application equipment.

NOTICE: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

Manufactured by:

IBC Manufacturing Company
416 East Brooks Road
Memphis, TN 38109
901-344-5350

List of User Friendly Statements to be added to Final Print Label:

Brand Extensions

HG
H&G
Lawn
Outdoor

Use Statements

Insecticide
Insecticide / Termiticide
Termiticide
Lawn & Ornamental
Soil Insect & Termite Killer
Fire Ant Killer
Ant & Roach Killer
Outdoor Perimeter
For Outdoor Residential Use Only

Control Statements

Kills Ticks that may cause Lyme disease
Kills Home Invading Pests (Outdoor)
Kills Named Insects
An easy to apply product to control Home Invading and Lawn Insects
An easy to apply product to control wood invading insects
Termiticide Formulation for Homeowner Use (Outdoor Use Only)
Termite Insecticide (Outdoor Use Only)
For Homeowner Use (Outdoor Use Only)
Controls Wood-Destroying Insects

Icons / Pictures

(Icon of People)
(Icon of Sprinkling Can)
(Icon of Hand Held Sprayer)
(Tree Icon)
(Plant Icon)
UPC Symbology
Picture Fire Ant Mound
Picture of a residential home
Picture of a lawn
Pictures of one or more of the following insects: Termites, ants (including Fire Ant mounds and Carpenter Ants), cockroach, spider, scorpion, fleas, tick, mosquito

Other

Treats up to (#will depend on application) Square Feet
Made in the USA
Formulated and Packaged in the USA
How to Apply
When to Apply
Amount to Use