

TUCO

Net Weight 10 pounds
9-630-1

Acti-dione® RZ

Broad Spectrum Turf Fungicide

Active Ingredients
Cycloheximide 13% w/w
(3-[2-(3,5-dimethyl-2-oxocyclohexyl)-2-hydroxyethyl]-glutarimide) 75.0% w/w
Inert Ingredients 23.7% w/w
100.00% w/w

For Use On:
Bluegrass, Bentgrass, Bermudagrass, Fescue, Ryegrass, and St. Augustinegrass
For the Prevention and Control of:
Large Brown Patch, Warming-Out, Rust, Dollarspot, Fading Out, Grease Spot, Jersey Wilting, Leafspot (*HelmiaBergerii*) and Gray Leafspot
Also for Aerial Petal Blight

KEEP OUT OF REACH OF CHILDREN

WARNING

MAY BE FATAL IF SWALLOWED

(See Side Panel for further Precautionary Statements)

811 231 001 EPA Reg. No. 1023-10
USE BEFORE EPA Est. No.
LOT

Packed for
TUCO - DIVISION OF THE UPJOHN COMPANY
KALAMAZOO, MICHIGAN 49001 U.S.A.

E229P
EDP 696717

ACCEPTED
SEP 16 1983
Made the Federal Insecticide
Fungicide and Rodenticide Act
1033-10

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Do not breathe dust. May be irritating if contacted with skin, eyes or clothing. Avoid contact with skin, eyes or clothing.

Statement of Practical Treatment
IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES OR ON SKIN: In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply where weather or strong winds drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Acti-dione RZ 10 pounds

TUCO

GENERAL INFORMATION

Read all label directions and warnings before using Acti-dione RZ. It is a broad spectrum turf fungicide designed for the prevention and control of most of the common fungus diseases on the major turfgrasses. It is effective for use on all turfgrasses. Do not use on or around the home.

SELECTIVE FOR USE

It is a selective fungicide for use on turfgrasses in a manner suggested with its labeling.

USE

Apply as a liquid spray or a hand or power sprayer, or a broadcast granular application. Use the recommended amount of Acti-dione RZ. It is a selective fungicide for use on turfgrasses. Do not use on or around the home. Do not use on or around the home.

For Control of: Large Brown Patch, Warming Out, Fading Out, Grease Spot, Leafspot, and Gray Leafspot.

Preventive Treatment: Apply Acti-dione RZ early in the spring before disease symptoms appear. Then follow a regular schedule of 2 to 4 applications throughout the season when conditions are favorable for disease development. Use 4 ounces of Acti-dione RZ in 25 to 35 gallons of water per 1,000 sq. ft. of turf area.

Curative Treatment: When disease is a ready problem or disease develops a preventive spray program, use 4 ounces of Acti-dione RZ in 25 to 35 gallons of water per 1,000 sq. ft. of turf area. Repeat treatment at 7 to 14 day intervals until the disease is controlled.

For Control of: Rust, Fungus Blight, Fading Out, Warming Out, and Leafspot (HelmiaBergerii), and Gray Leafspot.

Preventive Treatment: Apply Acti-dione RZ early in the spring before disease symptoms appear. Then follow a regular schedule of 2 to 4 applications throughout the season when conditions are favorable for disease development. Use 4 ounces of Acti-dione RZ in 25 to 35 gallons of water per 1,000 sq. ft. of turf area.

Curative Treatment: When disease is a ready problem or disease develops a preventive spray program, use 4 ounces of Acti-dione RZ in 25 to 35 gallons of water per 1,000 sq. ft. of turf area. Repeat treatment at 7 to 14 day intervals until the disease is controlled.

COMBINATION: Acti-dione RZ may be applied in combination with other fungicides. Fungicide combinations may be used to the same extent as the rate of 4 ounces per 1,000 sq. ft. of turf area. Do not mix Acti-dione RZ with other fungicides or with other chemicals such as insecticides, herbicides, and fertilizers.

SALES BY THE GALLON: When application is made by hand or power sprayer, use 4 ounces of Acti-dione RZ in 25 to 35 gallons of water per 1,000 sq. ft. of turf area. Do not mix Acti-dione RZ with other fungicides or with other chemicals such as insecticides, herbicides, and fertilizers.

When to Apply	Use this Amount of Acti-dione RZ	Frequency
4 to 6 weeks before disease (for application only)	One (1) ounce per square yard of the surface	Single spray or 4 to 6 times as directed
Pre-season and Season (Repeat applications as for preventive treatment)	1 to 4 ounces per 1,000 sq. ft. of turf area	Repeat at 7 to 14 day intervals or as new disease occurs

When using backpack sprayers, reduce the concentration to 1/4 of a level teaspoon per gallon or 8 ounces per 100 gallons and use 2 to 4 times as above. Do not use Acti-dione RZ on or around the home. Do not use on or around the home.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container and keep closed. Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide spray should be disposed of in accordance with Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Unions of bags according to approved Federal, State and local procedures under the Resource Conservation and Recovery Act. Do not reuse container.

IMPORTANT—READ BEFORE USE

By using this product user accepts the following conditions, warranty, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, because of extreme weather and soil conditions, neither of use and other factors beyond TUCO's control, it is impossible for TUCO to eliminate all risks associated with use of this product. As a result, crop injury or ineffectiveness is always possible.

WARRANTY AND DISCLAIMER OF WARRANTIES: TUCO warrants that this material conforms to the chemical description on the label and is reasonably fit for use under the directions and conditions on this label. TUCO AND SELLER MAKE NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS, MERCHANTABILITY OR OTHERWISE.

LIMITATIONS OF LIABILITY: The liability of TUCO or Seller for damages arising from the use of this product is limited to the replacement cost of the product used and shall not include any consequential damages such as loss of profits or other.

NO CHARGES AUTHORIZED: No one (other than an authorized agent of TUCO) is authorized to make any other warranty, or change the above conditions, disclaimer or limitations and then only if in writing and with a specific reference to this label.

REST DOCUMENT AVAILABLE