1023-51	ACCEPTED 5/19/1975 HUBILION	Page 172
9-533-1	MAY 1 9 1975 Net Weight 2 Pounds AR 18 1975	No. 9597
	Under the Folder State Add PROXOL® 80 SP Insecticide EPA Reg. No. /023-57	e •

Active Ingredient

WARNING: Keep out of reach of children.

May be fatal if swallowed. Harmful if inhaled, or absorbed through the skin. Do not get in eyes or on skin. Avoid breathing mist or dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Do not contaminate food or feed. Keep bag closed. Use with adequate ventilation.

FIRST AID: Exposure will result in cholinesterase depression. If swallowed, induce vomiting. In case of contact, remove contaminated clothing. Wash skin with soap and water; for eyes, flush with water for at least 15 minutes. Obtain prompt medical aid.

ANTIDOTE: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

PRECAUTION: Keep children and pets off treated areas until spray has dried. Do not use on other crops used for food or forage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse. This product is toxic to bees and should not be applied when bees are actively visiting the area.

WILDLIFE: This product is toxic to birds and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONTAINER & WASTE DISPOSAL: Rinse equipment and dispose of wastes by burying in noncrop land away from water supplies. Dispose of empty container by burying with wastes or by burning (Keep out of smoke).


COMPATIBILITY: Proxol 80 SP may be applied with most commonly used insecticides. Do not mix with alkaline materials such as lime and lime sulfur.

EPA Reg. No. 1023-51-AA

Manufactured for TUCO, Division of The Upjohn Company • Kalamazoo, Michigan 49001, U.S.A.

> Page 1 of 11 (<u>Revised 3-18-75</u>)

PROXOL 80 SP



## RECOMMENDED APPLICATIONS

CROP	INSECT	Ounces PROXOL 80 SP	REMARKS	
FRUIT	Fruit tree leafroller Omnivorous leafroller		Apply specified dosage per acre in ground equipment using 200 to 500 gallons of water for good coverage. Up to three applications may be made per season. Do not apply within 21 days of harvest.	
Citrus (California only)	Avocado leafroller Orange Tortrix Citrus cutworm Citrus looper	80		
VEGETABLES Western bean cutworm 10 to 20		Apply specified dosage per acre		
Beans (dry & snap)	Armyworms Dipterous leaf miners Lygus bugs Mexican bean beetle Stink bugs Variegated cutworm	20 to 30	using sufficient water for com- plete coverage but not less than l gallon per acre. Repeat as necessary. Do not apply within 14 days of harvest.	
Lima beans	Western bean cutworm	10 to 20	Apply specified dosage per acre using sufficient water for com- plete coverage but not less than l gallon per acre. Up to five applications may be made per season. Do not apply within 3 days of harvest.	
	Armyworms Dipterous leaf miners Lygus bugs Mexican bean beetle Stink bugs Variegated cutworm	20 to 30		
Cowpeas (Southern peas, Black- eyed peas, Crowder peas)	Armyworms Dipterous leaf miners Imported cabbageworm Lygus bugs Mexican bean beetle Stink bugs Variegated cutworm	20 to 30	Apply specified dosage per acre using sufficient water for com- plete coverage but not less that l gallon per acre. Do not appl more than twice per season nor within 14 days of harvest.	
Brussels sprouts	<b>y</b>		Apply specified dosage per acre using sufficient water for com-	
Cabbage Cauliflower	Imported cabbageworm Variegated cutworm	10 to 20	plete coverage but not less that l gallon per acre. Repeat as necessary. Do not apply within 21 days of harvest.	

Page 6 of 11 (<u>Revised 3-18-75</u>)

1