

**WARNING**

HARMFUL IF SWALLOWED. Contact with skin can cause toxic symptoms. In case of contact with skin, wash with soap and water. Avoid breathing spray mist. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Do not use Chlordane on grazing areas or on crops after edible parts start to form. Do not feed treated plants to poultry, dairy cattle or animals being finished for slaughter. Do not use in dairy barns. Do not use on forage. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Do not contaminate feed and foodstuffs. During commercial or prolonged exposure in mixing and loading operations, wear clean, synthetic rubber gloves and a mask or respirator of a type found suitable by the U. S. Bureau of Mines for protection against Chlordane. Also see cautions under directions for specific uses.

**ENVIRONMENTAL CAUTIONS**

**TOXIC TO FISH AND WILDLIFE**—Birds feeding on treated areas may be killed. Do not contaminate any body of water with this material. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees. Don't apply when they are actively visiting the area. Apply this product only as specified on label. Do not apply when weather conditions favor drift from areas treated.

**CONTAINER DISPOSAL**

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place. Large drums should be returned to drum reconditioner.

**DIRECTIONS FOR USE**

- Chlordane 4E can be used in conventional hydraulic sprayers and hand sprayers. It emulsifies readily with water. Agitate the mixture before use and while spraying.

**Emulsible Insecticide**

<b>ACTIVE INGREDIENTS:</b>	<b>By Weight</b>
*Technical Chlordane . . . . .	45%
Petroleum Distillate . . . . .	51%
<b>INERT INGREDIENTS . . . . .</b>	<b>4%</b>
<b>TOTAL . . . . .</b>	<b>100%</b>

\*Equivalent to 27.0% Octachloro 4,7-Methanotetrahydroindane and 18.0% Related Compounds.

**Contains 4 lbs. Technical Chlordane per Gallon**

**CAUTION  
KEEP OUT OF REACH OF CHILDREN**

See Additional Cautionary Information On Side Panels.  
Study Entire Label Before Opening or Handling.

**NOTICE**

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME**

**ACCEPTED**  
 JUN 4 1976  
 Under the Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 161-101

**SOIL TREATMENTS:** Follow state and federal recommendations.

Never use less than 20 volumes water per volume of concentrate.

**Broadcast:** Apply before planting. Mix the required amounts of Concentrate in sufficient water for uniform coverage of soil surface. Till promptly to mix the chemical into top 3 to 6 inches of soil.

**Row, Band or Furrow:** Mix the required amounts of Concentrate in sufficient water for uniform coverage of row when planting or seed furrow before closing.

Corn (incl. sweet corn), Sorghum, Soybeans

White Fringed Beetle Larvae: For use only in 8 Southeast States (AL, FL, GA, LA, MS, NC, SC, TN)	White Grubs: For use only in the State of Michigan
---	---

Broadcast 1 1/4 gallons per acre or make row or furrow treatment of 1 to 2 quarts per acre.  
Note: Use on Corn is suspended effective 8/1/76.

**SUBTERRANEAN TERMITES** — Directions for Non-Professional Use:

Prepare a 1% Chlordane water emulsion by mixing one volume of Chlordane-4E with 47 volumes of water, and use as follows:

**New Construction:** Apply 1 gallon per 10 sq. ft. to the soil surface before laying slabs which are adjacent to the building.

**Buildings Having Basements or Crawl Spaces:** Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundations, dig a trench to, but not below, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment. It is suggested that you consult the State Extension Service for advice.

4/26, 76)

Contents	EPA Est. 1029-1A-1	MANUFACTURED BY	<b>AIDEX</b>	OMAHA, NEBRASKA	EPA Reg. No.	LOT
----------	--------------------	-----------------	--------------	-----------------	--------------	-----