

## DIRECTIONS

Apply at the rate of 10 to 25 pounds per acre using a standard aerial or ground granular application equipment.

**For control of EUROPEAN CORN BORER (SWEET CORN)**—Make first application when 25% of plants show evidence of leaf feeding and continue at 7 day intervals as long as leaf feeding is evident. From one to three treatments may be required. For control of second brood borer, apply when eggs begin to hatch.

**FIELD CORN**—Apply 10 days after eggs first hatch or when 75% of plants show evidence of feeding in leaf whorls. Repeat in 7 to 10 days if new evidence of feeding is present. For control of second brood borer, apply when eggs begin to hatch.

**NOTE: NEVER APPLY EPN WITHIN 14 DAYS OF HARVEST.**

**EXTREMELY HAZARDOUS BY SKIN CONTACT, INHALATION OR SWALLOWING. RAPIDLY ABSORBED THROUGH SKIN. REPEATED EXPOSURE MAY, WITHOUT SYMPTOMS, BE INCREASINGLY HAZARDOUS.**

Do not get on skin, eyes, or clothing. Do not breathe vapor or loose dust. Wear clean, heavy, natural rubber gloves, rubber boots, and clean, waterproof or freshly laundered clothing, coveralls, cap, and goggles when applying or handling. Destroy and replace gloves frequently. Wear an effective mask or respirator of a type passed by the U.S. Bureau of Mines for protection.

After applying or handling, wash hands, arms, and face thoroughly with soap and water before eating or smoking. After work, bathe at home and change all clothing. Wash clothing with soap and hot water thoroughly before reuse.

In case of any contact or a possible case of poison symptoms, apply FIRST AID immediately. Follow directions on page

## NOTICE

# AIDEX EPN-2G

## 2% EPN GRANULAR INSECTICIDE

For Control of European Corn Borer

For control of European Corn Borer (SWEET CORN) and Field Corn. Apply at the rate of 10 to 25 pounds per acre using a standard aerial or ground granular application equipment.

## WARNING

Keep animals and unprotected humans away from area during treatment and when there is a danger from drift or residues.

**THIS PRODUCT IS TOXIC TO FISH, BIRDS, AND OTHER WILDLIFE.** Do not contaminate any body of water or any spot specified on label with this material or its spillage. Do not use cleaning equipment to dispose of waste in any other manner.

Apply this product only as specified on the label.

Do not apply when weather conditions favor drift from area being treated. Do not apply where runoff is likely to occur.

Store in cool, dry place, away from feed, foodstuff, and domestic animals. Keep container closed.

Do not contaminate Feed or Foodstuff.

Do not retain or reuse empty container.

Crush and bury when empty or return drum to rental conditioner. Bury any spillage and clean up area with strong lye solution.

**CAUTION**—This product is highly toxic to bees, exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

## Symptoms of Poisoning

Symptoms of poisoning by any organophosphorus insecticide, including EPN, are likely to be as follows:

- 1. Digestive upset, nausea, vomiting, and diarrhea.
  - 2. Pressure under breastbone, diarrhea.
  - 3. Swelling and excessive salivation.
  - 4. Headache, dizziness, headache.
  - 5. Muscle tremors, twitching, and convulsions.
  - 6. Weakness, loss of coordination, and loss of reflexes.
  - 7. In severe cases, respiratory paralysis, and death.
- Do not allow contact with any organophosphorus insecticide, including EPN, on your skin, clothing, or hair.



## DIRECTIONS

Apply at the rate of 10 to 25 pounds per acre using standard aerial or ground granular application equipment.

**For control of EUROPEAN CORN BORER: SWEET CORN** - Make first application when 25% of plants show evidence of leaf feeding and continue at 5-day intervals as long as leaf feeding is evident. From one to three treatments may be required. For control of second brood borers, apply when eggs begin to hatch.

**FIELD CORN** - Apply 10 days after eggs first hatch or when 75% of plants show evidence of feeding in leaf whorls. Repeat in 7 to 10 days if new evidence of feeding is present. For control of second brood borers, apply when eggs begin to hatch.

**NOTE: NEVER APPLY EPN WITHIN 14 DAYS OF HARVEST.**

## DANGER!

EXTREMELY HAZARDOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING. RAPIDLY ABSORBED THROUGH SKIN. REPEATED EXPOSURE MAY, WITHOUT SYMPTOMS, BE INCREASINGLY HAZARDOUS.

Do not get on skin, eyes, or clothing. Do not breathe vapor or loose dust. Wear clean, heavy, natural rubber gloves, rubber boots, and clean, waterproof or freshly laundered clothing (coveralls, cap), and goggles when applying or handling. Destroy and replace gloves frequently. Wear an effective mask or respirator of a type passed by the U.S. Bureau of Mines for protection.

After applying or handling wash hands, arms, and face thoroughly with soap and water before eating or smoking. After work bathe at once and change all clothing. Wash clothing with soap and hot water thoroughly before reuse.

In case of **any contact** or appearance of **poison symptoms** apply **FIRST AID** measures shown on front panel.

## NOTICE

See the quantity of material listed on the front of the label and apply the rate of application accordingly. Do not use more than the amount specified on the label. Do not use the product if the label is missing or if the label is illegible.

# AIDEX EPN-2G

## 2% EPN GRANULAR INSECTICIDE

For Control of European Corn Borer

### ACTIVE INGREDIENTS

O-Ethyl O-p-nitrophenyl phenylphosphonothioate 2

INERT INGREDIENTS 98

From technical EPN

Total 100

DANGER



POISON

KEEP OUT OF REACH OF CHILDREN

## ANTIDOTE AND FIRST AID

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON

Internal

External

Note to obtaining antidote

## WARNING

Keep animals and unprotected humans away from area during treatment and while there is danger from drift or residues.

THIS PRODUCT IS TOXIC TO FISH, BIRDS AND OTHER WILDLIFE. Do not contaminate any body of water or area not specified on label with this material, by spillage, drift, cleaning equipment, disposal of wastes, or in any other manner.

Apply this product only as specified on this label.

Do not apply when weather conditions favor drift from area being treated. Do not apply where runoff is likely to occur.

Store in cool, dry place, away from Feed, Foodstuffs, and domestic animals. Keep container closed.

Do not contaminate Feed or Foodstuffs.

Do not retain or reuse empty container.

Crush and bury when empty or return drums to reconditioner. Bury any spillage and clean up area with strong lye solution.

CAUTION: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

## Symptoms of Poisoning

Symptoms of poisoning by any organophosphorus insecticides, including EPN, are likely to be as follows:

- 1 - Digestive upsets: nausea, vomiting, cramps, feeling of pressure under breastbone, diarrhea.
- 2 - Sweating and excessive salivation.
- 3 - Restlessness, giddiness, headache.
- 4 - Muscular tremors, twitching of facial and eye muscles.

If any of these symptoms are felt remove victim from area and apply First Aid measures. Do not allow victim to resume operations with any organophosphorus insecticide until physician has cleared him after testing blood for organophosphorus effect.

