

10% TOXAPHENE GRANULAR INSECTICIDE

AN INSECTICIDE CONCENTRATE ESPECIALLY PREPARED FOR
EUROPEAN CORN BORER CONTROL.

ACTIVE INGREDIENT:	Percent by Weight
Toxaphene (technical chlorinated camphene containing 67-69% chlorine)	10
INERT INGREDIENTS:	90
TOTAL	100

DIRECTIONS FOR USE

European Corn Borer Control - Apply 20 pounds per acre by either ground equipment or by airplane. A second application may be required 7 to 10 days later. A third application may be required for control of second-brood borers. Consult State Extension Service specialists for most effective application dates.

CAUTION: Do not feed ensilage made from treated corn plants to dairy animals or animals being finished for slaughter. Do not allow dairy animals or animals being finished for slaughter to graze or feed on treated stover within 4 weeks of slaughter.

NOTICE. Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and agents the product under these conditions.

CAUTION


May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Can be absorbed through the skin. Wash thoroughly after using. Do not breathe dust. Avoid contamination of feed or foodstuffs. Do not use around dairy barns. Do not use on cucurbits.

This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of lakes, streams, and ponds. Apply only as specified on the label. Dispose of wastes by burying on noncrop lands away from water supplies. Dispose of containers by burning. Keep out of smoke.

ANTIDOTE

Internal - Give sodium chloride 15 g. (one tablespoon of table salt) in glass of warm water and repeat until vomit is clear. Call doctor immediately.

External - Wash with soap and water.



Warranted
REGISTERED AND APPROVED PRODUCT

Manufactured For

EPA Reg. No. 1029-73

