

# AIDEX

## 2% MALATHION

ACCEPTED

AUG 7 1974

UNDER FEDERAL PESTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 1029-59 SUBJECT  
 TO ATTACHED COMMENTS.

FOR BACKRUBBERS

Lice And Hornflies On Cattle  
Lice On Swine

FOR FOGGING AND SPRAYING  
 GRAIN STORAGE SURFACES

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
 See other panel for additional cautions

Active Ingredients:  
 Malathion\* 2.00  
 Xylene 1.00  
 Mineral Spirits 1.00  
 Kerosene 1.00  
 Inert Ingredients 1.00

\* Malathion is a cholinesterase inhibitor.

**DIRECTIONS**

FOR BACKRUBBERS

**CATTLE:** To Aid In Reduction Of Lice And Hornflies---A ready-to-use 2% Malathion solution. Do not dilute further with oil. For use in backrubber tank type machines. The amount of material needed will depend on the size of the reservoir. These devices should be made continuously accessible, one to each 35-45 head of cattle. Retreat the sides every 2-3 weeks or as necessary for control.

Do not treat animals under one month of age. Do not make accessible to dairy animals.

**SWINE:** To Aid In Reduction Of Lice---Dilute 1 gallon of 2% Malathion with 5 gallons of fuel oil or other suitable oil for use in tank type machines. The amount of material will depend on the size of the reservoir. These devices should be made continuously accessible. Retreat the sides every 2-3 weeks or as necessary for control.

Do not treat animals under one month of age.

FOR SPRAYING AND FOGGING

**GRAIN STORAGE SURFACES:** For control of confused flour beetle, rice weevil, granary weevil, saw-toothed grain beetle, flat grain beetle, lesser grain beetle, rusty grain beetle, lesser grain borer, Indian meal moth and other insects on grain storage surfaces. Thoroughly clean the surface. Apply Malathion at a point of high humidity. Allow treated surfaces to dry completely before use.

CATTLE

Harmful if swallowed. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and feed-stuffs.

Notice to Physician: Malathion contact may cause cholinesterase inhibition. Atropine is antidotal.

Highly toxic to fish. Do not contaminate any body of water by cleaning of equipment or disposal of wastes and containers or.

Do not reuse empty drum. Refrain from consuming any food or straw by perforating, crushing and burning it. Keep away from water supplies.

Malathion is a cholinesterase inhibitor. It may cause production on animals simultaneously or within a few days after treatment with an exposure to cholinesterase-inhibiting drugs, pesticides or chemicals. Consult a veterinarian at the first sign of an adverse reaction. Atropine is antidotal.

**NOTICE**

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME**

NET CONTENTS \_\_\_\_\_ GALLONS