

LIVESTOCK USES

DAIRY AND BEEF CATTLE, HOGS, HORSES, SHEEP, GOATS:

Control of Face Flies, Horn Flies and Stable Flies:

FOR WET SPRAYING: Dilute 1 quart of CIVAP with 3 1/2 gallons of water (equals 9 oz. per gallon). Spray all parts of the animal including legs with 1 to 2 pints of spray. For more effective Face Fly control add 1 pound of sugar per 14 gallons of finished spray. Repeat as necessary but not more often than once weekly.

FOR MIST SPRAYING: Dilute 1 quart of CIVAP with 1 3/4 gallons of water (equal to 1 pint per 7 pints). Apply as a fine mist at the rate of 2 oz. per animal. Cover all parts of the animal thoroughly. Do not exceed 2 oz. per animal per day.


FOR POUR-ON---To Control Horn Flies: Dilute 1 quart of CIVAP with 3 1/2 gallons of water (equal to 9 oz. per gallon). Apply by pouring 1 quart of the diluted emulsion along the backline of the animal. Repeat as necessary but not more often than once weekly.

FOR BACKRUBBER---To Control Horn Flies on Dairy and Beef Cattle: Dilute 1 quart of CIVAP with 4 1/2 gallons of fuel oil or other petroleum carrier, and charge backrubber device or sack or cloth as occasion requires. Retreat backrubber as needed. Do not spray animals with this oil solution.

Control of Lice and Ticks: Dilute 1 quart of CIVAP with 12 1/2 gallons of water and spray animal thoroughly. Make second application in 14 days if needed. Repeat as necessary but not more often than once weekly.

NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.



Ciodrin-DDVP Emulsible Concentrate
RESIDUAL DAIRY LIVESTOCK SPRAY
and
BACKRUBBER CONCENTRATE

For Use On

DAIRY AND BEEF CATTLE - HOGS - HORSES - SHEEP - GOATS

CONTROLS

FACE FLIES, HORN FLIES, STABLE FLIES, LICE AND TICKS

Active Ingredients: Dimethyl Phosphate of Alpha-Methyl-Benzyl 3-Hydroxy-cis-crotonate 15.00%; 2,2-Dichlorovinyl Dimethyl Phosphate 0.93%; related compounds 0.07%; Xylene 71.25%

Inert Ingredients: 12.75%

*T/M Shell Chemical Co.

WARNING: Keep Out Of Reach Of Children.
See side panel for other warnings.

BARN FLY CONTROL

As an aid in the reduction of House Flies, Face Flies and Stable Flies in dairy barns, animal shelters, hog barns and poultry houses, and other places where flies are a problem, mix 1 quart of CIVAP with 6 gallons (1 pint per 3 gallons) of water. Spray mixture on surfaces where flies alight and feed. Spray surface to run-off. To improve control, add 1 pound of sugar to 6 gallons of spray solution.

WARNING: Hazardous if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or clothing. Wash with soap and water after using. Avoid contaminating of feed, water, and foodstuffs. During commercial or prolonged exposure in spray mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type found suitable by the U.S. Bureau of Mines for protection against Ciodrin-DDVP mixtures. Keep away from heat and open flame.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard it in a safe place.

Do not use this material on Brahman or Brahman cross cattle.

ANTIDOTE

External---In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water, for eyes, get medical attention.

Internal----Call a physician immediately, give emetic of mustard. Large doses of Atropine are antidotal.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

**NET CONTENTS:
ONE GALLON**

MANUFACTURED BY



OMAHA, NEBRASKA

EPA Reg. No. 1029-95

ACCEPTED
JAN 18 1974
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR THE CONTROL OF PESTS IN THE UNITED STATES SUBJECT TO ATTACHED COMMENTS.

1029-95