

DIRECTIONS FOR FLY CONTROL

For control of houseflies in dairy barns, stables, hog pens, sheds, colic barns, battery houses and other farm buildings.

Residual Wall Spray: Mix one gallon of DIMEX 267 in 32 gallons of water. Apply with a mist blower, etc. Thoroughly wet all fly resting areas to the point of runoff. One gallon of spray will cover 200 sq. ft. of surface. Repeat applications should be made when necessary. Remove all animals from building when applying residual wall spray.

Spot Spray: Mix one pint of DIMEX 267 with water to make 3 1/2 gallons of spray. Use residual control spray with a knapsack or similar type sprayer in areas frequented by flies such as dairies, around windows, etc. Repeat applications should be made when necessary. Good sanitation is a necessary part of any effective fly control program.

Housefly Maggots: Mix 1 pint of DIMEX 267 with water to make 3 1/2 gallons of spray. Apply with a spray or with a sprinkling can in fly breeding areas, such as poultry droppings in cages, layer houses, garbage dumps and manure piles. Repeat application as additional manure or garbage is added.

General Outside Use: Mix one gallon of DIMEX 267 in 18 gallons of water. Thoroughly spray exposed surfaces such as walls, fences, garbage and refuse containers. Repeat applications should be made when necessary. Do not use in edible product areas of meat and food processing plants.

DIMEX-267 IS ACCEPTABLE FOR USE AS A RESIDUAL-TYPE INSECTICIDE IN AND AROUND FEDERALLY INSPECTED MEAT PACKING PLANTS EXCEPT IN AREAS WHERE MEAT FOOD PRODUCTS ARE PROCESSED OR HANDLED. Do not use in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

DIMEX 267

Systemic Insecticide

Contains 2.67 lbs. Dimethoate Per Gallon
In Water Miscible Form.

Active Ingredient:
Dimethoate (O, O-dimethyl S- [(methylcarbamoyl) methyl] phosphorothioate) **30.5%**

Inert Ingredients: **69.5%**

Total 100.0%

WARNING: Keep Out Of Reach Of Children.
See side panel for antidote and other warnings.

NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

DO NOT STORE BELOW FREEZING TEMPERATURES

DIRECTIONS

This product is intended for use by the commercial grower or producer in accordance with the label instructions for each crop. It is not intended for use in household or other non-commercial situations. Do not use in areas where food is prepared or processed. Do not use in areas where food is stored. Do not use in areas where food is sold. Do not use in areas where food is served. Do not use in areas where food is displayed. Do not use in areas where food is consumed. Do not use in areas where food is stored. Do not use in areas where food is sold. Do not use in areas where food is served. Do not use in areas where food is displayed. Do not use in areas where food is consumed.

CROP	PEST CONTROLLED	RATE	INTERVAL DAYS BETWEEN LAST APPLICATION AND HARVEST
FRUIT			
Apple	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Orange	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
VEGETABLE CROPS			
Asparagus	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Brussels Sprouts	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Broccoli	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Carrots	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Cauliflower	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Chard	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Chicory	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Endive	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Kale	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Kohlrabi	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Lettuces	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Spinach	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Swiss Chard	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Turnips	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
MILLS			
Wheat	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
ORCHARD			
Apple	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Orange	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Peach	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Pineapple	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14

WARNING

Vapor harmful. May be fatal if swallowed, inhaled or absorbed through the skin. Cause eye irritation. Avoid breathing vapor or spray mist. Use only with adequate ventilation. Do not get in eyes. If in contact with eyes, wash thoroughly after handling and flush with plenty of water. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

FIRST AID: If swallowed, induce vomiting immediately and get medical attention. In case of contact with eyes, flush eyes with water for at least 15 minutes. Use medical attention.

Note: This product may cause cholinesterase inhibition. Atropine is antidote.

Preventive measure: 2-PAM, PROTOPAM or diisopropyl fluorophosphate may be effective as an adjunct to atropine. Use according to label directions.

Do not reuse empty container. Destroy by perforating or crushing, and burning in a safe place away from water supplies. Large drums may be returned to the drum conditioner.

This product is not to be fed to birds. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment into or about it. This product is toxic to bees. Restrict application to the period after dark when bees are inside the hive or in the early morning before the bees are foraging in the fields. Protective information may be obtained from your cooperative agricultural extension service.

Not for use on animals. Do not contaminate feed and foodstuffs, drinking fountains, later or feed troughs. Do not use in milk processing rooms, milk houses or milk storage rooms. Do not use in the home.

The Buyer of this material (including its ultimate user) is granted a license under U.S. Patent No. 3,210,242, limited to the use of the within contents, a prepaid royalty being included in the selling price. A non-transferable license under such patent may be obtained from the patent owner on current royalty terms whatever the source of Dimethoate used.

CROP	PEST CONTROLLED	RATE	INTERVAL DAYS BETWEEN LAST APPLICATION AND HARVEST
Apple	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Orange	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Peach	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Pineapple	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
FRUIT			
Apple	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Orange	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Peach	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Pineapple	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
VEGETABLE CROPS			
Asparagus	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Brussels Sprouts	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Broccoli	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Carrots	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Cauliflower	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Chard	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Chicory	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Endive	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Kale	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Kohlrabi	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Lettuces	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Spinach	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Swiss Chard	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Turnips	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
MILLS			
Wheat	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
ORCHARD			
Apple	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Orange	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Peach	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14
Pineapple	Apply at 20-30% bloom	1/2 to 3/4 gal per acre	14



1029-108
ACCEPTED
JUN 4 1973
BY THE FEDERAL INSECTICIDE
OFFICE OF THE FEDERAL AGRI-
CULTURE
1029-108

DIRECTIONS FOR FLY CONTROL

Use in control of houseflies in dairy barns, stables, hog pens, sheds, corrals, pig-breeding pens and other farm buildings.

Residual Wall Spray. Mix one gallon of DIMEX-267 in 20 gallons of water. Spray on walls, window sills, etc. This spray will kill flies on contact. The point of contact should be sprayed with water. Repeat application if necessary. Repeat application if flies are still present when the spray has dried. Repeat application if necessary.

Spot Spray. Mix one pint of DIMEX-267 with water to make 5 gallons. Spray on areas where flies are most abundant. This includes racks in dairies, areas frequented by flies, and areas around windows. Repeat application if necessary. Repeat application if flies are still present when the spray has dried.

Housefly Maggots. Mix one pint of DIMEX-267 with water to make 5 gallons. Apply to areas where maggots are most abundant. This includes areas around manure piles, garbage dumps, and manure piles. Repeat application if maggots are still present when the spray has dried.

General Outside Use: Mix one gallon of DIMEX-267 in 18 gallons of water. This spray is applied to exterior surfaces such as walls, fences, garbage and refuse containers. Repeat applications if necessary. Do not use in edible product areas of meat and food processing plants.

DIMEX-267 IS ACCEPTABLE FOR USE AS A RESIDUAL-TYPE INSECTICIDE IN AND AROUND FEDERALLY INSPECTED MEAT PACKING PLANTS EXCEPT IN AREAS WHERE MEAT FOOD PRODUCTS ARE PROCESSED OR HANDLED. Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

DO NOT STORE BELOW FREEZING TEMPERATURES

DIMEX

Systemic

Contains 2.67 'lbs. 2
In Water M

Active Ingredient:

Dimethoate (O, O-dimethyl S-methyl] phosph

Inert ingredients:

WARNING: Keep Out Of
See side panel

NO

Seller's guarantee shall be limited
ject thereto the buyer assumes any
of use or handling and accepts the

DO NOT USE, POUR, SPILL OR ST

DIRECTIONS

For use in control of houseflies in dairy barns, stables, hog pens, sheds, corrals, pig-breeding pens and other farm buildings.

CROP	PESTS CONTROLLED	RATE	INTERVAL DAYS BETWEEN LAST APPLICATION AND HARVEST
FRUIT			
Apples	Apple maggot, codling moth, European spruce sawfly, fruitworm, etc.	1.0-2.0 gal. per acre	14-21 days
Pears	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Grapefruit	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Lemons	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Oranges	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Tangerines	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Citrus and bearing			
Citrus and bearing	Same as Apples	1.0-2.0 gal. per acre	14-21 days
NUTS			
Pecans	Same as Apples	1.0-2.0 gal. per acre	14-21 days
VEGETABLE CROPS			
Beans green lima	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Beans dry	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Broccoli	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Cauliflower	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Cabbage	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Head lettuce	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Leaf lettuce	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Spinach	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Catnip	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Rice	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Turnip greens and roots	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Mustard greens	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Sweet chard	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Endive (chicory)	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Melons (except watermelons)	Same as Apples	1.0-2.0 gal. per acre	14-21 days
Watermelons	Same as Apples	1.0-2.0 gal. per acre	14-21 days



1024-103

ACCEPTED