

DIRECTIONS FOR USE

FLY BAN PREMIUM is a ready to use, highly effective product that is artificially colored to prevent accidental usage. No mixing preparation or application equipment is required to use the material as a dry bait. Scatter the bait evenly on floors or other horizontal surfaces where flies alight or congregate in factories, picnic areas, patios, kennels, dairy barns, calf barns, hog barns, poultry houses, livestock loafing sheds, kennels, garbage areas, outdoor latrines and dumps. Do NOT use in edible products areas of food processing plants. Good sanitation measures and proper disposal of wastes are necessary for effective fly control.

APPLICATION RATES

Apply to floors or other areas mentioned above at the rate of 8 oz. per 1000 sq. feet. Apply evenly over manure piles and floor areas where flies breed. Applications should be made daily until the fly population is under control. Then applications should be repeated every few days as necessary. Baits should be placed in areas not accessible to children, pets, wildlife and domestic animals.

FOR MAGGOT CONTROL

Apply FLY BAN PREMIUM at a rate of 1 lb. per 2000 sq. ft. of fly breeding area, garbage dumps and manure piles. Repeated applications will be

FLY-BAN PREMIUM Fly Killer

READY-TO-USE DRY BAIT
Also for mixing as a liquid bait

CAUTION
KEEP OUT OF REACH OF CHILDREN
See other panel for additional cautions

ACTIVE INGREDIENTS:	By Weight
2,2-Dichlorovinyl dimethyl phosphate (DDVP)	0.465%
Related compounds	0.035%
Ronnel (O, O-dimethyl 0-2,4,5- trichlorophenyl phosphorothioate)	0.250%
INERT INGREDIENTS:	99.250%
Total	100.000%

STORE IN A COOL DRY PLACE
Close Container Tightly After Each Use

CONTAINS
DDVP and RONNEL

NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk of handling and accepts the product on these conditions.

Net Weight 100 Pounds

DIRECTIONS (cont'd)

necessary as additional garbage or manure is added.

AS LIQUID BAIT

Use 4 oz. of FLY BAN PREMIUM per gallon of water and sprinkle on floors of above mentioned buildings. Do not exceed a rate of 1 gallon per 1000 sq. ft. Application of the liquid bait should be made daily until flies are under control and then 2 to 3 times per week as required.

CAUTION

Avoid contact with skin. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Harmful if swallowed. Do not apply the bait in large amounts at any one place and it will be impossible for domestic animals to lick up the bait. Do not contaminate livestock feed. Do not use in human dwellings or food processing plants. Do not contaminate milk or milking utensils.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy when empty.

ACCEPTED
AUG 26 1975
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 1029-117

EPA Est.
1029-1A-1

MANUFACTURED BY

AIDEX

OMAHA, NEBRASKA

EPA
Reg. No.
1029-117

CC (6/26/75) 2C