

1029-123

6-3-1975

LEISUR-AID[®]

44215-45

LAWN INSECT CONTROL

CONTAINING DURSBAN*

19713-341

Controls Chinch Bugs, Sod Webworms (Lawn Moths), Ants, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Sowbugs (Pillbugs) Certain White Grubs and Hyperodes (Turfgrass) Weevils Infesting Home Lawns and Other Ornamental Turf Areas

ACTIVE INGREDIENT:
Chlorpyrifos | O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate | 0.5%
INERT INGREDIENTS: 99.5%

GRANULAR INSECTICIDE

CAUTION

KEEP OUT OF REACH OF CHILDREN
MAY BE HARMFUL IF SWALLOWED

DIRECTIONS FOR USE

Apply LAWN INSECT CONTROL granular insecticide uniformly to home lawn and other ornamental and recreational turf grass areas using any hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear using LAWN INSECT CONTROL at the rate of 200 pounds per acre. To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle. Immediately after application sprinkle the turf with 300 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine amounts of LAWN INSECT CONTROL and water needed for smaller sized areas. Re-treat as necessary.

RATES TO USE

| Area to be Treated | Pounds of LAWN INSECT CONTROL | Gallons of Water to Sprinkle on Turf after Treatment |
|--------------------|-------------------------------|------------------------------------------------------|
| 1000 sq. ft. | 5 | 30 |
| 5000 sq. ft. | 25 | 150 |
| 10700 sq. ft. | 50 | 325 |

*To control certain white grubs (larvae of European chafer and Japanese beetle) use two to four times the amount of LAWN INSECT CONTROL shown in the table, i.e. 10 to 20 pounds per 1,000 sq. ft. Apply the granules uniformly over the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

*To control Hyperodes (turfgrass) weevils use twice the rate of LAWN INSECT CONTROL shown in the table. Treat suspected problem areas in mid-April and again in mid-May. Consult your local agricultural Extension Service specialist for best time to treat in your area.

ACCEPTED
JUN 3 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered under EPA Reg. No. 1029-123

Transferred to
44215-45
on 1/29/82

USE PRECAUTIONS

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

CAUTION

Do not take internally. Avoid eye contact. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before re-use. Keep away from food, feedstuffs and domestic water supplies. Do not induce vomiting if swallowed. Call a physician immediately. NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label. Do not apply when weather conditions favor drift from areas treated.

CONTAINER DISPOSAL

Dispose of empty container and wastes by burning or burying in non-croplands away from water supplies.

NET CONTENTS
50
POUNDS

NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

*DURSBAN Trademark of
THE DOW CHEMICAL COMPANY
U.S. Patent No. 3,244,586

(1-28/75)

EPA Est.
1029-1A-1

MANUFACTURED BY



OMAHA, NEBRASKA

EPA
Reg. No.
1029-123