

pm 32

1043-19

page 171

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, promptly drink a large quantity of water. Avoid alcohol. Do not induce vomiting. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. General and Tuberculous Disinfection, Elimination of Fungi and Viruses*; Spray hard, non-porous environmental surfaces from any angle, even upside down, from a distance of 6-8 inches for 2-3 seconds or until surface is thoroughly wet. Allow surfaces to remain wet for 10 minutes or allow to air dry without wiping or rinsing. Gross filth or heavy soils must be removed before disinfection.

KILLS HIV-1 (AIDS VIRUS) ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surface/object likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (HUMAN IMMUNODEFICIENCY VIRUS OR AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:
Personal Protection: Wear appropriate barrier protection such as latex gloves, gowns, masks, or eye coverings. Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application. Apply solution as directed in paragraph above. Contact Time: While the HIV-1 virus is inactivated in 1 minute, use a 10 minute contact time for disinfection of all organisms on this label.

Infectious Material Disposal: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious disposal.

To Deodorize, and Refreshen the Air: Direct Spray upward toward the center of the room for 3-5 seconds.

Staphene®
DISINFECTANT SPRAY
AND DEODORIZER

ACTIVE INGREDIENTS

o-benzyl-p-chlorophenol	0.077%
p-tertiary amyphenol	0.074%
o-phenylphenol	0.041%
N-alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethylbenzyl ammonium chlorides	0.042%
N-alkyl (30% C ₁₂ , 30% C ₁₄ , 17% C ₁₆ , 3% C ₁₈) dimethyl ethylbenzyl ammonium chlorides	0.042%
ethanol	53.096%
INERT INGREDIENTS	46.628%

E.P.A. Reg. No. 1043-19

Germicidal	Tuberculoicidal
Air Deodorizer	Fungicidal
Virucidal*	

SPRAY FROM ANY ANGLE, EVEN UPSIDE DOWN

KEEP OUT OF REACH OF CHILDREN

CAUTION

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperature above 130°F may cause bursting. Never throw container into incinerator.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Do not use on surfaces which will come in prolonged contact with skin. Avoid prolonged inhalation of spray and mist and spraying of food, skin or varnished surfaces.

See side panel for additional precautionary and practical treatment statements.

Net Contents 16 oz. (1 lb.) (453g)
Can Size 20 fl. oz.

Staphene®
DISINFECTANT SPRAY
AND DEODORIZER
Germicidal* Tuberculoicidal
Air Deodorizer* Fungicidal* Virucidal*

Staphene Disinfectant Spray and Air Deodorizer is an aerosol-dispersed, spot disinfectant and deodorant for schools, medical, dental and business offices, homes, sick rooms, nursing homes and general hospital use.

- *Bedside and office furniture
- *Waste cans and laundry hampers
- *Trash carts
- *Telephones
- *Sinks

Germicidal: Passes A.O.A.C. Germicidal Spray Products Test (Staphylococcus aureus, Salmonella choleraesuis, Pseudomonas aeruginosa) in the presence of 5% blood serum, 10 minutes at 20°C.

Tuberculoicidal: Passes A.O.A.C. Germicidal Spray Products Test (Mycobacterium tuberculosis var. bovis) in the presence of 5% blood serum, 10 minutes at room temperature (25°C/77°F) on hard inanimate surfaces. Gross filth or heavy soils must be removed before disinfection.

Fungicidal: Passes A.O.A.C. Germicidal Spray Products Test (Trichophyton mantagrophytes) in the presence of 5% blood serum, 10 minutes at 20°C.

Virucidal: Passes virucidal assay (EPA Proposed Method) [Influenza A, Herpes simplex Type 2, Rhinovirus Type 1A] in the presence of 5% blood serum, 10 minutes at 20°C on hard inanimate surfaces.


When tested by an EPA-approved Dilution Method, the HIV-1 (AIDS) virus, with added 10% organic soil (serum), was completely inactivated in 60 seconds at 20-25°C. Although efficacy at 1 minute contact time has been shown to be adequate against HIV-1, this time would not be sufficient for other organisms. Use a 10-minute contact time for disinfection against all of the organisms claimed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not use or store near heat or open flame. Open dumping is prohibited. Do not reuse empty container. Wrap container and put in trash.

ConvaTec
Division of E.R. Squibb & Sons, Inc.
St. Louis, Mo 63110

EPA Est. No. 33915-MI-1



(Logo) ConvaTec
A Bristol Myers-Squibb Company

MAR 25 3 05 PM '97
RECD ENV/DPD1