

EPA REGISTRATION NO. 1043-26-AA

1 STROKE ENVIRON combines the high detergent action of a general purpose cleaner with the non-selective germicidal action of a general purpose disinfectant. It effectively cleans, disinfects and deodorizes food, animal and poultry installations.

GERMICIDAL QUALIFICATION (AOAC Use Dilution Confirmation).....1:256
 TUBERCULOCIDAL ACTIVITY TEST (AOAC, M. tuberculosis).....1:256
 FUNGICIDAL TEST (AOAC, T. interdigitale).....1:256
 GERMICIDAL, PSEUDOMONACIDAL, TUBERCULOCIDAL, AND FUNGICIDAL BY OFFICIAL AOAC QUALIFICATION IN SOFT OR HARD WATER (AOAC 400 PPM as CaCO₃)

ACTIVE INGREDIENTS: o-phenylphenol 10.0%, o-benzyl-p-chlorophenol 8.5%, p-tertiary-amylphenol 2.0%.

TOTAL INERT INGREDIENTS79.5%

DIRECTIONS FOR USE—FOR ALL GENERAL CLEANING AND DISINFECTION, INCLUDING TUBERCULOUS AREAS, USE 1:256 SOLUTION (½ OZ. PER GALLON OF WATER). For simplicity and accurate solution control, use this concentrate with Vestal's One-Stroke Dispenser. Use 1 dispenser stroke (up and down) for each gallon of solution prepared (each stroke delivers ½ oz.). When used as directed, 1 STROKE ENVIRON is harmless to fabrics, resilient flooring, tile, terrazzo, painted and varnished surfaces. When cleaning and disinfecting food, animal or poultry facilities, first remove all livestock from the premises, cover or remove all feed and foodstuffs, and remove all dirt, litter, and debris. Then thoroughly saturate all surfaces with 1 STROKE ENVIRON use solution. (Rinse all surfaces contacting food, and flush feeders and waterers with potable water after disinfecting.) Allow disinfected surfaces to dry before rehousing livestock.

Read carefully the following statement which applies to the contents of this container.

DANGER: Corrosive to tissues. This concentrate causes eye and skin damage. Harmful or fatal if swallowed. Do not get in eyes, on skin or on clothing. Wear goggles and rubber gloves when handling. Avoid contamination of food. Do not reuse empty container.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, do not induce vomiting, drink large quantities of fluid and call a physician immediately.

VESTAL LABORATORIES • DIVISION OF CHEMED CORPORATION • ST. LOUIS, MO. 63110



ria and
ubercu-
viruses.

LDREN
cautions

ALLON)