

ACCEPTED  
MAY 30 1991  
This product is registered under the Fungicide, Fungicide, and  
Rodenticide Act, as amended, for the  
purpose registered under  
EPA Reg. No. 1043-26

Center Panel

# 1STROKE LENVIRON®

U.S. PATENT NO. 3,538,217

EPA REGISTRATION NO. 1043-26

Germicide • Fungicide • Pseudomonasicide  
Tuberculocide • Virucide • Deodorant

ACTIVE INGREDIENTS:	
sodium o-phenvionate .....	11.3%
sodium o-benzyl-o-chlorophenolate .....	9.4%
sodium o-tertiary-amyphenate .....	2.3%
INERT INGREDIENTS .....	77.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT: (FIRST AID): In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, do not induce vomiting. Drink promptly a large quantity of water. Avoid alcohol. Call physician immediately. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory respiratory depression, and convulsion may be needed.

NET CONTENTS: 1 U.S. GALLON (3.78 Liters)

### 1-Stroke Environ

1-Stroke Environ is a detergent based germicide developed especially to control cross contamination of treated surfaces in the agricultural and food industries, and by the veterinary profession. It is intended for use on precleaned hard non-porous surfaces including floors, walls, counters, equipment in food processing areas, feed racks, troughs, and automatic feeders. It may be used in food processing, preparation and service facilities, in animal and avian husbandry, poultry and feed processing facilities, veterinary clinics, and animal research facilities.

When used as directed, this product is effective against pathogenic bacteria such as *S. aureus*, *S. choleraesuis*, *Ps. aeruginosa*, *M. tuberculosis* and fungi (T. mentagrophytes) when tested at 1:256 by official A.O.A.C. methods in either hard (400ppm) or soft water; and \*viruses (Influenza A<sub>2</sub>, Herpes simplex, Type 2 Adenovirus, Vaccinia, Newcastle disease, Infectious<sup>2</sup> laryngotracheitis, Infectious bronchitis, Marek's Agent (JM-V), Hog Cholera) on inanimate environmental surfaces in 10 minutes at 20°C.

When tested according to A.O.A.C. Use Dilution Testing against *Salmonella enteritidis*, *Pasteurella multocida*, *Bordetella avium* and *Mycoplasma gallisepticum*, 1-Stroke Environ was shown to be effective when diluted 1:256 (1/2 oz. per gallon) in 400ppm A.O.A.C. hard water and 5% organic soil (serum), in 10 minutes at 20°C.

GERMICIDAL QUALIFICATION (A.O.A.C. Use Dilution Confirmation), 10 minutes at 20°C when diluted 1:256 in (400ppm CaCO<sub>3</sub>) A.O.A. C. hard water.

Tuberculocidal Activity Test: A.O.A.C. *M. tuberculosis* (BCG): effective in 10 minutes at 20°C at 1:256 in soft or hard water (400ppm as CaCO<sub>3</sub>). Remove heavy soil and preclean surface before application of product.

Fungicidal Test (A.O.A.C. T. Mentagrophytes), 10 minutes at 20°C when diluted 1:256 in (400ppm CaCO<sub>3</sub>) A.O.A.C. hard water.

GERMICIDAL, PSEUDOMONACIDAL, TUBERCULOCIDAL AND FUNGICIDAL BY OFFICIAL A.O.A.C. QUALIFICATION IN SOFT OR HARD WATER (A.O.A.C. 400PPM AS CaCO<sub>3</sub>) IN TEN MINUTES AT 20°C.

This product is recommended for use on washable hard surfaces such as counters, equipment, resilient tile flooring, terrazzo and painted or varnished surfaces.

1-Stroke Environ helps in the control of cross contamination and odors of bacterial origin.

Primary skin and eye irritation scores of 1:256 dilutions -- 0 when tested according to 16 CFR 1500.41 and 1500.42.

DIVISION OF CALGON CORPORATION  
SUBSIDIARY OF MERCK & CO., INC.  
ST. LOUIS, MO 63133

5397-06B



1-Stroke Environ--Left Panel

Precautionary Statements

Hazard to Humans and Domestic Animals:

DANGER

Read carefully the following statement which applies to the contents of this container: Corrosive to tissue. Causes eye and skin damage. Harmful or fatal if swallowed. Do not get in eyes, on skin, or on clothing. Wear goggles and rubber gloves when handling. Avoid contamination of food.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Pesticide Disposal: This germicide, its solutions or rinsings from empty containers should be disposed of in a toilet or service sink served by a sanitary sewer or in a land fill approved for pesticides.

Container Disposal: Triple rinse and deposit in a waste container for incineration or burial in a land fill approved for pesticide containers.

General: Consult Federal, State, or Local disposal authorities for approved alternative procedures such as limited open burning, if frozen, thaw, and remix before use.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For use on all washable, precleaned, hard environmental surfaces including tuberculous areas, use a 1:256 (1/2 ounce/gallon of water). For accurate solution preparation, use Calgon Vestal's 1-Stroke Dispenser (one stroke delivers 1/2 oz. of product). This is a complete product. Do not add other chemicals. Use only as directed.

CLEANING AND DISINFECTING ANIMAL AND POULTRY FACILITIES: Do not use in milking stall, milking parlors, or milk houses. Cover or remove all food, feedstuffs, packaging material, and animals. Remove all dirt, litter, and debris from surfaces to be disinfected. Thoroughly clean all surfaces with soap or detergent, and rinse with water. Apply use solution with a sponge, mop, or coarse spray to thoroughly wet all surfaces. Allow the surface to remain wet for 10 minutes and remove excess solution with a wrung-out applicator. Discard solutions when they become soiled and replace with a fresh solution of this product. Wash all feeders and waterers with a detergent, and rinse with portable water before reuse. Ventilate buildings, coops, and other closed spaces. Allow disinfected areas to dry before rehousing livestock. Always remove gross filth and heavy soil before application of the solutions of this product.

(Left Panel cont.)

**CLEANING AND DISINFECTING FOOD CONTACT SURFACES, TOOLS, AND EQUIPMENT:** Cover or remove all food and packaging materials. Thoroughly clean surfaces to be disinfected before application of the solution of this product. Apply use solution to the surface being disinfected with a sponge, mop, cloth or coarse spray and allow to remain wet for 10 minutes. Discard soiled solutions and replace with fresh use solutions. Rinse all surfaces with a potable water rinse after disinfection, and allow to air dry before reuse.

**FOGGING AS AN ADJUNCT TO REGULAR CLEANING AND DISINFECTING:**  
This product may be used in fogging (wet misting) as an adjunct either proceeding or following regular cleaning and disinfecting procedures in poultry operations, hatcheries, swine operations, animal care facilities, animal research facilities, livestock operations, veterinary facilities, zoos and kennels. Use a 1:256 dilution (1/2 ounce per gallon of water). Do not use in milking stalls, parlors or houses. Remove all animals and feed from areas to be fogged. Remove all litter and manure from surfaces or areas traversed by animals. Empty all trough, racks and other feeding and watering applicances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Fog the desired area at 32 to 64 ounces of use dilution per 1000 cubic feet using equipment with an automatic timer. Do not remain in treated areas; allow at least two hours after fogging is complete before reentering fogged area. Before fogging remove or cover any food or packaging material with waterproof coverings. Treated surfaces must be thoroughly scrubbed with 1-Stroke Environ at a 1:256 dilution and rinsed with potable water before reuse. Do not rehouse livestock or employ equipment until treatment has dried.

**MUSHROOM FARM USE:** A normal use-solution (1/2 ounce per gallon of water) may be used between crops on surfaces that do not come in contact with the crop. Use on tanks, air conditioners, air ducts, fans, and breezeways, compost wharf, concreted surfaces, inside and outside walls of mushroom houses, lofts, floors, spawning machines, tampers, casing rings, storage sheds and track alleys before spawning. Thoroughly clean all surfaces with soap or detergent and rinse with water. Then apply use-solution with cloth, sponge, mop or coarse spray to thoroughly wet all surfaces; allow to remain wet for 10 minutes. Apply ONLY to surfaces that do not come in contact with the crop. Do not apply to the crop, compost or casings.