

Active ingredients: Potassium o-benzyl-p chlorophenate.  
5.57%; potassium p-tertiary amyphenate. 5.52%; potassium  
o-phenyphenate. 5.51%.  
Inert ingredients.....83.4%

EPA REGISTRATION NO 1043-32

**DISINFECTS • DEODORIZES**  
**KEEP OUT OF REACH OF CHILDREN**  
**DANGER**

**Statement of Practical Treatment**

In case of eye contact, flush with water for 15 minutes. Get medical attention.  
For skin contact, flush thoroughly with water. See side panel for additional  
precautionary statements, and first aid.

**CONTENTS: ONE U.S. GALLON (3.78 Liters)**

NOV 06 1968

1043-32

This product is a concentrated, phosphate-free, multi-purpose germicide. It is intended for use on floors, walls, equipment, sickroom receptacles, hard rubber goods and instruments. When used as directed, it is harmless to resilient flooring, tile, terrazzo, painted and varnished surfaces.

A 1/2% use-solution is effective against the following pathogenic bacteria and fungi including *S. aureus*, *S. choleraesuis*, *Ps. aeruginosa*, *M. tuberculosis*, *T. mentagrophytes* and viruses such as Influenza A<sub>2</sub>, Herpes Simplex, Type 2 adenovirus and Vaccinia in 10 minutes at 20°C on hard, non-porous environmental surfaces.

GERMICIDAL QUALIFICATION: A.O.A.C. Use-Dilution Confirmation, Effective in 10 minutes at 20°C when diluted at 1:200.

TUBERCULOCIDAL ACTIVITY TEST: A.O.A.C., *M. tuberculosis*, Effective in 10 minutes at 20°C when diluted at 1:200. Remove heavy soil and gross filth and thoroughly clean surfaces.

FUNGICIDAL TEST: A.O.A.C., *T. mentagrophytes*, Effective in 10 minutes at 20°C when diluted at 1:200.

#### DILUTION TABLE

Quantity of Staphene Concentrate to make 1/2% Use Dilutions		
Water	1 Gallon	5 Gallons
CONCENTRATE	2/3 fl. oz.	3 1/3 fl. oz.

Classified by Underwriter's Laboratories, Inc. as to electrical conductivity when used on conductive floors and spontaneous heating. Staphene—For use with listed electrically conductive flooring of the vinyl type.

942 P

6388-06 A




DIVISION OF CALGON CORPORATION  
SUBSIDIARY OF MERCK & CO., INC

St. Louis, Missouri 63133

EPA EST. N. 143 MO 1

RIGHT PANEL



do not induce vomiting, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available drink large quantities of water and call a physician immediately. **Note to Physician:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Disinfecting Washable, Precleaned Hard Nonporous Surfaces:** Use 1/2% solution, 2/3 ounce per gallon of water for disinfection, including tuberculous areas. Apply by wiping, mopping or scrubbing to wet surfaces thoroughly with the use solution. Treated surfaces should remain wet for ten minutes. Remove excess solution with a wrung-out applicator. Discard solution when it becomes dirty, and replace with a fresh solution. Always remove gross filth or heavy soil and preclean surface prior to applying the solutions of this product. Heavily soiled surfaces should be precleaned and rinsed before applying a Staphene solution.

**Instrument Disinfection:** Pre-clean instruments thoroughly, removing all soils including blood and serous exudates using a 1%-2% STAPHENE solution (1/2 - 1/3 oz./gallon). Rinse and rough dry objects before immersion in solution. Clean and rinse the lumens of hollow instruments before filling with solution. Then soak in a fresh 1/2% STAPHENE solution for 15 minutes.

**Terrazzo Care:** Use at 1:128 (1 ounce per gallon of water) to increase patina and highlight the natural beauty of terrazzo floors. This is a complete product. Use only as directed. Do not add other chemicals.


## STORAGE AND DISPOSAL

**Prohibitions:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

**Pesticide Disposal:** This germicide, its solutions or rinsings from empty containers should be disposed of in a toilet or service sink served by a sanitary sewer, or in a landfill approved for pesticides.

**Container Disposal:** Triple rinse and deposit in a waste container for incineration or burial in a landfill approved for pesticide containers.

**General:** Consult Federal, State or Local disposal authorities for approved alternative procedures such as limited open burning. If frozen, thaw and remix before use.



PM 32 1043-32

1/3

CENTER  
PANEL

CONCENTRATED

# Staphene.

HOSPITAL GERMICIDE

Active ingredients: Potassium o-benzyl-p-chlorophenate, 5.57%; potassium o-tertiary amyphenate, 5.52%; potassium o-phenyphenate, 5.51%.  
 Inert ingredients.....83.4%

EPA REGISTRATION NO. 1043-32

**DISINFECTS • DEODORIZES**

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**

Statement of Practical Treatment


In case of eye contact, flush with water for 15 minutes. Get medical attention.  
 For skin contact, flush thoroughly with water. See side panel for additional  
 precautionary statements, and first aid.

**CONTENTS: ONE U.S. GALLON (3.78 Liters)**

ACCEPTED

NOV 06 1989

U.S. Environmental Protection Agency  
 Office of Pesticide Programs  
 Washington, D.C. 20460  
 EPA Reg. No. 1043-32



This product is a concentrated, phosphate-free, multi-purpose germicide. It is intended for use on floors, walls, equipment, sickroom receptacles, hard rubber goods and instruments. When used as directed, it is harmless to resilient flooring, tile, terrazzo, painted and varnished surfaces.

A 1/2% use-solution is effective against the following pathogenic bacteria and fungi including *S. aureus*, *S. choleraesuis*, *Ps. aeruginosa*, *M. tuberculosis*, *T. mentagrophytes* and viruses such as Influenza A<sub>2</sub>, Herpes Simplex, Type 2 adenovirus and Vaccinia in 10 minutes at 20°C on hard, non-porous environmental surfaces.

GERMICIDAL QUALIFICATION: A.O.A.C. Use-Dilution Confirmation, Effective in 10 minutes at 20°C when diluted at 1:200.

TUBERCULOCIDAL ACTIVITY TEST: A.O.A.C., *M. tuberculosis*, Effective in 10 minutes at 20°C when diluted at 1:200. Remove heavy soil and gross filth and thoroughly clean surfaces.

FUNGICIDAL TEST: A.O.A.C., *T. mentagrophytes*, Effective in 10 minutes at 20°C when diluted at 1:200.

**DILUTION TABLE**

Quantity of Staphene Concentrate to make 1/2% Use Dilutions		
Water	1 Gallon	5 Gallons
CONCENTRATE	2/3 fl. oz.	3 1/3 fl. oz.

Classified by Underwriter's Laboratories, Inc. as to electrical conductivity when used on conductive floors and spontaneous heating. Staphene—For use with listed electrically conductive flooring of the vinyl type. 942 P.

6388-06 A



DIVISION OF CALGON CORPORATION  
SUBSIDIARY OF MERCK & CO., INC.  
St. Louis, Missouri 63133

EPA Est. No. 1043-MO-1

RIGHT PANEL

**ACCEPTED**  
NOV 6 1993  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1043-32

**FIRST AID:** In case of contact with eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, do not induce vomiting, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available drink large quantities of water and call a physician immediately. Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Disinfecting Washable, Precleaned Hard Nonporous Surfaces:** Use 1/2% solution, 2/3 ounce per gallon of water for disinfection, including tuberculous areas. Apply by wiping, mopping or scrubbing to wet surfaces thoroughly with the use solution. Treated surfaces should remain wet for ten minutes. Remove excess solution with a wrung-out applicator. Discard solution when it becomes dirty, and replace with a fresh solution. Always remove gross filth or heavy soil and preclean surface prior to applying the solutions of this product. Heavily soiled surfaces should be precleaned and rinsed before applying a Staphene solution.

**Cold Instrument Disinfection:** Pre-clean instruments throughly, removing all soils including blood and serous exudates using a 1%-2% STAPHENE solution (1 1/4 - 2 1/2 oz./gallon). Rinse and rough dry objects before immersion in solution. Clean and rinse the lumens of hollow instruments before filling with solution. Then soak in a fresh 1/2% STAPHENE solution for 15 minutes.

**Terrazzo Care:** Use at 1:128 (1 ounce per gallon of water) to increase patina and highlight the natural beauty of terrazzo floors. This is a complete product. Use only as directed. Do not add other chemicals.

## STORAGE AND DISPOSAL

**Prohibitions:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

**Pesticide Disposal:** This germicide, its solutions or rinsings from empty containers should be disposed of in a toilet or service sink served by a sanitary sewer, or in a landfill approved for pesticides.

**Container Disposal:** Triple rinse and deposit in a waste container for incineration or burial in a landfill approved for pesticide containers.

**General:** Consult Federal, State or Local disposal authorities for approved alternative procedures such as limited open burning. If frozen, thaw and remix before use.

